

Forecasting range shifts of a dioecious plant species under climate change

Jacob K. Moutouama 0000-0003-1599-1671^{a,c,1}, Aldo Compagnoni 0000-0001-8302-7492^b, and Tom E.X. Miller^a

^aProgram in Ecology and Evolutionary Biology, Department of BioSciences, Rice University, Houston, Texas, USA; ^bInstitute of Biology, Martin Luther University Halle-Wittenberg, Halle, Germany; and German Centre for Integrative Biodiversity Research (iDiv), Leipzig, Germany

Global climate change has triggered an urgent need for predicting the reorganization of Earth's biodiversity. For dioecious species (in which female and male reproductive organs are not on the same individual), it is unclear how commonly unique climate sensitivities of females and males could influence projections for species-level responses to climate change. We developed demographic models of range limitation, parameterized from geographically distributed common garden experiments, with females and males of a dioecious grass species (*Poa arachnifera*) throughout and beyond its range in the south-central U.S. Female-dominant and two-sex model versions both predict that future climate change will alter population viability and will induce a poleward niche shift beyond current northern limits. However, the magnitude of niche shift was underestimated by the female-dominant model, because females have broader temperature tolerance than males and become mate-limited under female-biased sex ratios. Our result illustrate how explicit accounting for both sexes could enhance population viability forecasts and conservation planning for dioecious species in response to climate change.

demography | forecasting | global warming | matrix projection model | population dynamics | sex ratio | range limits |

Rising temperatures and extreme drought events associated with global climate change are leading to increased concern about how species will become redistributed across the globe under future climate conditions (1). Species' range limits, when not driven by dispersal limitation, should generally reflect the limits of the ecological niche (2). Niches and geographic ranges are often limited by climatic factors including temperature and precipitation (3). Therefore, any substantial changes in the magnitude of these climatic factors could impact population viability, with implications for range expansions or contractions based on which regions of a species' range become more or less suitable (4).

Forecasting range shifts for dioecious species (most animals and ca. 7% of plant species) is complicated by the potential for sexual niche differentiation, i.e. distinct responses of females and males to shared climate drivers (5). The lower cost of reproduction for one sex (male or female) may allow that sex to invest its energy in other functions that produce higher growth rates, greater clonality, or even higher survival rates compared to the other sex, leading to sexual niche differentiation (6). Accounting for sexual niche differentiation is a long-standing challenge in accurately predicting which sex will successfully track environmental change and how this will impact population viability and range shifts (7). Populations in which males are rare under current climatic conditions could experience low reproductive success due to sperm or pollen limitation that may lead to population decline in response to climate change that disproportionately favors females (8). In contrast, climate change could expand male habitat

suitability (e.g. upslope movement), which might increase seed set for mate-limited females and favor range expansion (9). Across dioecious plants, for example, studies suggest that future climate change toward hotter and drier conditions may favor male-biased sex ratios (10). Although the response of species to climate warming is an urgent and active area of research, few studies have disentangled the interaction between sex and climate drivers to understand their combined effects on population dynamics and range shifts, despite calls for such an approach (11).

The vast majority of theory and models in population biology, including those used to forecast biodiversity responses to climate change, ignore the complication of sex structure (but see 12). Traditional approaches instead focus exclusively on females, assuming that males are in sufficient supply as to never limit female fertility. In contrast, "two-sex" models are required to fully account for demographic differences between females and males and sex-specific responses to shared climate drivers (13). Sex differences in maturation, reproduction, and mortality schedules can generate skew in the operational sex ratio (OSR; sex ratio of individuals available for mating) even if the birth sex ratio is 1:1 (14). Climate and other environmental drivers can therefore influence the OSR via their influence on sex-specific demographic rates. In a two-sex framework, demographic rates both influence and respond to the OSR in a feedback loop that makes two-sex models inherently nonlinear and more data-hungry than corresponding female-dominant models. Given the additional complexity and data needs, forecasts of range dynamics for dioecious species under future climate change that explicitly account for females, males, and their inter-dependence are limited (15).

Tracking the impact of climate change on population viability

Significance Statement

The vast majority of models used to forecast population viability and range shifts in response to climate change overlook the complexity of sex structure, and thus the potential for females and males to differ in their sensitivity to climate drivers. Here, we combined a common garden experiment with mathematical models to demonstrate that accounting for only one sex could lead to an underestimation of the impact of climate change on dioecious species, particularly in regions of their range that are biased toward one sex.

J.K.M., A.C. and T.E.X.M. designed the study.
A.C. and T.E.X.M. collected the data.
All authors conducted the statistical analyses and modeling.
J.K.M. drafted the manuscript, and T.E.X.M. contributed to revisions.

The authors declare no conflict of interest

To whom correspondence should be addressed. E-mail: jkmoutouama@gmail.com

ity (λ) and distributional limits of dioecious taxa depends on our ability to build mechanistic models that take into account the spatial and temporal context of sex specific response to climate change, while accounting for sources of uncertainty (? ?). Structured population models built from demographic data collected from geographically distributed observations or common garden experiments provide several advantages for studying the impact of climate change on species' range shifts (? ? ?). First, demographic models link individual-level life history events (mortality, development, and regeneration) to population demography, allowing the investigation of factors explaining vital rate responses to environmental drivers (? ? ?). Second, demographic models have a natural interface with statistical estimation of individual-level vital rates that provide quantitative measures of uncertainty and isolate different sources of variation, features that can be propagated to population-level predictions (? ? ?). Finally, structured demographic models can be used to identify which aspects of climate are the most important drivers of population dynamics. For example, Life Table Response Experiments (LTRE) built from structured models have become widely used to understand the relative importance of covariates in explaining variation in population growth rate (? ? ?).

ACKNOWLEDGMENTS. Please include your acknowledgments here, set in a single paragraph. Please do not include any acknowledgments in the Supporting Information, or anywhere else in the manuscript.

1. Bertran R, et al. (2011) Changes in plant community composition lag behind climate warming in lowland forests. *Nature* 479(7374):517–520.
2. Gamelon M, et al. (2017) Interactions between demography and environmental effects are important determinants of population dynamics. *Science Advances* 3(2):e1602298.
3. Smith MD, et al. (2024) Extreme drought impacts have been underestimated in grasslands and shrublands globally. *Proceedings of the National Academy of Sciences* 121(4):e2309881120.
4. Lee-Yaw JA, et al. (2016) A synthesis of transplant experiments and ecological niche models suggests that range limits are often niche limits. *Ecology letters* 19(6):710–722.
5. Sexton JP, McIntyre PJ, Angert AL, Rice KJ (2009) Evolution and ecology of species range limits. *Annu. Rev. Ecol. Syst.* 40:415–436.
6. Davis MB, Shaw RG (2001) Range shifts and adaptive responses to quaternary climate change. *Science* 292(5517):673–679.
7. Pease CM, Lande R, Bull J (1989) A model of population growth, dispersal and evolution in a changing environment. *Ecology* 70(6):1657–1664.
8. Tognetti R (2012) Adaptation to climate change of dioecious plants: does gender balance matter? *Tree Physiology* 32(11):1321–1324.
9. Pottier P, Burke S, Drobniak SM, Lagisz M, Nakagawa S (2021) Sexual (in) equality? a meta-analysis of sex differences in thermal acclimation capacity across ectotherms. *Functional Ecology* 35(12):2663–2678.
10. Hultine KR, et al. (2016) Climate change perils for dioecious plant species. *Nature Plants* 2(8):1–8.
11. Morrison CA, Robinson RA, Clark JA, Gill JA (2016) Causes and consequences of spatial variation in sex ratios in a declining bird species. *Journal of Animal Ecology* 85(5):1298–1306.
12. Bruijning M, et al. (2017) Surviving in a cosexual world: A cost-benefit analysis of dioecy in tropical trees. *The American Naturalist* 189(3):297–314.
13. Jones MH, Macdonald SE, Henry GH (1999) Sex- and habitat-specific responses of a high arctic willow, *Salix arctica*, to experimental climate change. *Oikos* pp. 129–138.
14. Gissi E, et al. (2023) Exploring climate-induced sex-based differences in aquatic and terrestrial ecosystems to mitigate biodiversity loss. *nature communications* 14(1):4787.
15. Eberhart-Phillips LJ, et al. (2017) Sex-specific early survival drives adult sex ratio bias in snowy plovers and impacts mating system and population growth. *Proceedings of the National Academy of Sciences* 114(27):E5474–E5481.
16. Petry WK, et al. (2016) Sex-specific responses to climate change in plants alter population sex ratio and performance. *Science* 353(6294):69–71.
17. Field DL, Pickup M, Barrett SC (2013) Comparative analyses of sex-ratio variation in dioecious flowering plants. *Evolution* 67(3):661–672.
18. Ellis RP, et al. (2017) Does sex really matter? explaining intraspecific variation in ocean acidification responses. *Biology letters* 13(2):20160761.
19. Gissi E, Bowyer RT, Bleich VC (2024) Sex-based differences affect conservation. *Science* 384(6702):1309–1310.
20. Gerber LR, White ER (2014) Two-sex matrix models in assessing population viability: when do male dynamics matter? *Journal of Applied Ecology* 51(1):270–278.
21. Miller TE, Shaw AK, Inouye BD, Neubert MG (2011) Sex-biased dispersal and the speed of two-sex invasions. *The American Naturalist* 177(5):549–561.
22. Shelton AO (2010) The ecological and evolutionary drivers of female-biased sex ratios: two-sex models of perennial seagrasses. *The American Naturalist* 175(3):302–315.

23. Lynch HJ, et al. (2014) How climate extremes—not means—define a species' geographic range boundary via a demographic tipping point. *Ecological Monographs* 84(1):131–149.
24. Evans ME, Merow C, Record S, McMahon SM, Enquist BJ (2016) Towards process-based range modeling of many species. *Trends in Ecology & Evolution* 31(11):860–871.
25. Merow C, Bois ST, Allen JM, Xie Y, Silander Jr JA (2017) Climate change both facilitates and inhibits invasive plant ranges in new england. *Proceedings of the National Academy of Sciences* 114(16):E3276–E3284.
26. Schwinning S, Lortie CJ, Esque TC, DeFalco LA (2022) What common-garden experiments tell us about climate responses in plants.
27. Schultz EL, et al. (2022) Climate-driven, but dynamic and complex? a reconciliation of competing hypotheses for species' distributions. *Ecology letters* 25(1):38–51.
28. Ehrlén J, Morris WF (2015) Predicting changes in the distribution and abundance of species under environmental change. *Ecology letters* 18(3):303–314.
29. Louthan AM, et al. (2022) Climate change weakens the impact of disturbance interval on the growth rate of natural populations of venus flytrap. *Ecological Monographs* 92(4):e1528.
30. Dahlgren JP, Bengtsson K, Ehrlén J (2016) The demography of climate-driven and density-regulated population dynamics in a perennial plant. *Ecology* 97(4):899–907.
31. Elder BD, Miller TE (2016) Quantifying demographic uncertainty: Bayesian methods for integral projection models. *Ecological Monographs* 86(1):125–144.
32. Ellner SP, et al. (2022) A critical comparison of integral projection and matrix projection models for demographic analysis: Comment. *Ecology*.
33. Ellner SP, Childs DZ, Rees M, et al. (2016) Data-driven modelling of structured populations. *A practical guide to the Integral Projection Model*. Cham: Springer.
34. Hernández CM, Ellner SP, Adler PB, Hooker G, Snyder RE (2023) An exact version of life table response experiment analysis, and the r package exactltre. *Methods in Ecology and Evolution* 14(3):939–951.
35. Czachura K, Miller TE (2020) Demographic back-casting reveals that subtle dimensions of climate change have strong effects on population viability. *Journal of Ecology* 108(6):2557–2570.