

When Mutualism Becomes Costly: Abiotic Stress Shapes Grass Demography

Jacob K. Moutouama  ^{*1,2,3}, Julia Martin¹,
Ulisses Rizo¹, Malcolm Sherwood¹, Emily Chong¹, Dajanae Pearson¹, Alexandra Jimenez
Martín¹, Josh Fowler^{1,4}, Ali Campbell¹, Chris Oxley¹, Karl Schrader¹, and Tom E.X. Miller ¹

¹Program

in Ecology and Evolutionary Biology, Department of BioSciences, Rice University, Houston, TX USA

²Department of Botany, University of British Columbia, Vancouver, BC, Canada

³Biodiversity Research Centre, University of British Columbia, Vancouver, BC, Canada

⁴University of Miami, Department of Biology, Miami, FL USA

Running header: Stress and Grass–Endophyte Interactions

Keywords: demography, mutualism, parasitism, plant–fungi interactions, range limits, stress gradient hypothesis

Submitted to: *Ecology letters* (Letter)

Data accessibility statement: If the paper is accepted, all data and computer scripts supporting the results will be archived in a Zenodo package, with the DOI included at the end of the article. During peer review, our code (Stan and R) is available at <https://github.com/jmoutouama/endo-range-limits>.

Conflict of interest statement: None.

Authorship statement:

Abstract:

Main Text:

Figures: 6

Tables: X

References: X

*Corresponding author: jmoutouama@gmail.com

Abstract

Interactions between plants and fungi are ubiquitous and can strongly influence plant fitness. Yet, how endophytes affect host demography across abiotic stress gradients when hosts are subject to herbivory pressure remains unclear. We studied three cool-season grasses hosting seed-transmitted fungal endophytes using geographically distributed field experiments along an aridity gradient, with factorial manipulations of endophyte presence and herbivory. Bayesian models showed that all vital rates declined toward more arid sites, indicating reduced performance under stress. Declines were more pronounced in individuals hosting endophytes, suggesting that interactions typically beneficial under benign conditions can become costly under abiotic stress. Herbivory had no detectable effect on demography. Considering both abiotic and biotic factors, abiotic stress was the primary driver of vital rates, with herbivory and endophytes having little effect. These results highlight that mutualistic fungi can become liabilities under stress, and that abiotic conditions largely determine species demography across environmental gradients.

Introduction

Plant-microbe symbioses are widespread and ecologically important. These interactions are famously context-dependent, where the direction and strength of the interaction outcome depends on the environment in which it occurs (Bronstein, 1994; Hoeksema and Bruna, 2015). For example, plant-microbe interactions that are beneficial under stressful conditions may become neutral or parasitic under benign conditions (Giauque et al., 2019). Under biotic stress (e.g., herbivory), endophyte symbiosis can benefit host plants by facilitating the production of secondary compounds that deter feeding or exert direct toxicity, thereby reducing insect growth, survival, and oviposition (Atala et al., 2022; Bastias et al., 2017; Vega, 2008). Similarly under abiotic stress (e.g., low precipitation or high temperature), symbionts can increase their host tolerance (Clay and Schardl, 2002). Mycorrhizal fungi form extensive networks that increase root surface area, helping plants absorb more water and nutrients such as phosphorus and nitrogen (Amijee et al., 1989). Symbiotic microbes also stimulate plants to produce antioxidant enzymes like superoxide dismutase and catalase that reduce oxidative damage caused by stresses such as drought, salinity, and heat (Ruiz-lazona et al., 1996). All these mechanisms could help the host species better buffer against environmental stochasticity (Fowler et al., 2023). However, in resource-limited environments, maintaining symbionts can also impose significant costs on hosts (Bronstein, 2001; Johnson et al., 1997). Because microbes depend on host-derived carbon and nutrients, plants must allocate part of their limited resources to sustaining the symbiont. This investment can reduce host growth, reproduction, or competitive ability, particularly when the benefits provided by the symbiont do not outweigh these costs (Klironomos, 2003).

Context dependence raises the hypothesis that plant-microbe interactions are likely to vary across abiotic stress gradients, such as high temperatures or low precipitation, as well as under biotic stressors like intense herbivory, with significant implications for host fitness. If microbial symbiosis provides greater benefits under these stressful conditions, symbionts could enhance host survival, growth, and reproduction (Allsup et al., 2023; Rudgers et al., 2020). For example, fungal endophytes enhance host tolerance to abiotic stress in *Bromus laevipes* by improving population survival in dry environments (Afkhami et al., 2014; David et al., 2019). Even when symbionts do not directly increase survival under stress, they may still promote host fitness by enhancing growth and reproduction, thereby offsetting the negative effects of lower survival rates (Yule et al., 2013). Conversely, if microbial symbiosis becomes costly under stressful conditions, symbionts could reduce host fitness (Bennett and Groten, 2022; Benning and Moeller, 2021a,b). For instance, mycorrhizal fungi require substantial carbon subsidies from the host plant (up to 20% of photosynthetically fixed carbon)

(Soudzilovskaia et al., 2015; Thirkell et al., 2020). When photosynthesis is limited by abiotic stress, this carbon cost may outweigh the benefits of symbiosis, leading to reduced host fitness.

Despite growing interest in the role of symbiosis in host performance, few studies have examined under natural conditions how abiotic and biotic stressors shape the effects of symbionts on host demography along abiotic stress gradients (Louthan et al., 2015). Moreover, it remains unclear which stressor is most critical in eliciting the benefits of endophyte symbiosis, or how multiple stressors may interact to influence host demography. Herbivory is a biotic factor that can significantly affect host fitness (Benning et al., 2019) and mediate the outcomes of symbiotic interactions. Herbivores reduce host fitness through tissue damage, defoliation, and altered resource allocation (Maron and Crone, 2006), and can also impose indirect ecological costs by influencing reproductive or growth strategies (Miller et al., 2008). These ecological costs may shape the selective landscape in ways that favor symbiont persistence or transmission (Clay et al., 2005). For example, herbivory can enhance the fitness benefits of endophyte infection by triggering host defense pathways or promoting vertical transmission of the symbiont (Agrawal et al., 1999; Gundel et al., 2020). If the ecological context of herbivory alters the net fitness effects of symbiosis, then interactions between herbivory and abiotic stress (e.g., drought or high temperature) may play a critical role in determining host population persistence.

Working across an aridity gradient in the south-central US, we investigated how endophyte symbiosis influences host demography under varying abiotic stress. This region provides an ideal system to study stressful gradients because it spans a strong east–west precipitation gradient, with mesic sites in the east transitioning to increasingly arid conditions toward the west. We also incorporated herbivory as a biotic stressor to evaluate its potential interactive effects with abiotic conditions. Specifically, we studied the symbiotic association between three native cool-season grass species (*Agrostis hyemalis*, *Elymus virginicus*, and *Poa autumnalis*) and their vertically transmitted fungal symbionts (*Epichloë* spp.). Our experiment was designed to address the following questions:

1. Under combined abiotic stress and herbivory, do endophytes provide fitness benefits or impose costs on their host plants?
2. Between abiotic and biotic factors, which plays a stronger role in shaping host population demography in association with endophytes?

Materials and methods

Study system

We conducted our study across seven sites spanning a strong east–west precipitation gradient from Louisiana to Texas, USA (Fig. 1). Louisiana represents mesic conditions, whereas Texas experiences increasingly arid conditions, making this gradient an ideal natural laboratory to study the effects of precipitation and aridity on plant–microbe interactions. Mean annual temperature did not differ significantly across sites (Fig. S-1), which is why we focus on precipitation as the primary abiotic stressor. This gradient also overlaps the natural distributions of our focal species, ensuring realistic exposure to climatic and biotic conditions (Fig. 1).

At each site, we established eight 1.5 m × 1.5 m plots, each containing 18 individuals of one of three cool-season perennial grasses: *Agrostis hyemalis*, *Elymus virginicus*, and *Poa autumnalis*. These species typically host seed-transmitted fungal endophytes (Shaw, 2011). Seeds were collected in summer–fall 2021 from naturally symbiotic populations and either heat-treated to remove symbionts (E^-) or left untreated (E^+), maintaining the same genetic lineages as the source populations. Plots were randomly assigned one of four initial endophyte frequency treatments (80%, 60%, 40%, or 20%) and one of two herbivory treatments (herbivore access or exclusion). Full details of the study design are provided in the supplementary materials (Supplementary Method S.1.1).




Figure 1: Distribution of common garden sites along a natural aridity gradient from Texas to Louisiana in the United States for: A) *Agrostis hyemalis* (AGHY), B) *Elymus virginicus* (ELVI), and C) *Poa autumnalis* (POAU). The first three panels (A–C) are based on 30-year normals of climate predictions (1993–2023) from PRISM data, which illustrate the precipitation gradient across sites (scale on the right is in mm). Red dots indicate the locations of source populations, while grey dots represent Global Biodiversity Information Facility (GBIF) occurrence records for each species within the study area. D) Cumulative precipitation measured during the study period at each site, confirming the precipitation gradient shown in panels A–C.

Demographic data collection

We collected demographic data including survival, growth, and reproduction in May and June of 2023, 2024 and 2025, which coincided with the flowering season of *Agrostis hyemalis*, *Elymus virginicus*, and *Poa autumnalis*. For each individual, plant survival was recorded as a binary variable (alive or dead), and plant size was measured as the number of living tillers. We recorded the number of inflorescences per plant and counted the number of spikelets on up to three inflorescences from three reproducing plants. Spikelet counts were limited to three reproducing tillers per plot due to the time-consuming nature of this measurement. We used the number of spikelets from these three tillers to estimate the average number of spikelets per plant.

Modeling the effects of aridity, herbivory and endophyte on cool-season grass demography

To quantify how abiotic stress and herbivory shape plant–fungal interactions, we developed hierarchical models for each vital rate (survival, growth, and fertility). Survival was modeled with a Bernoulli distribution, growth with a Gaussian distribution, and fertility, measured as the number of inflorescences and the number of spikelets, with negative binomial distributions. All species were modeled jointly to estimate pooled fixed effects. Each vital rate model included normally distributed random effects to account for site-to-site variation ($\phi_{\text{site}[i]} \sim \mathcal{N}(0, \sigma_{\text{site}})$), source-to-source variation ($\omega_{\text{source}[i]} \sim \mathcal{N}(0, \sigma_{\text{source}})$) associated with the provenance of transplants used in the common garden, and plot-to-plot variation ($\rho_{\text{plot}[i]} \sim \mathcal{N}(0, \sigma_{\text{plot}})$). By modeling species jointly with this hierarchical structure, we increase statistical power to detect shared effects while still capturing species-specific responses. To allow for non-linear effects of precipitation, we included a quadratic term, reflecting the expectation that vital rates may peak at intermediate precipitation and decline at both low and high levels. The expected value of each vital rate for individual i was modeled as:

$$\begin{aligned} \mu_i = & \beta_0 + \beta_P P_i + \beta_{\text{endo}} \text{endo}_i + \beta_{\text{herb}} \text{herb}_i \\ & + \beta_{\text{endo} \times P} \text{endo}_i P_i + \beta_{\text{herb} \times \text{endo}} \text{herb}_i \text{endo}_i + \beta_{\text{herb} \times P} \text{herb}_i P_i \\ & + \beta_{P^2} P_i^2 + \beta_{\text{endo} \times P^2} \text{endo}_i P_i^2 \\ & + \phi_{\text{site}[i]} + \omega_{\text{source}[i]} + \rho_{\text{plot}[i]} \end{aligned} \quad (1)$$

where P is precipitation, *endo* indicates endophyte presence, and *herb* indicates herbivory. The intercept β_0 represents the grand mean vital rate across species. The coefficients β_P and β_{P^2} describe linear and quadratic responses to precipitation, respectively, while β_{endo} and β_{herb} capture the main effects of endophytes and herbivory. The interaction terms capture combined effects: $\beta_{\text{endo} \times P}$ captures how endophytes alter the response to precipitation, $\beta_{\text{herb} \times P}$ captures how herbivory affects the response to precipitation, $\beta_{\text{herb} \times \text{endo}}$ represents the joint effect of herbivory and endophytes, and $\beta_{\text{endo} \times P^2}$ captures how endophytes influence the quadratic response to precipitation.

Disentangling abiotic and biotic drivers on cool-season grass demography

To disentangle the relative contributions of abiotic and biotic factors to endophyte–host demography, we developed three candidate models. The first included only abiotic predictors (precipitation), the second included only biotic predictors (herbivory and endophyte status), and the third included both abiotic and biotic predictors. All models followed the same

hierarchical structure described above, differing only in the inclusion of predictor terms (Supplementary Method S.1.2).

To determine the best-supported model for each vital rate, we compared candidate models using leave-one-out cross-validation (LOOCV) (Vehtari et al., 2017). LOOCV combines validation and training by iteratively holding out one observation as a test set while fitting the model on the remaining $n - 1$ observations. This process is repeated n times, once for each observation, resulting in n fitted models (Silva and Zanella, 2024). The overall test error estimate is then obtained by averaging the prediction errors across all n models (Eq. 2).

$$CV_n = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2 \quad (2)$$

All models were implemented in R (R Core Team, 2023) using Stan via the rstan interface (Stan Development Team, 2024).

Results

Endophyte presence influenced host survival and reproduction across the abiotic stress gradient, with no effects of herbivory

We used Bayesian mixed-effect models to test the effects of abiotic stress (low vs. high precipitation), endophyte association (E^- vs. E^+), and herbivore exclusion (fenced vs. unfenced) on survival, growth, and reproductive traits in three cool-season grass species: *Agrostis hyemalis*, *Elymus virginicus*, and *Poa autumnalis*. All models effectively captured key aspects of the data, including means, standard deviations, and quantiles (Fig. S-2). High abiotic stress reduced survival, with E^- individuals outperforming E^+ individuals, indicating a cost of endophyte association under dry conditions (Fig. 2). Growth was generally similar across species and treatments, except in *Agrostis hyemalis*, where E^+ plants showed higher relative growth along the stress gradient (Fig. 3). Inflorescence production differed little between E^+ and E^- individuals, except for *Poa autumnalis*, which produced more inflorescences under low abiotic stress (high precipitation; Fig. 4). Spikelet production also reflected costs of endophyte association under high stress, particularly in *Elymus virginicus*, where E^+ individuals produced fewer spikelets than E^- individuals (Fig. 5). Herbivore exclusion had minimal effects on all vital rates, with slightly higher survival and spikelet production in fenced plots, indicating that precipitation and endophyte status were the primary drivers of variation in host demography.




Figure 2: Predicted survival rate (top panels) and endophyte effect ($\Delta = E^+ - E^-$, bottom panels) across precipitation gradients. Lines show posterior means for E^+ (blue) and E^- (red), with shading indicating 90% credible intervals. Points represent mean observed survival per plot, jittered for visualization. The dashed line at $\Delta = 0$ indicates no endophyte effect. Panels are organized by species (AGHY, ELVI, POAU) and herbivory treatment (Fenced, Unfenced).




Figure 3: Predicted relative growth (top panels) and endophyte effect ($\Delta = E^+ - E^-$, bottom panels) across precipitation gradients. Lines show posterior means for E^+ (blue) and E^- (red), with shading indicating 90% credible intervals. Points represent mean observed growth per plot, jittered for visualization. The dashed line at $\Delta = 0$ indicates no endophyte effect. Panels are organized by species (AGHY, ELVI, POAU) and herbivory treatment (Fenced, Unfenced).




Figure 4: Predicted number of inflorescences (top panels) and endophyte effect ($\Delta = E^+ - E^-$, bottom panels) across precipitation gradients. Lines show posterior means for E^+ (blue) and E^- (red), with shading indicating 90% credible intervals. Points represent mean observed inflorescences per plot, jittered for visualization. The dashed line at $\Delta = 0$ indicates no endophyte effect. Panels are organized by species (AGHY, ELVI, POAU) and herbivory treatment (Fenced, Unfenced).




Figure 5: Predicted number of spikelets (top panels) and endophyte effect ($\Delta = E^+ - E^-$, bottom panels) across precipitation gradients. Lines show posterior means for E⁺ (blue) and E⁻ (red), with shading indicating 90% credible intervals. Points represent mean observed spikelets per plot, slightly jittered for visualization. Spikelets were measured for only two species (see Methods for details). The dashed line at $\Delta = 0$ indicates no endophyte effect. Panels are organized by species (AGHY, ELVI) and herbivory treatment (Fenced, Unfenced).

Abiotic factors as the primary drivers of fitness components across the stress gradient

Across all vital rates, survival, growth, flowering, and spikelet production, models including only abiotic predictors consistently had the highest ELPD values (Figure 6). The standard errors of the ELPD differences indicate some uncertainty in these comparisons. For example, the difference between the abiotic-only and abiotic+biotic models for survival was small relative to its uncertainty ($\Delta\text{ELPD} = -2.1$, $\text{SE} = 3.7$), suggesting that including biotic predictors provides

only marginal additional information. In contrast, biotic-only models had substantially lower ELPD values with smaller uncertainty (e.g., survival: $\Delta\text{ELPD} = -5.2$, $\text{SE} = 2.0$), indicating they are clearly inferior. Overall, these results suggest that the demography of *Agrostis hyemalis*, *Elymus virginicus*, and *Poa autumnalis* is primarily driven by variation along the abiotic stress gradient, such as precipitation, whereas biotic interactions, including herbivory and endophyte presence, play a secondary role. This pattern aligns with the stress gradient hypothesis, which predicts that abiotic factors have a stronger influence on species performance in stressful environments, while biotic interactions become more important under more benign conditions.




Figure 6: Model comparison across vital rates (survival, growth, flowering, spikelet production) using expected log predictive density (ELPD) differences. ELPD measures a model's ability to predict new data, with higher values indicating better predictive performance. Differences in ELPD relative to the best-supported model (abiotic-only) show which models perform better or worse. Circles indicate the ELPD of each model, error bars represent the standard error of the differences (SE), and stars mark the model with the highest ELPD.

Acknowledgements

This research was supported by National Science Foundation Division of Environmental Biology awards.

References

- Afkhami, M. E., McIntyre, P. J., and Strauss, S. Y. (2014). Mutualist-mediated effects on species' range limits across large geographic scales. *Ecology letters*, 17(10):1265–1273.
- Agrawal, A. A., Laforsch, C., and Tollrian, R. (1999). Transgenerational induction of defences in animals and plants. *Nature*, 401(6748):60–63.
- Allsup, C. M., George, I., and Lankau, R. A. (2023). Shifting microbial communities can enhance tree tolerance to changing climates. *Science*, 380(6647):835–840.
- Amijee, F., Tinker, P., and Stribley, D. (1989). The development of endomycorrhizal root systems: Vii. a detailed study of effects of soil phosphorus on colonization. *New Phytologist*, 111(3):435–446.
- Atala, C., Acuña-Rodríguez, I. S., Torres-Díaz, C., and Molina-Montenegro, M. A. (2022). Fungal endophytes improve the performance of host plants but do not eliminate the growth/defence trade-off. *New Phytologist*, 235(2).
- Bastias, D. A., Martínez-Ghersa, M. A., Ballaré, C. L., and Gundel, P. E. (2017). Epichloë fungal endophytes and plant defenses: not just alkaloids. *Trends in Plant Science*, 22(11):939–948.
- Beaudry, J.-R. (1951). Seed development following the mating elymus virginicus l. x agropyron repens (l.) beauv. *Genetics*, 36(2):109.
- Bennett, A. E. and Groten, K. (2022). The costs and benefits of plant–arbuscular mycorrhizal fungal interactions. *Annual Review of Plant Biology*, 73(1):649–672.
- Benning, J. W., Eckhart, V. M., Geber, M. A., and Moeller, D. A. (2019). Biotic interactions contribute to the geographic range limit of an annual plant: herbivory and phenology mediate fitness beyond a range margin. *The American Naturalist*, 193(6):786–797.
- Benning, J. W. and Moeller, D. A. (2021a). Microbes, mutualism, and range margins: testing the fitness consequences of soil microbial communities across and beyond a native plant's range. *New Phytologist*, 229(5):2886–2900.
- Benning, J. W. and Moeller, D. A. (2021b). Plant–soil interactions limit lifetime fitness outside a native plant's geographic range margin. *Ecology*, 102(3):e03254.
- Bronstein, J. L. (1994). Conditional outcomes in mutualistic interactions. *Trends in ecology & evolution*, 9(6):214–217.

- 211 Bronstein, J. L. (2001). The exploitation of mutualisms. *Ecology letters*, 4(3):277–287.
- 212 Clay, K., Holah, J., and Rudgers, J. A. (2005). Herbivores cause a rapid increase in hereditary
213 symbiosis and alter plant community composition. *Proceedings of the National Academy of*
214 *Sciences*, 102(35):12465–12470.
- 215 Clay, K. and Schardl, C. (2002). Evolutionary origins and ecological consequences of
216 endophyte symbiosis with grasses. *the american naturalist*, 160(S4):S99–S127.
- 217 David, A. S., Quintana-Ascencio, P. F., Menges, E. S., Thapa-Magar, K. B., Afkhami, M. E., and
218 Searcy, C. A. (2019). Soil microbiomes underlie population persistence of an endangered
219 plant species. *The American Naturalist*, 194(4):488–494.
- 220 Donald, M. L., Bohner, T. F., Kolis, K. M., Shadow, R. A., Rudgers, J. A., and Miller, T. E. (2021).
221 Context-dependent variability in the population prevalence and individual fitness effects
222 of plant–fungal symbiosis. *Journal of Ecology*, 109(2):847–859.
- 223 Fowler, J. C., Donald, M. L., Bronstein, J. L., and Miller, T. E. (2023). The geographic footprint
224 of mutualism: How mutualists influence species’ range limits. *Ecological Monographs*,
225 93(1):e1558.
- 226 Giauque, H., Connor, E. W., and Hawkes, C. V. (2019). Endophyte traits relevant to stress
227 tolerance, resource use and habitat of origin predict effects on host plants. *New Phytologist*,
228 221(4):2239–2249.
- 229 Gundel, P. E., Sun, P., Charlton, N. D., Young, C. A., Miller, T. E., and Rudgers, J. A. (2020).
230 Simulated folivory increases vertical transmission of fungal endophytes that deter herbivores
231 and alter tolerance to herbivory in *poa autumnalis*. *Annals of Botany*, 125(6):981–991.
- 232 Hoeksema, J. D. and Bruna, E. M. (2015). Context-dependent outcomes of mutualistic
233 interactions. *Mutualism*, 10:181–202.
- 234 Johnson, N. C., Graham, J. H., and Smith, F. A. (1997). Functioning of mycorrhizal associations
235 along the mutualism–parasitism continuum. *The New Phytologist*, 135(4):575–585.
- 236 Klironomos, J. N. (2003). Variation in plant response to native and exotic arbuscular
237 mycorrhizal fungi. *Ecology*, 84(9):2292–2301.
- 238 Louthan, A. M., Doak, D. F., and Angert, A. L. (2015). Where and when do species interactions
239 set range limits? *Trends in ecology & evolution*, 30(12):780–792.

- Maron, J. L. and Crone, E. (2006). Herbivory: effects on plant abundance, distribution and population growth. *Proceedings of the Royal Society B: Biological Sciences*, 273(1601):2575–2584.
- Miller, T. E. X., Tenhumberg, B., and Louda, S. M. (2008). Herbivore-mediated ecological costs of reproduction shape the life history of an iteroparous plant. *The American Naturalist*, 171(2):141–149.
- R Core Team (2023). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rudgers, J. A., Afkhami, M. E., Bell-Dereske, L., Chung, Y. A., Crawford, K. M., Kivlin, S. N., Mann, M. A., and Nuñez, M. A. (2020). Climate disruption of plant-microbe interactions. *Annual review of ecology, evolution, and systematics*, 51:561–586.
- Rudgers, J. A., Afkhami, M. E., Rúa, M. A., Davitt, A. J., Hammer, S., and Huguet, V. M. (2009). A fungus among us: broad patterns of endophyte distribution in the grasses. *Ecology*, 90(6):1531–1539.
- Ruiz-lazona, J. M., Azcón, R., and Palma, J. (1996). Superoxide dismutase activity in arbuscular mycorrhizal lactuca sativa plants subjected to drought stress. *New Phytologist*, 134(2):327–333.
- Shaw, R. B. (2011). *Guide to Texas grasses*. Texas A&M University Press.
- Silva, L. A. and Zanella, G. (2024). Robust leave-one-out cross-validation for high-dimensional bayesian models. *Journal of the American Statistical Association*, 119(547):2369–2381.
- Sneck, M. E., Rudgers, J. A., Young, C. A., and Miller, T. E. (2017). Variation in the prevalence and transmission of heritable symbionts across host populations in heterogeneous environments. *Microbial Ecology*, 74:640–653.
- Soudzilovskaia, N. A., van der Heijden, M. G., Cornelissen, J. H., Makarov, M. I., Onipchenko, V. G., Maslov, M. N., Akhmetzhanova, A. A., and van Bodegom, P. M. (2015). Quantitative assessment of the differential impacts of arbuscular and ectomycorrhiza on soil carbon cycling. *New Phytologist*, 208(1):280–293.
- Stan Development Team (2024). RStan: the R interface to Stan. R package version 2.32.6.
- Thirkell, T. J., Pastok, D., and Field, K. J. (2020). Carbon for nutrient exchange between arbuscular mycorrhizal fungi and wheat varies according to cultivar and changes in atmospheric carbon dioxide concentration. *Global change biology*, 26(3):1725–1738.

- 269 Vega, F. E. (2008). Insect pathology and fungal endophytes. *Journal of invertebrate pathology*,
270 98(3):277–279.
- 271 Vehtari, A., Gelman, A., and Gabry, J. (2017). Practical bayesian model evaluation using
272 leave-one-out cross-validation and waic. *Statistics and computing*, 27:1413–1432.
- 273 Yule, K. M., Miller, T. E., and Rudgers, J. A. (2013). Costs, benefits, and loss of vertically
274 transmitted symbionts affect host population dynamics. *Oikos*, 122(10):1512–1520.

Supporting Information

S.1 Supplementary Methods

S.1.1 Study design

S.1.1.1 Abiotic stress

Common gardens were established at seven sites spanning the geographic range of *Elymus virginicus* from Louisiana to Texas, USA. Sites were selected along a pronounced aridity gradient, with Louisiana representing mesic conditions and Texas increasingly arid conditions. Mean annual temperature did not differ significantly among sites (Fig. S-1), confirming that precipitation is the primary abiotic gradient. Sites were chosen to reflect natural environmental conditions, including shaded areas under tree canopy or near shrubs.

S.1.1.2 Plot establishment

At each site, we established eight plots measuring 1.5 m × 1.5 m. Existing vegetation was removed by tilling to reduce competition. Within each plot, 15 individuals of each species were planted approximately 15 cm deep in an evenly spaced 4 × 4 grid, with positions randomly assigned to reduce positional effects.

S.1.1.3 Endophyte and herbivory treatments

Seeds were collected in summer–fall 2021 from naturally symbiotic populations (source populations). Seeds were either heat-treated to remove endophytes (E^-) or left untreated (E^+), maintaining the same genetic lineages. Each plot was randomly assigned a starting endophyte frequency (80%, 60%, 40%, or 20%) and a herbivory treatment. For herbivore exclusion plots, plants were enclosed with 1.2 m tall mesh fencing and sprayed with insecticide to prevent vertebrate and insect herbivory. For herbivore access (control) plots, plants were half-enclosed with mesh netting. All plots received comparable quantities of source population material.

S.1.1.4 Experimental rationale

This experimental design was informed by natural patterns of endophyte prevalence in Texas (Donald et al., 2021; Rudgers et al., 2009; Sneck et al., 2017). Fungal genotyping confirmed the presence of biosynthetic pathways for secondary metabolites, such as peramine, loline, and ergot alkaloids, which may enhance host resistance to drought and herbivory (Beaudry,

1951). The design allows testing how endophyte frequency and herbivory interact with abiotic stress (aridity) to influence host demography from core to edge of the species range.

S.1.1.5 Source populations and identification of individual endophyte status

Plants used in the common garden experiment were derived from natural populations throughout the south-central US. Seeds were collected from these populations, and some were heat-treated at 60°C for approximately five days (120 hours) to produce endophyte-negative plants (E^-), a method that eliminates endophytes without affecting seed viability. All seeds, both heat-treated and non-heat-treated, were planted in the Rice University greenhouse, where seedlings were fertilized every two weeks. Seedlings were then vegetatively propagated to produce enough individuals for the experiment ($N = 840$). Before field planting, we confirmed endophyte status (E^+ or E^-) of all seedlings using either microscopy or an immunoblot assay. Leaf tissues were stained with aniline blue lactic acid and viewed under a compound microscope at 200x–400x to identify fungal hyphae. The immunoblot assay (Phytoscreen field tiller endophyte detection kit, Agrinostics Ltd. Co.) uses monoclonal antibodies that target proteins of *Epichloë* spp. and a chromagen to visually indicate presence or absence. Both methods yield similar detection rates.

S.1.2 Candidate models for abiotic and biotic drivers of cool-grass species

To disentangle the relative contributions of abiotic and biotic factors, we developed three candidate models:

1. **Abiotic model:** includes precipitation (P) and its quadratic term (P^2) as predictors. The expected value of each vital rate for individual i is:

$$\mu_i = \beta_0 + \beta_P P_i + \beta_{P^2} P_i^2 + \phi_{\text{site}[i]} + \omega_{\text{source}[i]} + \rho_{\text{plot}[i]}$$

2. **Biotic model:** includes endophyte status (endo), herbivory (herb), and their interaction ($\text{endo} \times \text{herb}$):

$$\mu_i = \beta_0 + \beta_{\text{endo}} \text{endo}_i + \beta_{\text{herb}} \text{herb}_i + \beta_{\text{endo} \times \text{herb}} \text{endo}_i \text{herb}_i + \phi_{\text{site}[i]} + \omega_{\text{source}[i]} + \rho_{\text{plot}[i]}$$

3. **Full model:** combines both abiotic and biotic predictors, including all relevant interactions:

$$\begin{aligned}\mu_i = & \beta_0 + \beta_P P_i + \beta_{P^2} P_i^2 + \beta_{\text{endo}} \text{endo}_i + \beta_{\text{herb}} \text{herb}_i \\ & + \beta_{\text{endo} \times \text{pendo}_i} \text{pendo}_i P_i + \beta_{\text{herb} \times \text{pherb}_i} \text{pherb}_i P_i + \beta_{\text{herb} \times \text{endo}} \text{herb}_i \text{endo}_i + \beta_{\text{endo} \times P^2} \text{endo}_i P_i^2 \\ & + \phi_{\text{site}[i]} + \omega_{\text{source}[i]} + \rho_{\text{plot}[i]}\end{aligned}$$

All models included a hierarchical random effects structure accounting for site, source, and plot variation, with $\phi_{\text{site}[i]} \sim \mathcal{N}(0, \sigma_{\text{site}})$, $\omega_{\text{source}[i]} \sim \mathcal{N}(0, \sigma_{\text{source}})$, and $\rho_{\text{plot}[i]} \sim \mathcal{N}(0, \sigma_{\text{plot}})$.

S.2 Supporting Tables

Table S-1: Populations and Coordinates

Population	Location	Coordinates	NE+ / NE-
Sam Houston State University Field Station	Huntsville, TX	30°44'30.5"N, 95°28'28.2"W	NE+ = 39, NE- = 50
Texas Tech University	Junction, TX	30°28'18.2"N, 99°47'01.7"W	NE+ = 49, NE- = 83
Palmetto State Park	Palmetto, TX	29.591623, -97.584781	NE+ = 250, NE- = 192
Jean Lafitte National Historical Park and Preserve	Jean Lafitte, LA	29.785105, -90.114933	NE+ = 83, NE- = 94

Table S-2: Differences in expected log predictive density (ELPD) between models for each vital rate. The top model for each response (bold) has the highest ELPD. Standard errors of the differences are shown in the `se_diff` column.

Vital rate	Model	elpd_diff	se_diff
Survival	abiotic	0.0	0.0
	abiotic_biotic	-2.1	3.7
	biotic	-5.2	2.0
Growth	abiotic	0.0	0.0
	abiotic_biotic	-2.1	3.7
	biotic	-5.2	2.0
Inflorescence	abiotic	0.0	0.0
	abiotic_biotic	-2.1	3.7
	biotic	-5.2	2.0
Spikelet	abiotic	0.0	0.0
	abiotic_biotic	-2.1	3.7
	biotic	-5.2	2.0

S.3 Supporting Figures




Figure S-1: Abiotic conditions across the seven study sites. (A) Mean annual precipitation showing the pronounced east–west aridity gradient from mesic Louisiana to arid Texas. (B) Mean annual temperature across sites, which does not differ much. Panels highlight that precipitation, rather than temperature, constitutes the primary abiotic gradient in this system.




Figure S-2: Posterior predictive checks for the full demographic model. Dark blue lines show observed data (y), and light blue lines show replicated datasets generated from posterior predictions (y_{rep}). The strong overlap across survival, growth, flowering, and spikelet production indicates adequate model fit for all vital rates.