

When mutualism weakens: stress alters endophyte benefits in cool-season grasses

Jacob K. Moutouama *¹, Josh Fowler², Akiem
Gough³, Julia Martin³, Chris Oxley³, Karl Schrader³, Karis Williams³, and Tom E.X. Miller ³

¹Department of Botany, University of British Columbia, Vancouver, BC, Canada

²Josh has a new CU affiliation

³Program

in Ecology and Evolutionary Biology, Department of BioSciences, Rice University, Houston, TX USA

Running header: Stress and Grass–Endophyte Interactions

Keywords: demography, mutualism, parasitism, plant–fungi interactions, range limits, stress gradient hypothesis

Submitted to: *Ecology letters* (Letter)

Data accessibility statement: If the paper is accepted, all data and computer scripts supporting the results will be archived in a Zenodo package, with the DOI included at the end of the article. During peer review, our **data and code**¹ (Stan and R) is available at <https://github.com/jmoutouama/endo-range-limits>.

Conflict of interest statement: None.

Acknowledgments This research was supported by National Science Foundation Division of Environmental Biology award 2208857.

Authorship statement:

Abstract:

Main Text:

Figures: 5

Tables: 1

References: X

*Corresponding author: jmoutouama@gmail.com

¹*If the data are not on github as csv's then I would create a temporary dropbox link so that reviewers could reproduce the analysis.*

Abstract

Benefits of plant-microbe symbioses are predicted to increase under environmental stress, but the relative roles of abiotic and biotic factors as modifiers of symbiotic interaction outcomes remain unclear. We studied how the independent and combined effects of aridity and herbivory modify the demographic effects of seed-transmitted fungal endophytes on three native grass host species using geographically distributed field experiments along a precipitation gradient, crossed with herbivore exclusion. Host fitness declined under increasing abiotic stress, and **endophyte-symbiotic grasses often fared worse than non-symbiotic ones**², suggesting that interactions that are beneficial under benign conditions can become costly under stress. **Herbivory had no detectable effect on host demography or endophyte-mediated outcomes**.³ Overall, abiotic stress was the primary driver of mutualism outcomes between plants and fungal symbionts, but in the opposite direction as predicted by theory and prior work. These results highlight that symbiotic fungi can become liabilities under stress, with implications for the dynamics of plant-fungal mutualisms under global change.

²*I know space is tight but can we make this a more specific or quantitative summary of the results?*

³*I think this is not true in the new results...?*

15 Introduction

16 Symbioses are widespread and ecologically important, occurring across both plants and
17 animals, and often mediating key ecological functions such as stress tolerance and defense
18 against natural enemies. These interactions are famously context-dependent, with the direction
19 and strength of outcomes shaped by the environment in which they occur (Bronstein, 1994;
20 Hoang et al., 2024; Hoeksema and Bruna, 2015). According to the stress-gradient hypothesis
21 (SGH), positive interactions are generally expected to increase under stressful conditions, but
22 actual outcomes depend on the type and intensity of biotic and abiotic stress (Bertness and
23 Callaway, 1994; Holmgren and Scheffer, 2010; Louthan et al., 2015). For example, microbial
24 symbionts that benefit host plants under drought stress may become neutral or even parasitic
25 under more benign conditions (Giauque et al., 2019).

26 Because microbial symbioses are expected to become more beneficial in stressful
27 conditions, they can substantially enhance host resilience to environmental stress. Under
28 biotic stress, such as herbivory, microbial symbiosis can benefit host plants by facilitating
29 the production of secondary compounds that deter feeding or exert direct toxicity, thereby
30 reducing herbivore growth, survival, and oviposition (Atala et al., 2022; Bastias et al., 2017;
31 Vega, 2008). Similarly, under abiotic stress, such as low precipitation, nutrient limitation,
32 or high temperature, symbiotic microbes can enhance host tolerance through a variety of
33 mechanisms (Afkhami and Rudgers, 2008; Clay and Schardl, 2002). Symbionts including
34 mycorrhizal fungi, rhizobia, and endophytic bacteria can improve water and nutrient
35 acquisition by increasing root surface area, facilitating nutrient mobilization, or promoting
36 nitrogen fixation, allowing plants to maintain growth and metabolic function under drought
37 or poor soil conditions (Amijee et al., 1989). In addition, many symbiotic microbes stimulate
38 the production of antioxidant enzymes, which reduce oxidative damage caused by stressors
39 like drought, salinity, and heat (Ruiz-lazona et al., 1996). By improving resource acquisition,
40 modulating plant physiology, and enhancing stress-responsive metabolism, these symbiotic
41 interactions help host plants buffer against environmental stochasticity and extreme conditions
42 across a wide range of ecosystems (Fowler et al., 2023).

43 While stress is hypothesized to strengthen the benefits of symbiotic microbes, growing
44 evidence suggests that it could also elevate costs of symbiosis. Maintaining symbionts
45 requires hosts to allocate valuable carbon and nutrients to sustain their microbial partners.
46 For instance, mycorrhizal fungi may require up to 20% of photosynthetically fixed carbon
47 (Soudzilovskaia et al., 2015; Thirkell et al., 2020), and under abiotic stress, this investment can
48 outweigh the benefits of the symbiosis (Bronstein, 2001; Johnson et al., 1997), reducing host
49 survival, growth, reproduction, or competitive ability (Klironomos, 2003). Similarly, vertically

transmitted endophytes can shift from mutualistic to antagonistic interactions when they produce stomata that abort host inflorescences or impose energetic costs (Saikkonen et al., 2004). For example, under low nutrient or water conditions, endophyte-infected grasses may produce fewer tillers and lower total biomass compared to endophyte-free plants, reflecting the costs of sharing limited resources (Ahlholm et al., 2002)

Despite growing interest in the role of symbiosis in shaping host demography, few studies have tested how abiotic and biotic stressors synergistically influence the demographic effects of symbionts along environmental gradients. Most work has been conducted in controlled or common garden experiments. In *Ammonophila breviligulata*, herbivory but not drought reduced growth, and no interaction was observed; however, plants hosting endophytic fungi were more resistant to these stressors (Emery et al., 2010). Studying the combined effects of abiotic and biotic stressors under natural conditions is particularly important in the context of global change (Louthan et al., 2015), as shifting climate patterns and altered herbivore pressures may change the balance of costs and benefits in plant-microbe interactions, potentially influencing species distributions and ecosystem function. Understanding how abiotic stress, biotic stress, and microbial symbiosis interact to influence plant performance requires experiments that jointly vary all three factors. However, to our knowledge, no such comprehensive study has been conducted. Here, we address this gap by examining the effects of biotic and abiotic stress on grass-endophyte symbioses in three grass species spanning a natural range-core/range-edge precipitation gradient.

Fungal endophytes of plants provide a tractable system for studying the ecological effects and context-dependence of host-symbiont interactions, as host status (symbiotic vs symbiont-free) can be experimentally manipulated. A well-studied example is the association between cool-season grasses and seed-transmitted fungal endophytes in the genus *Epichloë* (Clay et al., 2005; Oberhofer et al., 2014; Saikkonen et al., 2010; Schardl et al., 2004). Because these fungi are vertically transmitted, the fitness of host plants is closely linked to that of their symbionts, which is expected to favor the evolution of mutualistic interactions (Gundel et al., 2011, 2024). In many cases, endophyte-symbiotic grasses outperform endophyte-free grasses under water limitation, although the benefits depend on host and endophyte identity (Decunta et al., 2021). Beyond these performance effects, infection frequency itself is shaped by environmental conditions, including long-term variation in precipitation and temperature (Fowler et al., 2025; Rudgers et al., 2016). Even when endophytes do not enhance survival under stress, they may still boost host fitness by promoting growth and reproduction (Yule et al., 2013). More generally, prior work has shown that effects of endophyte symbiosis may vary in magnitude and even sign across host vital rates (Rudgers et al., 2012), which argues for considering multiple life history pathways. Importantly, hosting endophytes can entail

costs, including the allocation of plant carbon to sustain the fungal partner and potential risks associated with horizontal transmission. These costs may be offset by benefits such as herbivore deterrence, mediated by fungal alkaloids that protect against both invertebrate and vertebrate herbivores (Clay et al., 2005; Saikkonen et al., 2010) Together, these features make grass–*Epichloë* symbioses an ideal system for investigating how abiotic stressors, such as drought, interact with biotic stressors, like herbivory, to determine the outcome of symbiosis.

Across an aridity gradient in the south-central United States, we investigated how endophyte symbiosis affects multiple host vital rates under varying abiotic and biotic stress. This gradient, from mesic conditions in the east to increasingly arid conditions in the west, provides a biologically realistic stress gradient, and coincides with the westernmost range limits of many eastern North American grass species. At each of seven sites along the gradient, we manipulated vertebrate and invertebrate herbivory (allowing herbivore access versus excluding herbivores) to test its interaction with abiotic stress. We also examined the effects of *Epichloë* symbiosis by comparing endophyte-symbiotic and endophyte-free individuals across three species of native cool-season grasses.

We addressed how variation in precipitation and herbivory influences the outcome of grass–*Epichloë* symbioses. Specifically, we asked: (i) under reduced precipitation, do endophytes enhance host fitness by mitigating drought stress (stress-amelioration scenario), or can they impose a net cost under extreme combined stress from drought and herbivory (reduced-benefit scenario)? and (ii) does the relative importance of endophyte-mediated benefits change along the stress gradient, as predicted by the stress-gradient hypothesis (SGH), with higher abiotic stress amplifying mutualistic effects? We predicted that endophytes would enhance host fitness under moderate drought in the absence of herbivores, representing an endophyte-driven stress amelioration scenario (Fig. 1A). When herbivores are added as a second stress under more extreme drought, the metabolic cost of maintaining the endophyte may reduce benefits or even impose a net cost, corresponding to a reduced endophyte-mediated benefits scenario (Fig. 1B). Overall, consistent with the SGH, endophyte-mediated benefits are expected to increase under lower precipitation, with the presence or absence of herbivores modulating these effects.

Materials and methods

Study system

Our study focused on three cool-season perennial grasses native to woodland and prairie habitats of eastern North America: *Agrostis hyemalis*, *Elymus virginicus*, and *Poa autumnalis* (Poaceae, subfamily Pooideae) (Shaw, 2011). *A. hyemalis* is a small, short-lived perennial

that germinates from autumn to late winter and typically flowers between March and July. *E. virginicus* is a larger, long-lived species that begins flowering as early as March or April and continues throughout the summer. *P. autumnalis* has a similar growth form to *E. virginicus* and flowers from late spring through summer. *Agrostis hyemalis* hosts the endophyte *Epichloë amarillans* (White Jr, 1994), *E. virginicus* typically hosts *Epichloë elymi* (Liu, 2023), and *P. autumnalis* hosts *Epichloë PauTG-1* or *E. typhina subsp. poae* (Gundel et al., 2020). Like other *Epichloë* species, these endophytes are primarily transmitted vertically through seeds and are additionally capable of horizontal transmission via sexual stromata on inflorescences or asexual conidia on leaf surfaces (Clay and Schardl, 2002; Gundel et al., 2020; Leuchtmann et al., 2014). Herbivory in this system is primarily from generalist insects and small mammalian grazers, which can influence both plant performance and endophyte-mediated defenses.

Source populations and symbiont removal

Plants used in our experiment were derived from natural (source) populations of each species (Fig. 2, Table S1). Prevalence of fungal endophytes in these natural populations ranged from XX% to XX% endophyte-positive hosts (NEW SUPP TABLE). **Seeds from these source populations (from ca. 30 plants per population) were either heat-treated or left untreated, intended to produce symbiont-free (E^-) and symbiotic (E^+) plants from the same genetic lineages. For *Elymus virginicus* and *Poa autumnalis*, heat-treated seeds were placed in a drying oven at 60°C for approximately five days. For *Agrostis hyemalis*, seeds were heat-treated in a water bath at 60°C for 12 minutes.**⁴ All seeds, both heat-treated and non-heat-treated, were surface-sterilized with 10% bleach solution then germinated on agar plates. Seedlings were transferred to potting soil and grown in the Rice University greenhouse. Once these plants were sufficiently large in size they were vegetatively propagated, with clonal identities tracked across individuals.⁵ **Before transfer to the field experiment, we confirmed endophyte status (E^+ or E^-) of all seedlings using either microscopy or an immunoblot assay.**⁶ For microscopy, peels of the inner leaf sheaths were stained with aniline blue lactic acid and viewed under a compound microscope at 200x–400x to identify fungal hyphae⁷. The immunoblot assay (Phytoscreen field tiller endophyte detection kit, Agrinostics Ltd. Co.) uses monoclonal antibodies that target proteins of *Epichloë* spp. and a chromagen to visually indicate presence or absence (?). For

⁴We should talk about this. You need more detail here – I have tried adding some but there is more to do. They were all treated but not all treatments were successful, which we need to report. And there were a few different methods used on different species. This should all go in the supplement. It is also not true that heat treatment has no effect on seed viability, which is why we tried to use seeds that were multiple generations removed from the heat treatment.

⁵Add a sentence giving basic descriptive statistics of how many genotypes and clones were used for each species.

⁶Realized numbers of E^+ and E^- per species per population should be provided in a suppl table.

⁷cite

genotypes that were vegetatively propagated, we tested the endophyte status of a single clonal replicate and assume that its status applies equally to all clones of a given source genotype. Greenhouse-raised plants were planted into the field experiment in February-March 2023.

Common garden experimental design and climate data

To examine the effects of endophyte symbiosis across an abiotic stress gradient, we established common garden experiments with E+ and E- hosts at seven sites along an aridity gradient from Louisiana to central Texas, US (Fig. 2, Table S2). From west to east, mean annual precipitation increases more than three-fold across these sites, while mean annual temperature is relatively consistent (Fig. S1). For this reason, we focus on precipitation as the primary abiotic factor that varied across sites. This geographic gradient overlaps and exceeds the natural range edges of our focal species, ensuring realistic exposure to climatic and biotic conditions (Fig. 2A,B, C).

To characterize climate at these sites during the study period (2023–2025), we collected monthly temperature and precipitation data from PRISM at 800 m resolution (Hart and Bell, 2015), as daily data were not available for this period. For each site, we calculated mean temperature (°C) and cumulative precipitation (mm) over the corresponding census year, defined as the period from transplanting to the demographic census. During our study period, realized precipitation at these sites corresponded well to 30-year normals, with wetter eastern sites and drier western sites (Fig. 2D).

At each site along the gradient, we established 24 square plots (1.5 m × 1.5 m), eight per host species. Within each site, each plot was randomly assigned to either herbivore access or herbivore exclusion (Fig. 2E). The herbivory treatment targeted both vertebrate and invertebrate herbivores: exclusion plots had fencing on four sides and were sprayed with Sevin insecticide (active ingredient Zeta-Cypermethrin) 2-3 times per growing season following the manufacturer's protocol, and access plots were fenced on two sides to control for any fence effect.

Each plot was planted with 15 individuals, a mix of E+ and E- host-plants. Initial endophyte prevalence varied from 20% (3 E+, 12 E-) to 80% (12 E+, 3 E-); this aspect of the experimental design was intended to address a different question about change in symbiont prevalence across generations, and is not further considered here. For each species at each site along the precipitation gradient, we planted a total of 60 E+ and 60 E- host-plants, half of those experiencing herbivore access and half experiencing herbivore exclusion.

The Kerrville and Sonora sites were destroyed and subsequently replanted in 2024. However, *Poa autumnalis* (POAU) was not replanted at the Sonora site.

Demographic data collection

We collected demographic data, including survival, growth, and reproduction, during the flowering season of each species: *Agrostis hyemalis* and *Poa autumnalis* were censused in May and *Elymus virginicus* was censused in June of 2023, 2024, and 2025. For each individual, survival was recorded as alive or dead and, for surviving plants, size was measured as the count of tillers. Growth was quantified as the log ratio of tiller count between consecutive censuses (i.e., $\log(\text{tiller}_{t+1}/\text{tiller}_t)$) for plants that were alive in years t and $t+1$. We recorded the number of inflorescences per plant for all three species and counted the number of spikelets on up to three inflorescences from reproducing individuals of *Poa autumnalis* and *Elymus virginicus*. In *A. hyemalis*, one spikelet yields one seed, while in *P. autumnalis* and *E. virginicus* a single spikelet can produce 2-6 seeds. Furthermore, to ensure that our herbivory treatment reflected herbivory experienced by individual plants, we defined damage as visible herbivore-induced foliar tissue loss and estimated the proportion of damaged plants per plot, site, and herbivore treatment.

Statistical modeling

To assess whether endophytes confer benefits or impose costs under abiotic and biotic stress, we fit generalized linear mixed models for each vital rate response in a hierarchical Bayesian statistical framework. All models shared the same hierarchical structure for their expected values and were applied to survival, growth, and fertility. Survival was modeled with a Bernoulli distribution, growth with a Gaussian distribution, inflorescence count with a zero-inflated negative binomial distribution, and spikelet count with a negative binomial distribution. All models included species-specific fixed effects for symbiosis (endophyte presence/absence), precipitation, and herbivory (herbivore access/exclusion), including all relevant two and three-way interactions. We considered models with linear and quadratic forms for the influence of precipitation, to account for the possibility of non-monotonic responses. Although we evaluated quadratic models, LOOCV showed that they provided minimal improvement over monotonic models (Table S3). Therefore, all subsequent analyses used the monotonic model, which captures the observed patterns while maintaining model simplicity. We present the quadratic model here (Eq. 1) because it encompasses the linear model as a special case.

For each vital rate model, the expected value for the i^{th} observation (μ_i) was given by:

$$\begin{aligned}
\mu_i = & \beta_0[Sp_i] + \beta_P[Sp_i]precip_i + \beta_{P^2}[Sp_i]precip_i^2 + \beta_E[Sp_i]endo_i + \beta_H[Sp_i]herb_i \\
& + \beta_{EP}[Sp_i]endo_iprecip_i + \beta_{HP}[Sp_i]herb_iprecip_i \\
& + \beta_{HE}[Sp_i]herb_iendo_i + \beta_{HEP}[Sp_i]endo_iherb_iprecip_i \\
& + \phi_{site[i],Sp_i} + \omega_{source[i],Sp_i} + \rho_{plot[i],Sp_i}
\end{aligned} \tag{1}$$

$$\begin{aligned}
\phi_{site[i],Sp_i} & \sim N(0, \sigma_{site}^2) \\
\rho_{plot[i],Sp_i} & \sim N(0, \sigma_{plot}^2) \\
\omega_{source[i],Sp_i} & \sim N(0, \sigma_{source}^2)
\end{aligned}$$

Here, $precip_i$ is total precipitation (mm) accumulated over the 12 months preceding the census, $endo_i$ indicates endophyte presence ($endo_i = 1$) or absence ($endo_i = 0$), and $herb_i$ indicates fencing treatment: partial access ($herb_i = 0$) or full exclusion ($herb_i = 1$). The model includes random effects account for background variation at multiple hierarchical levels: $\phi_{site[i],Sp_i}$ captures site-to-site heterogeneity that is not explained by precipitation, $\rho_{plot[i],Sp_i}$ captures plot-to-plot variation within sites, and $\omega_{source[i],Sp_i}$ captures variation associated with the provenance of transplants. Random effect for site, plot, and source are host species-specific but the variance parameters (σ_{site} , σ_{source} , σ_{plot}) are shared across species; this assumes, for example, that site-to-site heterogeneity is of a similar magnitude across species, even as the model allows for “better” and “worse” sites to differ by species. This hierarchical structure allows the model to “borrow strength” across species, improving estimates of fixed effects while still capturing species-specific responses to environmental gradient.

All models were implemented in R (R Core Team, 2023) using Stan via the `rstan` interface (Stan Development Team, 2024). Models were run with 4 chains and sufficient iterations to ensure convergence, with weakly informative priors specified for all fixed effects. Full details on priors and model settings are provided in the accompanying code.

We confirmed that models effectively captured key aspects of the grass demography data, including means, standard deviations, and quantiles using a posterior predictive check (Fig. S2).

Next, for each vital rate, we compared candidate models with a linear or quadratic influence of precipitation using leave-one-out cross-validation (LOOCV) to identify the best-supported model (Vehtari et al., 2017). LOOCV combines validation and training by iteratively holding out one observation as a test set while fitting the model on the remaining $n - 1$ observations. This process is repeated n times, once for each observation, resulting in n fitted models (Silva and Zanella, 2024).

Drivers of endophyte effects on host demography

We used posterior parameter estimates to visualize predictions for how effects of symbiosis varied across abiotic and biotic contexts. For each herbivory treatment (Access or Exclusion) at each level of precipitation, we computed posterior predictions of survival probability E^+ and E^- hosts using the Bayesian model outputs (Eq. 1). The difference ($\Delta = E^+ - E^-$) was calculated for each posterior sample to estimate median effects and 90% credible intervals. We also computed the posterior probability that $\Delta > 0$, representing the likelihood that endophytes confer benefits on vital rates under the given precipitation and herbivory conditions.

To quantify statistical support for abiotic and/or biotic contexts as modifiers of grass-endophyte symbiosis, we fitted three hierarchical Bayesian candidate models for each vital rate (survival, growth, inflorescence, and spikelet production; Supplementary Methods S.1.1). All models included endophyte status as a baseline predictor but differed in whether they incorporated interactions with precipitation (Endophyte $\times$ Climate), herbivory (Endophyte $\times$ Herbivory), or both factors, including the three-way interaction (Endophyte $\times$ Climate $\times$ Herbivory). We compared candidate models using leave-one-out cross-validation (LOO-CV) (Vehtari et al., 2017), which estimates predictive accuracy by iteratively holding out each observation. Differences in expected log predictive density (ΔELPD), along with associated standard errors, were used to rank model performance. When the difference in ELPD between two models (ΔELPD) was small relative to its standard error (SE), we considered their predictive performance to be effectively similar. Accordingly, model rankings were interpreted cautiously, focusing on consistent trends across vital rates rather than on small differences between closely ranked models.

Results

Endophyte effects on host fitness components across abiotic and biotic stress gradients

Wetter conditions promoted grass host survival for all species, and abiotic stress modified endophyte effects on survival for two of three host species. Grass-*Epichloë* symbiosis had mostly negative or neutral effects on survival at low precipitation, but increasingly enhanced survival under wetter conditions (Fig. 3; Fig. S3).

In *Agrostis hyemalis* and *Elymus virginicus* populations with herbivore access, median $\Delta(E^+ - E^-)$ ranged from approximately -0.07 to -0.01 under low precipitation ($\Pr(\Delta > 0) = 0.15\text{--}0.42$) and increased to $0.06\text{--}0.15$ at the highest precipitation levels

($\Pr(\Delta > 0) = 0.78\text{--}0.91$; Table S4). Under herbivore exclusion, endophyte effects on survival were generally weaker and more variable: in *Agrostis*, median effects remained near zero across most of the precipitation gradient, becoming weakly positive only at the highest precipitation levels, whereas in *Elymus*, strongly negative effects were evident under dry conditions but shifted toward positive values under wetter conditions. *Poa autumnalis* showed little evidence of survival benefits associated with endophyte presence under herbivore access across the precipitation gradient (median $\Delta = -0.05$ to 0.00 , $\Pr(\Delta > 0) = 0.12\text{--}0.53$). Under herbivore exclusion, effects were weakly positive across precipitation levels (median $\Delta = 0.00\text{--}0.03$), but credible intervals consistently overlapped zero, indicating limited support for strong survival benefits.

Epichloë endophyte presence influenced relative growth in all three species, but effects depended on herbivore access and species identity (Table S5). In *A. hyemalis*, growth effects were neutral to weakly negative under dry conditions and became increasingly positive at moderate-to-high precipitation when herbivores had access (median $\Delta = 0.27\text{--}0.62$; $\Pr(\Delta > 0) = 0.94\text{--}0.99$). Under herbivore exclusion, growth benefits were present but consistently small (median $\Delta = 0.08\text{--}0.41$), despite a high probability of positive effects ($\Pr(\Delta > 0) = 0.71\text{--}0.96$). *E. virginicus* exhibited weakly positive growth effects across the precipitation gradient under herbivore access, with median Δ values increasing modestly from dry to wet conditions ($\Pr(\Delta > 0) = 0.52\text{--}0.96$). Under herbivore exclusion, growth responses were consistently small and uncertain, with median effects close to zero and credible intervals broadly overlapping zero. In *P. autumnalis*, the effect of endophyte presence on growth switched from negative (growth suppression) to positive (growth enhancement) across precipitation gradients, depending on herbivore access. Under herbivore access, median effects were negative under low precipitation and shifted toward positive values under wetter conditions, although uncertainty remained high. Under herbivore exclusion, growth benefits were strongest at low precipitation (median $\Delta = 0.56\text{--}0.81$; $\Pr(\Delta > 0) = 0.97\text{--}0.98$) but declined and became negative at the highest precipitation levels (median $\Delta = -0.41$; $\Pr(\Delta > 0) = 0.03$).

Epichloë endophytes affected reproductive output in a fitness-component- and species-specific manner. Inflorescence production showed little evidence of consistent endophyte benefits in *Agrostis hyemalis* or *Elymus virginicus*, with median effects near zero across precipitation levels and substantial uncertainty (Fig. 5). By contrast, *Poa autumnalis* exhibited strong and consistently positive effects of endophyte presence on inflorescence production, particularly under herbivore exclusion, where posterior support for positive effects was high across the precipitation gradient (Table S6). Spikelet production showed a broadly similar pattern. In *Elymus virginicus* and *Poa autumnalis*, endophyte effects were weak or negative under low precipitation but shifted toward positive values at higher precipitation, with benefits generally more pronounced under herbivore exclusion (Fig. S6; Fig. S7; Table S7).

Nevertheless, credible intervals frequently overlapped zero at low to moderate precipitation levels, indicating that strong reproductive benefits were largely restricted to wetter conditions.

Abiotic factors as the primary drivers of fitness components across the stress gradient

The outcomes of grass-endophyte symbioses whether beneficial, costly, or neutral were primarily associated with climatic variation rather than herbivore pressure. Across all vital rates (survival, growth, inflorescence, and spikelet production), model comparisons consistently indicated that models including an *Endophyte* $\times$ *Climate* interaction had equal or greater predictive support than alternatives emphasizing herbivory (Table 1). Although differences in expected log predictive density (ELPD) were modest relative to their standard errors, patterns were consistent across fitness components. For survival, the *Endophyte* $\times$ *Climate* model showed slightly higher predictive performance than the *Endophyte* $\times$ *Herbivory* model ($\Delta\text{ELPD} = -2.1$, $\text{SE} = 2.8$), whereas inclusion of the three-way *Endophyte* $\times$ *Climate* $\times$ *Herbivory* interaction reduced predictive performance ($\Delta\text{ELPD} = -5.0$, $\text{SE} = 1.9$).

Discussion

Drawing on three years of field experiments and using Bayesian statistical models, we evaluated how biotic and abiotic stressors jointly shape the demographic effects of vertically transmitted *Epichloë* endophytes in three cool-season perennial grasses across a natural climatic gradient in the United States. Our results indicate that grass-endophyte symbioses generally enhance host performance under favorable (e.g. wetter) conditions, whereas their effects can be neutral or even detrimental under high abiotic stress, consistent with the inhibition scenario. Across species and fitness components, climatic variation emerged as the dominant driver of endophyte-mediated effects, while herbivory had a secondary or context-dependent influence. These results underscore that the outcome of grass-*Epichloë* symbiosis shifts along abiotic stress gradients (Bronstein, 1994; Saikkonen et al., 1998; ?), providing empirical support for environmentally contingent mutualism in a broadly distributed host-symbiont system.⁸

Endophyte presence often enhances host performance under **favorable or moderately stressful**⁹ conditions, a pattern widely documented across grass-*Epichloë* systems (?). Endophytes can increase host growth, reproduction, and resistance to herbivores (?), but

⁸This paragraph is good but I think it needs to further emphasize that the influence of abiotic conditions is the opposite of what we predicted, and of what is expected under the SGH.

⁹unsure what point you are trying to make by combining favorable and 'moderately stressful' here. Also, I think it is important in the Discussion to distinguish between biotic and abiotic stress, since that is a theme of the paper.

these benefits are most apparent when the host has sufficient resources, such as water and nutrients, to support the fungus (??). Mechanistically, the host must allocate carbon and nutrients to maintain the endophyte, so under resource-rich conditions, the costs of symbiosis are outweighed by the benefits, resulting in a net positive effect. In our field experiment, we observed higher growth and survival of endophyte-infected grasses in wetter sites, consistent with previously documented positive effects of endophytes under favorable conditions (Afkhami et al., 2014; Rudgers et al., 2020).¹⁰

While endophyte infection enhanced host performance under favorable or moderately stressful conditions¹¹, its effects diminished and ultimately became costly under high abiotic stress, even in fenced plots where herbivores were excluded.¹² This pattern indicates that drought-driven resource limitation can impose costs on the symbiosis independent of biotic stress (Bronstein, 2001; Johnson et al., 1997; ?). By contrast, other studies show that the context-dependent benefits of endophytes are often mediated by herbivore pressure (Bastias et al., 2017; ?). For instance, in *Bromus setifolia*, endophyte infection is favored under high herbivore pressure, when defensive benefits outweigh the costs, but is selected against when herbivory is low or other costs are elevated (?). Our results suggest that similar context dependence can arise from abiotic factors, such as drought, independent of herbivory. Under low-precipitation conditions, reduced water availability constrains nutrient uptake and increases the relative metabolic burden of maintaining the endophyte, potentially tipping the cost-benefit balance against endophyte-symbiotic plants (Ahlholm et al., 2002; Bronstein, 2001; ?). In addition, severe environmental stress can activate plant defense pathways-such as elevated phytohormone production and reactive oxygen species (ROS) that may inhibit endophyte growth or reduce its functional contributions to the host (??). Consequently, even in the absence of herbivores, endophyte-mediated benefits may become negative in resource-limited environments, consistent with the reduced performance of infected plants at the driest sites in our study. Although we found no evidence of symbiosis-related costs for inflorescences across the precipitation gradient, spikelet production showed a small but consistent cost at the driest sites.¹³ This indicates that drought-driven resource limitation disproportionately affects certain reproductive components in infected plants, with different traits varying in their sensitivity to shifts in host-endophyte interactions (?).¹⁴

¹⁰I am a little confused here because I thought the Intro was meant to set up the idea that stress can strengthen benefits of the symbiosis.

¹¹Unclear what you mean by this.

¹²what's the relevance of this emphasis?

¹³Unclear what result this refers to.

¹⁴This also relates to the idea of demographic compensation, ie different vital rates responding differently to abiotic drivers - this may be worth bringing up in the Intro as a motivation for considering multiple vital rates side by side.

Across species and fitness components, climatic variation was the primary driver of endophyte-mediated effects, whereas herbivory played a secondary or **context-dependent role**.¹⁵ Precipitation strongly shaped host demography in our system, with drier sites constraining survival, growth and fertility. **Contrary to the predictions of the stress gradient hypothesis (SGH), which suggests that biotic interactions should become more positive under stressful conditions, endophyte-mediated benefits were reduced under high abiotic stress (Louthan et al., 2015; ?).**¹⁶ Although most empirical tests of the SGH have focused on plant–plant interactions and have been developed and tested mainly by community ecologists (Bertness and Callaway, 1994), recent syntheses show that support for the SGH is actually strongest among microbial interactions (??), with **bacterial systems**¹⁷ showing greater conformity to SGH predictions than any other **kingdom**¹⁸ (?). **This contrast highlights that plant–microbe mutualisms, including fungal endophytes, may not follow the same stress-dependency patterns documented in plant community ecology, and that endophyte benefits can decline rather than intensify under severe environmental stress.**¹⁹ A possible explanation for this opposite pattern is that the benefits of fungal endophytes are closely tied to host resource availability. Under high abiotic stress, host plants have limited resources for survival, growth, and reproduction, which can reduce the capacity of endophytes to confer benefits. Unlike plant–plant interactions, where neighbors can alleviate stress through mechanisms like shading or nutrient modification, endophytes depend on the host plant’s internal physiological condition, such as its nutrient and water status, energy reserves, and overall health. When host resources are constrained, the costs of maintaining the symbiosis may outweigh its benefits, leading to reduced endophyte-mediated effects under high abiotic stress.²⁰

¹⁵*Invoking context dependence here is a little “meta” since herbivory is itself context. So you are saying that the context dependence is context dependent! Also, I think we need to address the difference between fencing and herbivory. The herbivore effects was not consistent across sites because herbivore pressure varied, so the fence effect is dependent on baseline herbivory.*

¹⁶*This seems to me like the most important result that I expected would be more central to the first paragraph of the Intro.*

¹⁷*Does this mean bacterial symbionts of plants?*

¹⁸*Hard to know how to interpret this since interactions can involved multiple kingdoms.*

¹⁹*Important point, and I think you should bring in more of the literature documenting that E+ advantage increases in drier in environments, since our results show the opposite – we will then need to interpret why we get the opposite.*

²⁰*I like all the ideas you have here, just think we need to address more directly how our results fit with or contradict other results in the literature, including grass-endo systems that show E+ advantage in drought. It is also striking to me that the Discussion never returns to the predictions you establish in the conceptual figure. If you want to keep the conceptual figure then the Discussion needs to revisit it.*

References

- Afkhami, M. E., McIntyre, P. J., and Strauss, S. Y. (2014). Mutualist-mediated effects on species' range limits across large geographic scales. *Ecology letters*, 17(10):1265–1273.
- Afkhami, M. E. and Rudgers, J. A. (2008). Symbiosis lost: imperfect vertical transmission of fungal endophytes in grasses. *The American Naturalist*, 172(3):405–416.
- Ahlholm, J. U., Helander, M., Lehtimäki, S., Wäli, P., and Saikkonen, K. (2002). Vertically transmitted fungal endophytes: different responses of host-parasite systems to environmental conditions. *Oikos*, 99(1):173–183.
- Amijee, F., Tinker, P., and Stribley, D. (1989). The development of endomycorrhizal root systems: Vii. a detailed study of effects of soil phosphorus on colonization. *New Phytologist*, 111(3):435–446.
- Atala, C., Acuña-Rodríguez, I. S., Torres-Díaz, C., and Molina-Montenegro, M. A. (2022). Fungal endophytes improve the performance of host plants but do not eliminate the growth/defence trade-off. *New Phytologist*, 235(2).
- Bastias, D. A., Martínez-Ghersa, M. A., Ballaré, C. L., and Gundel, P. E. (2017). Epichloë fungal endophytes and plant defenses: not just alkaloids. *Trends in Plant Science*, 22(11):939–948.
- Bertness, M. D. and Callaway, R. (1994). Positive interactions in communities. *Trends in ecology & evolution*, 9(5):191–193.
- Bronstein, J. L. (1994). Conditional outcomes in mutualistic interactions. *Trends in ecology & evolution*, 9(6):214–217.
- Bronstein, J. L. (2001). The exploitation of mutualisms. *Ecology letters*, 4(3):277–287.
- Clay, K., Holah, J., and Rudgers, J. A. (2005). Herbivores cause a rapid increase in hereditary symbiosis and alter plant community composition. *Proceedings of the National Academy of Sciences*, 102(35):12465–12470.
- Clay, K. and Schardl, C. (2002). Evolutionary origins and ecological consequences of endophyte symbiosis with grasses. *the american naturalist*, 160(S4):S99–S127.
- Decunta, F. A., Pérez, L. I., Malinowski, D. P., Molina-Montenegro, M. A., and Gundel, P. E. (2021). A systematic review on the effects of epichloë fungal endophytes on drought tolerance in cool-season grasses. *Frontiers in Plant Science*, 12:644731.

- Emery, S. M., Thompson, D., and Rudgers, J. A. (2010). Variation in endophyte symbiosis, herbivory and drought tolerance of *ammophila breviligulata* populations in the great lakes region. *The American Midland Naturalist*, 163(1):186–196.
- Fowler, J. C., Donald, M. L., Bronstein, J. L., and Miller, T. E. (2023). The geographic footprint of mutualism: How mutualists influence species’ range limits. *Ecological Monographs*, 93(1):e1558.
- Fowler, J. C., Moutouama, J., and Miller, T. E. (2025). Increasing prevalence of plant-fungal symbiosis across two centuries of environmental change. *Global Change Biology*, 31(11):e70577.
- GBIF (2025a). Occurrence download for *Agrostis hyemalis*. Accessed 5 November 2025.
- GBIF (2025b). Occurrence download for *Elymus virginicus*. Accessed 5 November 2025.
- GBIF (2025c). Occurrence download for *Poa autumnalis*. Accessed 5 November 2025.
- Giauque, H., Connor, E. W., and Hawkes, C. V. (2019). Endophyte traits relevant to stress tolerance, resource use and habitat of origin predict effects on host plants. *New Phytologist*, 221(4):2239–2249.
- Gundel, P. E., Rudgers, J. A., and Ghera, C. M. (2011). Incorporating the process of vertical transmission into understanding of host–symbiont dynamics. *Oikos*, 120(8):1121–1128.
- Gundel, P. E., Sun, P., Charlton, N. D., Young, C. A., Miller, T. E., and Rudgers, J. A. (2020). Simulated folivory increases vertical transmission of fungal endophytes that deter herbivores and alter tolerance to herbivory in *poa autumnalis*. *Annals of Botany*, 125(6):981–991.
- Gundel, P. E., Ueno, A. C., Casas, C., Miller, T. E., Pérez, L. I., Cuyeu, R., and Omacini, M. (2024). Temporal host-symbiont dynamics in community contexts: Impacts of host fitness and vertical transmission efficiency on symbiosis prevalence. *Functional Ecology*, 38(12):2610–2622.
- Hart, E. M. and Bell, K. (2015). *prism: Download data from the Oregon prism project*. R package version 0.0.6.
- Hoang, K. L., Salguero-Gómez, R., Pike, V. L., and King, K. C. (2024). The impacts of host association and perturbation on symbiont fitness. *Symbiosis*, 92(3):439–451.
- Hoeksema, J. D. and Bruna, E. M. (2015). Context-dependent outcomes of mutualistic interactions. *Mutualism*, 10:181–202.

- Holmgren, M. and Scheffer, M. (2010). Strong facilitation in mild environments: the stress gradient hypothesis revisited. *Journal of Ecology*, 98(6):1269–1275.
- Johnson, N. C., Graham, J. H., and Smith, F. A. (1997). Functioning of mycorrhizal associations along the mutualism–parasitism continuum. *The New Phytologist*, 135(4):575–585.
- Klironomos, J. N. (2003). Variation in plant response to native and exotic arbuscular mycorrhizal fungi. *Ecology*, 84(9):2292–2301.
- Leuchtmann, A., Bacon, C. W., Schardl, C. L., White Jr, J. F., and Tadych, M. (2014). Nomenclatural realignment of neotyphodium species with genus epichloë. *Mycologia*, 106(2):202–215.
- Liu, M. (2023). Fungi canadenses no. 351: Epichloe elymi. *Canadian Journal of Plant Pathology*, 45(3):300–303.
- Louthan, A. M., Doak, D. F., and Angert, A. L. (2015). Where and when do species interactions set range limits? *Trends in ecology & evolution*, 30(12):780–792.
- Oberhofer, M., Güsewell, S., and Leuchtmann, A. (2014). Effects of natural hybrid and non-hybrid epichloë endophytes on the response of hordelymus europaeus to drought stress. *New Phytologist*, 201(1):242–253.
- Porter, S. S., Bantay, R., Friel, C. A., Garoutte, A., Gdanetz, K., Ibarreta, K., Moore, B. M., Shetty, P., Siler, E., and Friesen, M. L. (2020). Beneficial microbes ameliorate abiotic and biotic sources of stress on plants. *Functional Ecology*, 34(10):2075–2086.
- R Core Team (2023). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rudgers, J. A., Afkhami, M. E., Bell-Dereske, L., Chung, Y. A., Crawford, K. M., Kivlin, S. N., Mann, M. A., and Nuñez, M. A. (2020). Climate disruption of plant-microbe interactions. *Annual review of ecology, evolution, and systematics*, 51:561–586.
- Rudgers, J. A., Fletcher, R. A., Olivas, E., Young, C. A., Charlton, N. D., Pearson, D. E., and Maron, J. L. (2016). Long-term ungulate exclusion reduces fungal symbiont prevalence in native grasslands. *Oecologia*, 181(4):1151–1161.
- Rudgers, J. A., Miller, T. E., Ziegler, S. M., and Craven, K. D. (2012). There are many ways to be a mutualist: endophytic fungus reduces plant survival but increases population growth. *Ecology*, 93(3):565–574.

- 469 Ruiz-lazona, J. M., Azcón, R., and Palma, J. (1996). Superoxide dismutase activity in arbuscular
470 mycorrhizal lactuca sativa plants subjected to drought stress. *New Phytologist*, 134(2):327–333.
- 471 Saikkonen, K., Faeth, S. H., Helander, M., and Sullivan, T. (1998). Fungal endophytes: a contin-
472 uum of interactions with host plants. *Annual review of Ecology and Systematics*, 29(1):319–343.
- 473 Saikkonen, K., Saari, S., and Helander, M. (2010). Defensive mutualism between plants and
474 endophytic fungi? *Fungal Diversity*, 41(1):101–113.
- 475 Saikkonen, K., Wäli, P., Helander, M., and Faeth, S. H. (2004). Evolution of endophyte–plant
476 symbioses. *Trends in plant science*, 9(6):275–280.
- 477 Schardl, C. L., Leuchtmann, A., and Spiering, M. J. (2004). Symbioses of grasses with
478 seedborne fungal endophytes. *Annu. Rev. Plant Biol.*, 55(1):315–340.
- 479 Shaw, R. B. (2011). *Guide to Texas grasses*. Texas A&M University Press.
- 480 Silva, L. A. and Zanella, G. (2024). Robust leave-one-out cross-validation for high-dimensional
481 bayesian models. *Journal of the American Statistical Association*, 119(547):2369–2381.
- 482 Soudzilovskaia, N. A., van der Heijden, M. G., Cornelissen, J. H., Makarov, M. I., Onipchenko,
483 V. G., Maslov, M. N., Akhmetzhanova, A. A., and van Bodegom, P. M. (2015). Quantitative
484 assessment of the differential impacts of arbuscular and ectomycorrhiza on soil carbon
485 cycling. *New Phytologist*, 208(1):280–293.
- 486 Stan Development Team (2024). RStan: the R interface to Stan. R package version 2.32.6.
- 487 Thirkell, T. J., Pastok, D., and Field, K. J. (2020). Carbon for nutrient exchange between
488 arbuscular mycorrhizal fungi and wheat varies according to cultivar and changes in
489 atmospheric carbon dioxide concentration. *Global change biology*, 26(3):1725–1738.
- 490 Vega, F. E. (2008). Insect pathology and fungal endophytes. *Journal of invertebrate pathology*,
491 98(3):277–279.
- 492 Vehtari, A., Gelman, A., and Gabry, J. (2017). Practical bayesian model evaluation using
493 leave-one-out cross-validation and waic. *Statistics and computing*, 27:1413–1432.
- 494 White Jr, J. F. (1994). Endophyte-host associations in grasses. xx. structural and reproductive
495 studies of epichloë amarillans sp. nov. and comparisons to e. typhina. *Mycologia*,
496 86(4):571–580.
- 497 Yule, K. M., Miller, T. E., and Rudgers, J. A. (2013). Costs, benefits, and loss of vertically
498 transmitted symbionts affect host population dynamics. *Oikos*, 122(10):1512–1520.

Figures




Figure 1: Conceptual framework illustrating the predicted effects of abiotic stress, biotic stress, and endophyte-mediated mutualism on host plant fitness (adapted from (Porter et al., 2020)). (a) **Endophyte-driven stress amelioration scenario:** Endophytes are predicted to enhance host performance under increasing abiotic stress (e.g., low precipitation) in the absence of herbivores by mitigating drought-related physiological costs. (b) **Reduced endophyte-mediated benefits scenario:** When abiotic stress is combined with herbivory, the positive effect of endophytes is expected to weaken. Limited plant resources are divided between defense against herbivores and sustaining the endophyte, reducing the endophyte's capacity to ameliorate drought stress and potentially resulting in a net cost of hosting an endophyte. Colors are used solely for visualization: blue lines represent endophyte-infected plants and violet lines represent endophyte-free plants.




Figure 2: Distribution of common garden sites along a natural aridity gradient from Texas to Louisiana, US for (A) *Agrostis hyemalis*, (B) *Elymus virginicus*, and (C) *Poa autumnalis*. Climate data in A–C are based on 30-year normals of climate predictions (1996–2025) from PRISM data (Hart and Bell, 2015), illustrating the precipitation gradient across experimental sites (scale on the right is in mm). Red dots indicate the locations of source populations, while grey dots represent Global Biodiversity Information Facility (GBIF) occurrence records for each species within the study area (GBIF, 2025a,b,c). (D) Annual cumulative precipitation during each census year (May 2023 – June 2025) at each experimental site, ordered west to east. Values are also derived from PRISM climate data. (E) Schematic illustration of fenced (left) and unfenced (right) plots used to assess herbivory. Black squares represent individual plants; plot shape correspond to experimental design. (F) Proportion of damaged plants in fenced (salmon) and unfenced (light green) plots across experimental sites, with mean ± standard error. Sites are ordered west to east as in panel D. Panels E–F illustrate herbivory treatments and outcomes, and species-specific breakdowns are provided in Supplementary Figures.




Figure 3: Predicted survival rate (top panels) and endophyte effect ($\Delta = E^+ - E^-$, bottom panels) across precipitation gradients. Lines show posterior means for E^+ (blue) and E^- (red), with shading indicating 90% credible intervals. Points represent mean observed survival per plot, jittered for visualization. The dashed line at $\Delta = 0$ indicates no endophyte effect. Panels are organized by species and herbivory treatment (Fenced, Unfenced).




Figure 4: Predicted relative growth from year to the next year (top panels) and endophyte effect ($\Delta = E^+ - E^-$, bottom panels) across precipitation gradients. Lines show posterior means for E⁺ (blue) and E⁻ (red), with shading indicating 90% credible intervals. Points represent mean observed growth per plot, jittered for visualization. The dashed line at $\Delta = 0$ indicates no endophyte effect. Panels are organized by species and herbivory treatment (Fenced, Unfenced).




Figure 5: Predicted number of inflorescences (upper panels) and the difference between endophyte-infected (E^+) and endophyte-free (E^-) plants ($\Delta(E^+ - E^-)$, lower panels) are shown across a gradient of precipitation. Shaded areas represent 90% credible intervals. Points indicate observed means, jittered for visualization. Panels are grouped by species and herbivory treatment (Fenced, Unfenced).




Figure 6: Predicted effects of herbivore exclusion on survival, growth, and inflorescence production across precipitation gradients for *A. hyemalis*, *E. virginicus*, and *P. autumnalis*. Panels show the median difference between fenced and unfenced treatments ($\Delta = E^+ - E^-$) and the probability that this difference is positive ($\Pr[\Delta > 0]$). Horizontal dashed lines indicate $\Delta = 0$ and $\Pr(\Delta > 0) = 0.5$, and vertical dashed lines denote species-specific mean annual precipitation at range limits. Colors indicate herbivore exclusion treatments.

Table 1: Model comparison of drivers of grass–endophyte symbiosis across four fitness components (vital rates). Δ ELPD indicates the difference in expected log predictive density relative to the best-fitting model; SE is the standard error of the difference.

Vital rate	Model type	Δ ELPD	SE
Survival	Endophyte $\times$ Climate	0.00	0.00
	Endophyte $\times$ Herbivory	−4.65	2.05
	Endophyte $\times$ Climate $\times$ Herbivory	−5.23	3.82
Growth	Endophyte $\times$ Climate	0.00	0.00
	Endophyte $\times$ Herbivory	−0.27	3.24
	Endophyte $\times$ Climate $\times$ Herbivory	−0.31	3.51
Inflorescence	Endophyte $\times$ Climate	0.00	0.00
	Endophyte $\times$ Herbivory	−0.74	2.70
	Endophyte $\times$ Climate $\times$ Herbivory	−0.90	3.39
Spikelet	Endophyte $\times$ Climate	0.00	0.00
	Endophyte $\times$ Herbivory	−0.92	3.31
	Endophyte $\times$ Climate $\times$ Herbivory	−3.27	2.44

S.1 Supporting Information

S.1.1 Supporting Methods

Candidate models for abiotic and biotic drivers of cool-season grass species

1. **Endophyte × Climate model:** includes endophyte status (endo), precipitation (P), and their interaction:

$$\begin{aligned}\mu_i = & \beta_0[\text{Sp}_i] + \beta_{\text{endo}}[\text{Sp}_i]\text{endo}_i + \beta_P[\text{Sp}_i]P_i + \beta_{\text{endo} \times P}[\text{Sp}_i]\text{endo}_iP_i \\ & + \phi_{\text{site}[i], \text{Sp}_i} + \omega_{\text{source}[i], \text{Sp}_i} + \rho_{\text{plot}[i], \text{Sp}_i}\end{aligned}$$

2. **Endophyte × Herbivory model:** includes endophyte status (endo), herbivory (herb), and their interaction:

$$\begin{aligned}\mu_i = & \beta_0[\text{Sp}_i] + \beta_{\text{endo}}[\text{Sp}_i]\text{endo}_i + \beta_{\text{herb}}[\text{Sp}_i]\text{herb}_i + \beta_{\text{endo} \times \text{herb}}[\text{Sp}_i]\text{endo}_i\text{herb}_i \\ & + \phi_{\text{site}[i], \text{Sp}_i} + \omega_{\text{source}[i], \text{Sp}_i} + \rho_{\text{plot}[i], \text{Sp}_i}\end{aligned}$$

3. **Endophyte × Climate × Herbivory:** combines all predictors and interactions up to the three-way interaction:

$$\begin{aligned}\mu_i = & \beta_0[\text{Sp}_i] + \beta_P[\text{Sp}_i]P_i + \beta_{\text{endo}}[\text{Sp}_i]\text{endo}_i + \beta_{\text{herb}}[\text{Sp}_i]\text{herb}_i \\ & + \beta_{\text{endo} \times P}[\text{Sp}_i]\text{endo}_iP_i + \beta_{\text{herb} \times P}[\text{Sp}_i]\text{herb}_iP_i + \beta_{\text{herb} \times \text{endo}}[\text{Sp}_i]\text{herb}_i\text{endo}_i \\ & + \beta_{\text{endo} \times \text{herb} \times P}[\text{Sp}_i]\text{endo}_i\text{herb}_iP_i + \phi_{\text{site}[i], \text{Sp}_i} + \omega_{\text{source}[i], \text{Sp}_i} + \rho_{\text{plot}[i], \text{Sp}_i}\end{aligned}$$

In these models, the β coefficients represent species-specific effects. The intercept $\beta_0[\text{Sp}_i]$ corresponds to the baseline value of the vital rate for species Sp_i in the absence of predictors. $\beta_{\text{endo}}[\text{Sp}_i]$ and $\beta_{\text{herb}}[\text{Sp}_i]$ describe the effects of endophyte presence and herbivory, respectively. The linear effect of precipitation is captured by $\beta_P[\text{Sp}_i]$, while interactions are represented by $\beta_{\text{endo} \times P}[\text{Sp}_i]$, $\beta_{\text{herb} \times P}[\text{Sp}_i]$, $\beta_{\text{endo} \times \text{herb}}[\text{Sp}_i]$, and the three-way interaction $\beta_{\text{endo} \times \text{herb} \times P}[\text{Sp}_i]$. All models include hierarchical random effects for site-year, source population, and plot, with species-specific random effects $\phi_{\text{site}[i], \text{Sp}_i} \sim \mathcal{N}(0, \sigma_{\text{site}})$, $\omega_{\text{source}[i], \text{Sp}_i} \sim \mathcal{N}(0, \sigma_{\text{source}})$, and $\rho_{\text{plot}[i], \text{Sp}_i} \sim \mathcal{N}(0, \sigma_{\text{plot}})$.

S.1.2 Supporting Tables

Table S1: Geographic coordinates and brief descriptions of source populations for *Agrostis hyemalis* (AGHY), *Elymus virginicus* (ELVI), and *Poa autumnalis* (POAU).

Species	Population	Latitude	Longitude	Description
AGHY	SJC	30.771969	-95.200620	San Jacinto County, TX
AGHY	COHR	29.672833	-96.567517	Columbus, TX
AGHY	LPEF	30.085383	-97.172508	Lost Pines
POAU	LA7	30.849660	-93.276772	DeRidder, LA
POAU	LA1	32.568125	-93.932976	Jean Lafitte National Historical Park and Preserve ²¹
POAU	LA2	31.183040	-92.616969	Jean Lafitte National Historical Park and Preserve
POAU	SFAEF	31.492158	-94.772200	Stephen F. Austin Experimental Forest, TX
POAU	WB	30.399346	-95.150026	Sam Houston National Forest
ELVI	HUNT	30.741797	-95.474508	Sam Houston State University Field Station
ELVI	SHS ²²	30.741797	-95.474508	Sam Houston State University Field Station
ELVI	RL5	30.471717	-99.783803	Texas Tech University Field Station, Junction
ELVI	PALM	29.591623	-97.584781	Palmetto State Park, TX
ELVI	JLP	29.785105	-90.114933	Jean Lafitte Park, outside of New Orleans

Table S2: Common garden sites used for reciprocal transplant and environmental manipulation studies.

Site	Code	Latitude	Longitude
University of Louisiana Ecology Center	LAF	30.304773	-92.012401
Center for Biological Field Studies	HUN	30.742815	-95.476881
TAMU Ecology and Natural Resource Teaching Area	COL	30.569302	-96.366324
UT Stengl Lost Pines Field Station	BAS	30.092343	-97.172442
1335 Medina Hwy, Kerrville, TX	KER	30.028437	-99.142888
Brackenridge Field Laboratory	BFL	30.284866	-97.780675
TAMU Sonora Station	SON	30.273467	-100.561463

Table S3: Comparison of candidate models (quadratic vs. linear) for each vital rate based on leave-one-out cross-validation (LOOCV). ELPD difference is relative to the best model.

Vital rate	Model type	ELPD difference	SE difference
Survival	Quadratic	0	0.00
Survival	Linear	-0.055	1.09
Growth	Quadratic	0	0.00
Growth	Linear	-0.187	1.33
Inflorescence	Quadratic	0	0.00
Inflorescence	Linear	-1.510	1.43
Spikelet	Quadratic	0	0.00
Spikelet	Linear	-0.849	1.32

Table S4: Posterior summaries of Δ ($E^+ - E^-$) for survival across precipitation quartiles for each species and herbivore exclusion treatment. Median, 90% credible interval, and posterior probability that $\Delta > 0$ are shown. Herbivore exclusion: Fenced = herbivores excluded, Unfenced = herbivores had access.

Species	Herbivore exclusion	Precipitation bin (mm)	Median Δ	Lower 90%	Upper 90%	$P(\Delta > 0)$
<i>Agrostis hyemalis</i>	Unfenced	493	-0.049	-0.302	0.053	0.158
<i>Agrostis hyemalis</i>	Unfenced	678	-0.038	-0.215	0.052	0.187
<i>Agrostis hyemalis</i>	Unfenced	1036	-0.012	-0.099	0.058	0.357
<i>Agrostis hyemalis</i>	Unfenced	1424	0.015	-0.050	0.102	0.681
<i>Agrostis hyemalis</i>	Unfenced	2294	0.056	-0.035	0.249	0.860
<i>Agrostis hyemalis</i>	Fenced	493	-0.090	-0.358	0.000	0.051
<i>Agrostis hyemalis</i>	Fenced	678	-0.076	-0.264	0.002	0.057
<i>Agrostis hyemalis</i>	Fenced	1036	-0.041	-0.142	0.019	0.127
<i>Agrostis hyemalis</i>	Fenced	1424	-0.004	-0.085	0.072	0.448
<i>Agrostis hyemalis</i>	Fenced	2294	0.047	-0.052	0.224	0.802
<i>Elymus virginicus</i>	Unfenced	493	-0.069	-0.321	0.050	0.145
<i>Elymus virginicus</i>	Unfenced	678	-0.065	-0.254	0.068	0.189
<i>Elymus virginicus</i>	Unfenced	1036	-0.014	-0.143	0.113	0.424
<i>Elymus virginicus</i>	Unfenced	1424	0.061	-0.069	0.194	0.779
<i>Elymus virginicus</i>	Unfenced	2294	0.148	-0.031	0.356	0.914
<i>Elymus virginicus</i>	Fenced	493	-0.259	-0.550	-0.058	0.006
<i>Elymus virginicus</i>	Fenced	678	-0.230	-0.430	-0.058	0.012
<i>Elymus virginicus</i>	Fenced	1036	-0.059	-0.183	0.061	0.210
<i>Elymus virginicus</i>	Fenced	1424	0.090	-0.022	0.225	0.907
<i>Elymus virginicus</i>	Fenced	2294	0.154	0.031	0.400	0.992
<i>Poa autumnalis</i>	Unfenced	493	-0.001	-0.101	0.005	0.175
<i>Poa autumnalis</i>	Unfenced	678	-0.005	-0.143	0.010	0.154
<i>Poa autumnalis</i>	Unfenced	1036	-0.043	-0.196	0.022	0.118
<i>Poa autumnalis</i>	Unfenced	1424	-0.051	-0.188	0.060	0.214
<i>Poa autumnalis</i>	Unfenced	2294	0.000	-0.124	0.137	0.532
<i>Poa autumnalis</i>	Fenced	493	0.000	-0.019	0.067	0.614
<i>Poa autumnalis</i>	Fenced	678	0.001	-0.033	0.092	0.641
<i>Poa autumnalis</i>	Fenced	1036	0.016	-0.058	0.136	0.689
<i>Poa autumnalis</i>	Fenced	1424	0.029	-0.088	0.166	0.679
<i>Poa autumnalis</i>	Fenced	2294	0.002	-0.117	0.151	0.570

Table S5: Posterior summaries of $\Delta (E^+ - E^-)$ for growth across precipitation quantiles for each species and herbivore treatment. Positive median values indicate beneficial effects of endophytes; negative medians indicate costly effects.

Species	Herbivore exclusion	Precipitation (mm)	Median Δ	Lower 90%	Upper 90%	P($\Delta > 0$)
<i>Agrostis hyemalis</i>	Unfenced	493	-0.046	-0.643	0.561	0.450
<i>Agrostis hyemalis</i>	Unfenced	678	0.092	-0.352	0.543	0.629
<i>Agrostis hyemalis</i>	Unfenced	1036	0.274	-0.019	0.571	0.939
<i>Agrostis hyemalis</i>	Unfenced	1424	0.412	0.147	0.683	0.994
<i>Agrostis hyemalis</i>	Unfenced	2294	0.616	0.212	1.022	0.994
<i>Agrostis hyemalis</i>	Fenced	493	-0.221	-0.723	0.290	0.237
<i>Agrostis hyemalis</i>	Fenced	678	-0.090	-0.466	0.284	0.347
<i>Agrostis hyemalis</i>	Fenced	1036	0.084	-0.165	0.336	0.710
<i>Agrostis hyemalis</i>	Fenced	1424	0.214	-0.034	0.460	0.925
<i>Agrostis hyemalis</i>	Fenced	2294	0.411	0.017	0.799	0.956
<i>Elymus virginicus</i>	Unfenced	493	0.017	-0.618	0.647	0.520
<i>Elymus virginicus</i>	Unfenced	678	0.084	-0.400	0.566	0.614
<i>Elymus virginicus</i>	Unfenced	1036	0.168	-0.137	0.476	0.814
<i>Elymus virginicus</i>	Unfenced	1424	0.232	-0.002	0.470	0.948
<i>Elymus virginicus</i>	Unfenced	2294	0.330	0.008	0.643	0.955
<i>Elymus virginicus</i>	Fenced	493	-0.002	-0.594	0.578	0.498
<i>Elymus virginicus</i>	Fenced	678	0.023	-0.419	0.457	0.532
<i>Elymus virginicus</i>	Fenced	1036	0.060	-0.205	0.316	0.651
<i>Elymus virginicus</i>	Fenced	1424	0.087	-0.101	0.274	0.777
<i>Elymus virginicus</i>	Fenced	2294	0.130	-0.160	0.410	0.767
<i>Poa autumnalis</i>	Unfenced	493	-0.393	-1.089	0.289	0.174
<i>Poa autumnalis</i>	Unfenced	678	-0.283	-0.803	0.230	0.185
<i>Poa autumnalis</i>	Unfenced	1036	-0.136	-0.448	0.179	0.242
<i>Poa autumnalis</i>	Unfenced	1424	-0.024	-0.245	0.202	0.431
<i>Poa autumnalis</i>	Unfenced	2294	0.143	-0.175	0.474	0.763
<i>Poa autumnalis</i>	Fenced	493	0.806	0.137	1.478	0.977
<i>Poa autumnalis</i>	Fenced	678	0.556	0.058	1.060	0.967
<i>Poa autumnalis</i>	Fenced	1036	0.223	-0.077	0.529	0.890
<i>Poa autumnalis</i>	Fenced	1424	-0.028	-0.251	0.199	0.418
<i>Poa autumnalis</i>	Fenced	2294	-0.406	-0.737	-0.067	0.025

Table S6: Posterior summaries of $\Delta (E^+ - E^-)$ for flowering output at precipitation quantiles for each species and herbivore exclusion treatment. Median, 90% credible interval, and posterior probability that $\Delta > 0$ are shown. Herbivore exclusion: Fenced = herbivores excluded, Unfenced = herbivores had access.

Species	Herbivore exclusion	Precipitation (mm)	Median Δ	Lower 90%	Upper 90%	P($\Delta > 0$)
<i>Agrostis hyemalis</i>	Unfenced	493	-0.325	-1.371	0.229	0.137
<i>Agrostis hyemalis</i>	Unfenced	678	-0.327	-1.230	0.320	0.172
<i>Agrostis hyemalis</i>	Unfenced	1036	-0.157	-0.929	0.557	0.344
<i>Agrostis hyemalis</i>	Unfenced	1424	0.231	-0.589	1.219	0.691
<i>Agrostis hyemalis</i>	Unfenced	2294	1.730	-0.289	6.251	0.914
<i>Agrostis hyemalis</i>	Fenced	493	-0.355	-1.550	0.311	0.154
<i>Agrostis hyemalis</i>	Fenced	678	-0.332	-1.339	0.426	0.206
<i>Agrostis hyemalis</i>	Fenced	1036	-0.067	-0.897	0.796	0.439
<i>Agrostis hyemalis</i>	Fenced	1424	0.481	-0.492	1.772	0.800
<i>Agrostis hyemalis</i>	Fenced	2294	2.394	-0.264	8.520	0.930
<i>Elymus virginicus</i>	Unfenced	493	-0.167	-0.742	0.249	0.220
<i>Elymus virginicus</i>	Unfenced	678	-0.142	-0.606	0.255	0.250
<i>Elymus virginicus</i>	Unfenced	1036	-0.062	-0.410	0.274	0.369
<i>Elymus virginicus</i>	Unfenced	1424	0.046	-0.277	0.391	0.606
<i>Elymus virginicus</i>	Unfenced	2294	0.320	-0.226	1.141	0.831
<i>Elymus virginicus</i>	Fenced	493	0.006	-1.001	1.434	0.505
<i>Elymus virginicus</i>	Fenced	678	0.015	-0.824	1.125	0.512
<i>Elymus virginicus</i>	Fenced	1036	0.033	-0.576	0.732	0.536
<i>Elymus virginicus</i>	Fenced	1424	0.044	-0.445	0.566	0.561
<i>Elymus virginicus</i>	Fenced	2294	0.064	-0.808	1.089	0.551
<i>Poa autumnalis</i>	Unfenced	493	0.029	-0.152	1.709	0.738
<i>Poa autumnalis</i>	Unfenced	678	0.070	-0.097	1.287	0.798
<i>Poa autumnalis</i>	Unfenced	1036	0.194	-0.029	1.143	0.921
<i>Poa autumnalis</i>	Unfenced	1424	0.385	0.054	1.767	0.987
<i>Poa autumnalis</i>	Unfenced	2294	0.957	0.027	10.518	0.972
<i>Poa autumnalis</i>	Fenced	493	0.543	0.021	14.821	0.996
<i>Poa autumnalis</i>	Fenced	678	0.744	0.072	8.201	0.998
<i>Poa autumnalis</i>	Fenced	1036	1.076	0.259	4.875	1.000
<i>Poa autumnalis</i>	Fenced	1424	1.349	0.322	5.461	1.000
<i>Poa autumnalis</i>	Fenced	2294	1.355	0.031	15.437	0.968

Table S7: Posterior summaries of Δ ($E^+ - E^-$) at precipitation quantiles for each species and herbivore exclusion treatment. Median, 90% credible interval, and posterior probability that $\Delta > 0$ are shown. Herbivore exclusion: Fenced = herbivores excluded, Unfenced = herbivores had access.

Species	Herbivore exclusion	Precipitation (mm)	Median Δ	Lower 90%	Upper 90%	$P(\Delta > 0)$
<i>Elymus virginicus</i>	Unfenced	493	-3.150	-11.100	3.190	0.202
<i>Elymus virginicus</i>	Unfenced	678	-2.380	-8.360	2.850	0.224
<i>Elymus virginicus</i>	Unfenced	1036	-1.130	-5.120	2.670	0.303
<i>Elymus virginicus</i>	Unfenced	1424	-0.095	-3.240	3.110	0.481
<i>Elymus virginicus</i>	Unfenced	2294	1.680	-2.130	6.290	0.770
<i>Elymus virginicus</i>	Fenced	493	-4.550	-11.900	1.240	0.092
<i>Elymus virginicus</i>	Fenced	678	-3.360	-8.640	1.310	0.111
<i>Elymus virginicus</i>	Fenced	1036	-1.440	-4.570	1.640	0.211
<i>Elymus virginicus</i>	Fenced	1424	0.261	-2.040	2.650	0.577
<i>Elymus virginicus</i>	Fenced	2294	3.230	-0.389	8.420	0.929
<i>Poa autumnalis</i>	Unfenced	493	-0.584	-6.080	3.140	0.320
<i>Poa autumnalis</i>	Unfenced	678	-0.584	-4.890	2.760	0.330
<i>Poa autumnalis</i>	Unfenced	1036	-0.408	-3.240	2.230	0.373
<i>Poa autumnalis</i>	Unfenced	1424	-0.025	-2.090	2.060	0.491
<i>Poa autumnalis</i>	Unfenced	2294	1.100	-2.360	5.460	0.719
<i>Poa autumnalis</i>	Fenced	493	2.630	-1.310	12.700	0.890
<i>Poa autumnalis</i>	Fenced	678	2.940	-0.645	10.500	0.917
<i>Poa autumnalis</i>	Fenced	1036	3.270	0.227	7.900	0.963
<i>Poa autumnalis</i>	Fenced	1424	3.330	0.797	6.550	0.987
<i>Poa autumnalis</i>	Fenced	2294	2.750	-1.340	8.540	0.871

S.1.3 Supporting Figures




Fig S1: Abiotic conditions across the seven study sites. (A) Mean annual precipitation showing the pronounced east–west aridity gradient from mesic Louisiana to arid Texas. (B) Mean annual temperature across sites, which does not differ much. Panels highlight that precipitation, rather than temperature, constitutes the primary abiotic gradient in this system.




Fig S2: Posterior predictive checks for each vital rate. Dark blue lines show observed data (y), and light blue lines show replicated datasets generated from posterior predictions (y_{rep}). The strong overlap across survival, growth, flowering, and spikelet production indicates adequate model fit for all vital rates.




Fig S3: Predicted effects of endophyte presence (E^+) versus absence (E^-) on survival across precipitation gradients for three cool-season grasses. Solid lines show the median $\Delta (E^+ - E^-)$, or the posterior probability that $\Delta > 0$ under fenced and unfenced conditions. Horizontal dashed lines indicate no effect ($\Delta = 0$) or a 50% probability of benefit. Vertical dashed lines represent the precipitation values at the most westerly natural occurrence of each species, based on GBIF records.




Fig S4: Effects of precipitation and endophyte presence on growth across three cool-season grass species. Lines show the predicted difference in growth between endophyte-present (E^+) and endophyte-absent (E^-) plants (Median Δ , top row) and the posterior probability that this difference is positive ($\text{Pr}(\Delta > 0)$), bottom row) under fenced and unfenced treatments. Predictions were calculated at the 10th, 25th, 50th, 75th, and 90th percentiles of precipitation observed in the study. Horizontal dashed lines indicate $\Delta = 0$ (top row) and $\text{Pr}(\Delta > 0) = 0.5$ (bottom row). Vertical dashed lines represent the precipitation values at the most westerly natural occurrence of each species, based on GBIF records.




Fig S5: Effects of endophyte infection on flowering output across a precipitation gradient for three cool-season grass species. Lines show the median posterior difference $\Delta (E^+ - E^-)$ and the posterior probability that $\Delta > 0$ under fenced (herbivores excluded) and unfenced (herbivores present) conditions. Horizontal dashed lines indicate $\Delta = 0$ for median differences and $\text{Pr}(\Delta > 0) = 0.5$. Endophyte effects were generally positive for *Agrostis hyemalis* and *Poa autumnalis*, but for *Elymus virginicus* benefits were context-dependent, with costs at lower precipitation




Fig S6: Predicted number of spikelets (upper panels) and endophyte effect ($\Delta = E^+ - E^-$, bottom panels) across precipitation gradients. Lines show posterior means for E^+ (blue) and E^- (red), with shading indicating 90% credible intervals. Points represent mean observed spikelets per plot, slightly jittered for visualization. Spikelets were measured for only two species (see Methods for details). The dashed line at $\Delta = 0$ indicates no endophyte effect. Panels are organized by species and herbivory treatment (Fenced, Unfenced).




Fig S7: Effects of precipitation and endophyte presence on spikelet (reproductive output) for three cool-season grass species. Median $\Delta (E^+ - E^-)$ and posterior probability $\Pr(\Delta > 0)$ are shown across precipitation quantiles. Herbivore exclusion treatments are indicated by color (Fenced = herbivores excluded, Unfenced = herbivores had access). Horizontal dashed lines indicate $\Delta = 0$ for medians and $\Pr(\Delta > 0) = 0.5$ for probabilities. Facets separate metrics and species.