

When Mutualism Becomes Costly: Abiotic Stress Shapes Grass Demography

Jacob K. Moutouama  *^{1,2,3}, Julia Martin¹, Alexandra Jimenez
Martín¹, Josh Fowler^{1,4}, Ali Campbell¹, Chris Oxley¹, Karl Schrader¹, and Tom E.X. Miller ¹

¹Program

in Ecology and Evolutionary Biology, Department of BioSciences, Rice University, Houston, TX USA

²Department of Botany, University of British Columbia, Vancouver, BC, Canada

³University of Miami, Department of Biology, Miami, FL USA

Running header: Stress and Grass–Endophyte Interactions

Keywords: demography, mutualism, parasitism, plant–fungi interactions, range limits, stress gradient hypothesis

Submitted to: *Ecology letters* (Letter)

Data accessibility statement: If the paper is accepted, all data and computer scripts supporting the results will be archived in a Zenodo package, with the DOI included at the end of the article. During peer review, our code (Stan and R) is available at <https://github.com/jmoutouama/endo-range-limits>.

Conflict of interest statement: None.

Authorship statement:

Abstract:

Main Text:

Figures: 6

Tables: X

References: X

*Corresponding author: jmoutouama@gmail.com

Abstract

Beneficial interactions between plants and fungi are ubiquitous and foundational to terrestrial ecosystems. Plant-fungal mutualisms are expected to strengthen under environmental stress, yet the relative importance of abiotic and biotic factors in eliciting mutualistic benefits remains unclear. We studied how the demographic effects of seed-transmitted fungal endophytes on native three grass host species varied in response to abiotic and biotic factors using geographically distributed field experiments along an aridity gradient, cross with herbivore access or exclusion. Vital rate models showed that all components of host fitness (survival, growth and fertility) declined with increasing, indicating reduced performance under abiotic stress. Declines were more pronounced in individuals hosting endophytes, suggesting that interactions that are beneficial under benign conditions can become costly under abiotic stress. On the other hand, herbivory had no detectable effect on host demography or on the demographic effects of hosting endophytes. Our results indicate that abiotic stress was the primary determinant of interaction outcomes between plants and fungal symbionts, but in the opposite direction as predicted by theory and prior work. These results highlight that symbiotic fungi can become liabilities under stress, with implications for the dynamics of plant-fungal mutualisms under global change.

Introduction

Plant-microbe symbioses are widespread and ecologically important. These interactions are famously context-dependent, where the direction and strength of the interaction outcome depend on the environment in which it occurs (Bronstein, 1994; Hoeksema and Bruna, 2015). For example, plant-microbe interactions that are beneficial under stressful conditions may become neutral or parasitic under benign conditions (Giauque et al., 2019).

Stress may derive from biotic and/or abiotic factors. Under biotic stress, such as herbivory, microbial symbiosis can benefit host plants by facilitating the production of secondary compounds that deter feeding or exert direct toxicity, thereby reducing herbivore growth, survival, and oviposition (Atala et al., 2022; Bastias et al., 2017; Vega, 2008). Similarly under abiotic stress, such as low precipitation or high temperature, **symbionts can increase their host tolerance (Clay and Schardl, 2002)¹. Mycorrhizal fungi form extensive networks that increase root surface area, helping plants absorb more water and nutrients such as phosphorus and nitrogen (Amijee et al., 1989).**² Symbiotic microbes can also stimulate plants to produce antioxidant enzymes like superoxide dismutase and catalase that reduce oxidative damage caused by stresses such as drought, salinity, and heat (Ruiz-lazona et al., 1996). All these mechanisms could help the host species better buffer against environmental stochasticity (Fowler et al., 2023).

In resource-limited environments, maintaining symbionts can also impose significant costs on hosts (Bronstein, 2001; Johnson et al., 1997).³ Because microbes depend on host-derived carbon and nutrients, plants must allocate part of their limited resources to sustaining the symbiont. This investment can reduce host growth, reproduction, or competitive ability, particularly when the benefits provided by the symbiont do not outweigh these costs (Klironomos, 2003).

Context-dependence⁴ raises the hypothesis that plant-microbe interactions are likely to vary across abiotic stress gradients, such as high temperatures or low precipitation, as well as under biotic stressors like herbivory, with significant implications for host fitness in variable environments. **If microbial symbiosis provides greater benefits under these stressful**

¹*Can you expand this a bit? The previous sentence gave a more thorough explanation of how symbionts can benefit hosts exposed to herbivory. How do symbionts promote tolerance of abiotic stress? Through this section, keep the scope pretty wide, so don't just focus on the grass-endophyte literature.*

²*Unclear how this relates to the preceding sentences. Is this meant to be an example of how symbionts promote tolerance of abiotic stress? If so, it does not come through clearly. What is the importance of access to phosphorous and nitrogen, and how does this related to abiotic stress?*

³*You are introducing a new idea, so I suggest starting a new paragraph with a strong topic sentence, and more fully developing the idea of costs of symbiosis.*

⁴*What is the topic of this paragraph, and what is it intended to do differently than the preceding paragraphs? A lot of this material overlaps heavily with what you have already presented above. Think about the logic structure that you want for the intro, and make sure your paragraphs are taking readers through a logical progression of ideas.*

conditions, symbionts could enhance host survival, growth, and reproduction.⁵ (Allsup et al., 2023; Rudgers et al., 2020). For example, fungal endophytes of grasses enhance host tolerance to abiotic stress in *Bromus laevipes* by improving population survival in dry environments (Afkhami et al., 2014; David et al., 2019). Even when symbionts do not directly increase survival under stress, they may still promote host fitness by enhancing growth and reproduction, thereby offsetting the negative effects of lower survival rates (Yule et al., 2013). Conversely, if microbial symbiosis becomes costly under stressful conditions, symbionts could reduce host fitness⁶ (Bennett and Groten, 2022; Benning and Moeller, 2021a,b). For instance, mycorrhizal fungi require substantial carbon subsidies from the host plant (up to 20% of photosynthetically fixed carbon) (Soudzilovskaia et al., 2015; Thirkell et al., 2020). When photosynthesis is limited by abiotic stress, this carbon cost may outweigh the benefits of symbiosis, leading to reduced host fitness.

Despite growing interest in the role of symbiosis in host performance, few studies have examined under natural conditions how both abiotic and biotic stressors shape the effects of symbionts on host demography along stress gradients (Louthan et al., 2015). Moreover, it remains unclear which stressor is most critical in eliciting the benefits of endophyte symbiosis⁷, or how multiple stressors may interact to influence host demography. Herbivory is a biotic factor that can significantly affect host fitness (Benning et al., 2019) and mediate the outcomes of symbiotic interactions. Herbivores reduce host fitness through tissue damage, defoliation, and altered resource allocation (Maron and Crone, 2006), and can also impose indirect ecological costs by influencing reproductive or growth strategies (Miller et al., 2008). These ecological costs may shape the selective landscape in ways that favor symbiont persistence or transmission (Clay et al., 2005). For example, herbivory can enhance the fitness benefits of endophyte infection by triggering host defense pathways or promoting vertical transmission of the symbiont (Agrawal et al., 1999; Gundel et al., 2020). If the ecological context of herbivory alters the net fitness effects of symbiosis, then interactions between herbivory and abiotic stress (e.g., drought or high temperature) may play a critical role in determining host population persistence.⁸

⁵Statements like this are a little circular ("if symbionts are beneficial then the plant would benefit.") Can you sharpen this?

⁶Also circular.

⁷What is the intended scope at this point in the paper? Is this meant to be a general statement about microbial symbioses, or are we now talking specifically about *Epichloe* symbioses in grasses? I think the Intro can provide clearer sign-posting for this. There is a lot of literature on drought and herbivory in the *Epichloe* literature. I think it would be good to have a paragraph late in the Intro that introduces this system and briefly covers the literature on drought and herbivory, highlighting that few (or no?) previous studies have looked at both.

⁸The opening sentence suggested that this paragraph would be about the combined and relative roles of abiotic and biotic stress, but most of this paragraph is actually about herbivory. Again, bring attention to the way you structure paragraphs. Use clear topic sentences, and make sure the content of the paragraph stays true to the topic sentence.

Working across an aridity gradient in the south-central US, we investigated how endophyte symbiosis influences **host-plant demography under varying abiotic stress**¹⁰. This region provides an ideal system to study biologically realistic stress gradients because it spans a strong east–west precipitation gradient, with mesic conditions in the east transitioning to increasingly arid conditions in the west. At each site along the gradient we also tested herbivory as a biotic stressor to evaluate its potential interactive effects with abiotic conditions. Specifically, we studied the symbiotic association between three native cool-season grass species (*Agrostis hyemalis*, *Elymus virginicus*, and *Poa autumnalis*) and their vertically transmitted fungal symbionts (*Epichloë* spp.). Our experiment was designed to address the **following questions**¹¹:

1. Under combined abiotic stress and herbivory, do endophytes provide fitness benefits or impose costs on their host plants?
2. Between abiotic and biotic factors, which plays a stronger role in **shaping host population demography in association with endophytes**¹²?

We hypothesized...¹³

Materials and methods

Study system

Our study focused on three cool-season perennial grasses native to woodland and prairie habitats of eastern North America: *Agrostis hyemalis*, *Elymus virginicus*, and *Poa autumnalis* (subfamily Pooideae) (Shaw, 2011). *A. hyemalis* is a small, short-lived perennial that germinates from autumn to late winter and typically flowers between March and July. *E. virginicus* is a larger, long-lived species that begins flowering as early as March or April and continues throughout the summer. *P. autumnalis* has a similar growth form to *E. virginicus* and flowers from late spring through summer. All three species host *Epichloë* fungal endophytes, which are primarily transmitted vertically through seeds. *Agrostis hyemalis* is associated with

⁹Here is where I might dedicate a paragraph to describing the biology and context-dependence of the grass-endophyte symbioses.

¹⁰This paper includes multiple vital rate responses, and it might be strategic to highlight in the Intro the value of looking at multiple vital rates, since they may respond differently to abiotic or biotic stress (I am thinking of the demographic compensation ideas).

¹¹I think these questions can be improved but I will keep working through the methods and results, then return to this.

¹²This is vague. Can you articulate this question more precisely?

¹³Could be here or elsewhere, but I recommend including hypotheses or predictions toward the end of the Intro. Readers will need to know what to look for in the results. This occurred to me when I got to the methods section about modeling effects aridity, herbivory, and symbiosis. There is not enough provided here to anticipate what types of analyses you would need to do, and what your expectations would be. So that is the main point of this comment: set up expectations for readers.

Epichloë amarillans (White Jr, 1994), *E. virginicus* typically hosts *Epichloë elymi* (Liu, 2023), and *P. autumnalis* can also harbor endophyte taxa capable of horizontal transmission via sexual stromata on inflorescences or asexual conidia on leaf surfaces (Clay and Schardl, 2002; Gundel et al., 2020; Leuchtmann et al., 2014). Horizontal transmission is rare, influenced by environmental and genetic factors, and was not observed in our study system. To examine the effects of endophytes on host species across a stress gradient, we established eight plots (1.5 m × 1.5 m) at each of seven sites along an aridity gradient in Louisiana and Texas (Fig. 1A-D). Each plot contained 15 individuals of a single species, with half of the plots protected from herbivores using exclusion cages to assess the interactive effects of endophytes and herbivory (Fig. 1E). In summer–fall 2021, seeds were collected from naturally symbiotic source populations and either heat-treated to remove endophytes or left untreated, producing symbiont-free (E^-) and symbiotic (E^+) plants from the same genetic lineages.

Source populations and symbiont removal

Plants used in the common garden experiment were derived from natural (source) populations throughout the natural distribution of the species (Fig. 1, Table S-1). Seeds were collected from these source populations, and some were heat-treated at 60°C for approximately five days (120 hours) to produce endophyte-negative plants (E^-), a method that eliminates endophytes without affecting seed viability. All seeds, both heat-treated and non-heat-treated, were planted in the Rice University greenhouse, where seedlings were fertilized every two weeks. Seedlings were then vegetatively propagated to produce enough individuals for the experiment. Before field planting, we confirmed endophyte status (E^+ or E^-) of all seedlings using either microscopy or an immunoblot assay. Leaf tissues were stained with aniline blue lactic acid and viewed under a compound microscope at 200x–400x to identify fungal hyphae. The immunoblot assay (Phytoscreen field tiller endophyte detection kit, Agrinostics Ltd. Co.) uses monoclonal antibodies that target proteins of *Epichloë* spp. and a chromagen to visually indicate presence or absence. Both methods yield similar detection rates.

Common garden experimental design and climate data

We conducted our common experiment across spanning the precipitation gradient from Louisiana to central Texas, USA (Fig. 1, Table S-2). To characterize the abiotic stress gradient, we collected daily temperature and precipitation data from PRISM (Hart and Bell, 2015). These daily values were used to calculate the mean temperature (°C) and cumulative precipitation (mm) for each year, spanning from the time plants were transplanted to when demographic data were collected. During our study period (2023–2025) realized precipitation at these

129 sites corresponded well to 30-year normals, with wetter eastern sites and drier western sites.
130 Mean annual temperature did not differ significantly across sites (Fig. S-1), so we focus on
131 precipitation as the primary abiotic factor that varied across sites. This gradient also overlaps
132 and exceeds the natural distributions of our focal species, ensuring realistic exposure to
133 climatic and biotic conditions (Fig. 1).

134 Plots were randomly assigned one of four initial endophyte frequency treatments (80%,
135 60%, 40%, or 20%) and one of two herbivore access treatments (fenced to exclude herbivores
136 or unfenced to allow access). In the fenced plots, insecticides were applied periodically to
137 eliminate small herbivorous insects, as *Agrostis hyemalis*, *Epichloë amarillans*, and *Poa autumnalis*
138 are naturally subject to herbivory by both insects and larger mammals. To ensure that our
139 herbivory treatment reflected the level of herbivory experienced by individual plants, we
140 counted the number of damaged plants per plot and estimated the proportion of damaged
141 plants per site and herbivore treatment. Our results confirmed that the herbivory treatment
142 was effective (Fig. 1F).




Figure 1: Distribution of common garden sites along a natural aridity gradient from Texas to Louisiana, US for (A) *Agrostis hyemalis*, (B) *Elymus virginicus*, and (C) *Poa autumnalis*. Climate data in A–C are based on 30-year normals of climate predictions (1993–2023) from PRISM data (Hart and Bell, 2015), which illustrate the precipitation gradient across sites (scale on the right is in mm). Red dots indicate the locations of source populations, while grey dots represent Global Biodiversity Information Facility (GBIF) occurrence records for each species within the study area (GBIF, 2025a,b,c). D) Cumulative precipitation during the study period (May 2023 – June 2025) at each site, ordered west to east. Values are also derived from PRISM climate data (Hart and Bell, 2015).

Demographic data collection

We collected demographic data, including survival, growth, and reproduction, during the flowering season of each species: *Agrostis hyemalis* and *Poa autumnalis* were censused in May, and *Elymus virginicus* was censused in June of 2023, 2024, and 2025. For each individual, plant survival was recorded as alive or dead and, for surviving plants, size was measured as the number of tillers. Growth was quantified as the log ratio of the number of tillers between consecutive censuses (i.e., $\log(\text{tiller}_{t+1}/\text{tiller}_t)$) for plants that were alive in both years. We recorded the number of inflorescences per plant for all three species and counted the number of spikelets on up to three inflorescences from reproducing individuals of *Poa autumnalis* and *Elymus virginicus*.

Assessing whether endophytes confer benefits or impose costs under abiotic and biotic stress

To assess whether endophytes confer benefits or impose costs under abiotic and biotic stress, we first developed two candidate models for each vital rate: one quadratic and one monotonic. All models shared the same hierarchical structure and were applied to survival, growth, and fertility. Survival was modeled with a Bernoulli distribution, growth with a Gaussian distribution, the number of inflorescences with a zero-inflated negative binomial distribution, and the number of spikelets with a negative binomial distribution. We modeled all vital rates (survival, growth, and fertility) using species-specific fixed effects for symbiosis (endophyte presence), precipitation, and herbivore access, including all relevant two- and three-way interactions. Each observation i corresponds to a plant in a given year; individuals can appear in multiple years. For simplicity, we omit the full subscript notation for species, plot, population, and site-year, which is explicitly included in the Stan model. We present the quadratic model here (Eq. 1) because it encompasses the linear model as a special case.

$$\begin{aligned} \mu_i = & \beta_0[Sp_i] + \beta_P[Sp_i]P_i + \beta_{P^2}[Sp_i]P_i^2 + \beta_{endo}[Sp_i]endo_i + \beta_{herb}[Sp_i]herb_i \\ & + \beta_{endo \times P}[Sp_i]endo_iP_i + \beta_{endo \times P^2}[Sp_i]endo_iP_i^2 + \beta_{herb \times P}[Sp_i]herb_iP_i \\ & + \beta_{herb \times endo}[Sp_i]herb_iendo_i + \beta_{endo \times herb \times P}[Sp_i]endo_iherb_iP_i \\ & + \phi_{\text{site}[i], Sp_i} + \omega_{\text{source}[i], Sp_i} + \rho_{\text{plot}[i], Sp_i}. \end{aligned} \quad (1)$$

Here, P_i is total precipitation (mm) averaged over the 12 months preceding the census, $endo_i$ indicates endophyte presence, and $herb_i$ indicates herbivore access. Random effects account for background variation at multiple hierarchical levels: $\phi_{\text{site}[i], Sp_i}$ captures site-to-site heterogeneity not explained by precipitation, $\omega_{\text{source}[i], Sp_i}$ captures variation associated with the provenance of transplants, and $\rho_{\text{plot}[i], Sp_i}$ captures plot-to-plot variation. Each species has

its own random effect for each site, source, and plot, while the variance parameters (σ_{site} , σ_{source} , σ_{plot}) are shared across species. This hierarchical structure allows the model to “borrow strength” across species, improving estimates of fixed effects and background variation, while still capturing species-specific responses to environmental heterogeneity. We confirmed that models effectively captured key aspects of the grass demography data, including means, standard deviations, and quantiles using a posterior predictive check (Fig. S-2).

Secondly, to determine the best-supported model for each vital rate, we compared candidate models (quadratic and monotonic) using leave-one-out cross-validation (LOOCV) (Vehtari et al., 2017). LOOCV combines validation and training by iteratively holding out one observation as a test set while fitting the model on the remaining $n - 1$ observations. This process is repeated n times, once for each observation, resulting in n fitted models (Silva and Zanella, 2024). The overall test error estimate is then obtained by averaging the prediction errors across all n models (Eq. 2) for each vital rate.

$$CV_n = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2 \quad (2)$$

Although we evaluated quadratic models, LOOCV showed that they provided minimal improvement over monotonic models (Table S-3). Therefore, all subsequent analyses used the monotonic model, which captures the observed patterns while maintaining model simplicity.

Finally, for each vital rate, precipitation was summarized at key quantiles (10th, 25th, 50th, 75th, and 90th percentiles) to represent the range of site-level abiotic stress. For each species (*Agrostis hyemalis*, *Elymus virginicus*, *Poa autumnalis*) and herbivory treatment (Fenced or Unfenced), we computed posterior predictions of survival probability for individuals with endophytes (E^+) and without endophytes (E^-) using the best Bayesian model outputs (Eq. 1). The difference ($\Delta = E^+ - E^-$) was calculated for each posterior sample to estimate median effects and 90% credible intervals. We also computed the posterior probability that $\Delta > 0$, representing the likelihood that endophytes confer benefits on vital rates under the given climate and herbivory conditions.

Drivers of endophyte effects on cool-season grass demography

To determine the primary drivers of endophyte-mediated effects on grass performance, we focused on interactions with climate and herbivory to identify when endophytes confer benefits or costs under multiple stressors (abiotic and biotic). The endophyte $\times$ climate and endophyte $\times$ herbivory $\times$ climate interactions summarize how the benefits of endophyte presence on survival, growth, or reproduction change across precipitation gradients and

herbivory treatments. The endophyte $\times$ herbivory interaction indicates whether endophytes mitigate or exacerbate herbivore impacts, and together these interactions highlight the environmental and biotic contexts that enhance or reduce symbiotic benefits.

Effect sizes were quantified using the posterior distributions from the hierarchical Bayesian models (Eq. 1). For survival, coefficients on the logit scale were exponentiated to obtain odds ratios (OR), representing the multiplicative change in the probability of survival associated with endophyte presence under different conditions. For flowering and spikelet production, coefficients were exponentiated to obtain incidence rate ratios (IRR), representing the proportional change in reproductive output. Growth was modeled with a Gaussian distribution, and effect sizes were expressed as the expected change in log-transformed growth per unit change in the predictor. For each species and vital rate, we summarized effects as mean OR, IRR, or growth change with 95% credible intervals. We also computed the posterior probability that OR or IRR exceeded 1, or that growth effects were positive, providing a direct measure of the likelihood that endophytes confer benefits under specific combinations of climate and herbivory. This approach allows comparison of the magnitude and certainty of abiotic, biotic, and interactive drivers on grass demography in biologically interpretable units.

All models were implemented in R (R Core Team, 2023) using Stan via the rstan interface (Stan Development Team, 2024).

Results

Endophyte effects on host fitness components across abiotic and biotic stress gradients

Across fitness components, *Epichloë* endophytes generally enhanced host performance under wetter conditions, while their effects were neutral or occasionally costly under drier conditions or in the presence of herbivores. Benefits were most apparent when herbivores were excluded.

For survival, endophytes effect was neutral or slightly costly at low precipitation, but became increasingly positive with wetter conditions, particularly in unfenced plots (Fig. 2; Fig. S-3). In *Agrostis hyemalis* and *Elymus virginicus*, median $\Delta(E^+ - E^-)$ ranged from -0.71 to 0.10 under dry conditions ($\Pr(\Delta > 0) = 0.37\text{--}0.61$; Table S-4) and increased to $0.95\text{--}1.54$ with high precipitation in unfenced plots ($\Pr(\Delta > 0) = 0.69\text{--}0.84$; Table S-4). In fenced plots, positive effects emerged only at the highest precipitation and were weaker (Fig. S-3). In contrast, *Poa autumnalis* showed little benefit when herbivores were presented (median $\Delta = -0.71$ to 1.54 , $\Pr(\Delta > 0) = 0.02\text{--}0.08$), but weakly positive survival when herbivores were excluded (median $\Delta = -0.13$ to 0.11 , $\Pr(\Delta > 0) = 0.57\text{--}0.71$).

Epichloë endophytes enhanced growth with increasing precipitation across species, though magnitude and certainty varied with herbivory. In *A. hyemalis*, effects were neutral at low-to-moderate precipitation in unfenced plots (median $\Delta = 0.006$ – 0.093) and became more positive at high precipitation (median $\Delta = 0.170$ – 0.187 ; $\text{Pr}(\Delta > 0) = 0.56$ – 0.74 ; Table S-5). Fenced plots showed similar but smaller effects. *E. virginicus* showed minor or negative effects under dry conditions (median $\Delta = -0.123$ to -0.008 ; $\text{Pr}(\Delta > 0) = 0.35$ – 0.49), turning beneficial at higher precipitation, especially in unfenced plots (median $\Delta = 0.207$ – 0.238 ; $\text{Pr}(\Delta > 0) = 0.81$ – 0.82). In *P. autumnalis*, effects were context-dependent: unfenced plots showed neutral to mild benefits at low precipitation (median $\Delta = 0.192$ – 0.258 ; $\text{Pr}(\Delta > 0) = 0.69$ – 0.85), becoming strongly positive at high precipitation, while fencing amplified benefits across the gradient (median $\Delta = 0.461$ – 1.179 ; $\text{Pr}(\Delta > 0) = 0.98$ – 1.00).

Epichloë endophytes also increased flowering and spikelet production, with effects generally stronger under wetter conditions (Figs. 4, 5; Figs. S-6). Flowering benefits were consistent across species, though decreased under dry conditions, with herbivore exclusion having little effect for *A. hyemalis* and *E. virginicus*, but enhancing benefits for *P. autumnalis* (median Δ up to 6.84 ; $\text{Pr}(\Delta > 0) \sim 1.00$; Table S-6). Spikelet production was similarly context-dependent: *E. virginicus* and *P. autumnalis* showed negative or weak effects under low precipitation, shifting to positive effects under high precipitation, with fencing amplifying benefits across the gradient (Table S-7).




Figure 2: Predicted survival rate (top panels) and endophyte effect ($\Delta = E^+ - E^-$, bottom panels) across precipitation gradients. Lines show posterior means for E^+ (blue) and E^- (red), with shading indicating 90% credible intervals. Points represent mean observed survival per plot, jittered for visualization. The dashed line at $\Delta = 0$ indicates no endophyte effect. Panels are organized by species and herbivory treatment (Fenced, Unfenced).




Figure 3: Predicted relative growth from year to the next year (top panels) and endophyte effect ($\Delta = E^+ - E^-$, bottom panels) across precipitation gradients. Lines show posterior means for E^+ (blue) and E^- (red), with shading indicating 90% credible intervals. Points represent mean observed growth per plot, jittered for visualization. The dashed line at $\Delta = 0$ indicates no endophyte effect. Panels are organized by species and herbivory treatment (Fenced, Unfenced).




Figure 4: Predicted number of inflorescences (upper panels) and the difference between endophyte-infected (E^+) and endophyte-free (E^-) plants ($\Delta(E^+ - E^-)$, lower panels) are shown across a gradient of precipitation. Shaded areas represent 90% credible intervals. Points indicate observed means, jittered for visualization. Panels are grouped by species and herbivory treatment (unfenced = with herbivory; fenced = herbivory excluded).




Figure 5: Predicted number of spikelets (upper panels) and endophyte effect ($\Delta = E^+ - E^-$, bottom panels) across precipitation gradients. Lines show posterior means for E^+ (blue) and E^- (red), with shading indicating 90% credible intervals. Points represent mean observed spikelets per plot, slightly jittered for visualization. Spikelets were measured for only two species (see Methods for details). The dashed line at $\Delta = 0$ indicates no endophyte effect. Panels are organized by species and herbivory treatment (Fenced, Unfenced).

Abiotic factors as the primary drivers of fitness components across the stress gradient

Endophytes were most mutualistic under wet conditions or when hosts were exposed to herbivores, but they reduced growth or survival under dry conditions with low herbivory

(Fig. 6). Survival increased under wetter conditions due to the Endophyte $\times$ Climate interaction (OR = 1.21–1.36; Pr(OR > 1) = 0.68–0.81; Table S-8). Herbivory modified these benefits in a species-specific manner: *Poa autumnalis* showed strongly enhanced survival when herbivores were present (OR = 1.94; Pr = 0.88; Table S-8), whereas responses in *Agrostis hyemalis* and *Elymus virginicus* were weak or uncertain. The three-way interaction further boosted survival in *A. hyemalis* and *E. virginicus* under herbivore exclusion in wet climates (OR = 1.15–1.45; Pr = 0.63–0.77; Table S-8), but had little effect on *P. autumnalis*. Growth responses were modest. Endophyte $\times$ Climate effects were near zero in *A. hyemalis* and *E. virginicus* (Pr(>0) $\approx$ 0.46–0.49; Table S-8), but moderately positive in *P. autumnalis* (Pr = 0.87). Herbivory tended to reduce growth in the first two species (Pr $\approx$ 0.10–0.17) while increasing it in *P. autumnalis* (Pr = 0.93). The three-way interaction was only strongly negative in *P. autumnalis* (mean = –0.55; Pr < 0.01; Table S-8). Reproduction was enhanced under combined endophyte and climate or herbivory effects. Flowering and spikelet output increased with Endophyte $\times$ Climate (IRR = 1.09–1.18; Pr(IRR > 1) > 0.66; Table S-8) and Endophyte $\times$ Herbivory in *P. autumnalis* (IRR = 1.31–1.96; Pr > 0.96; Table S-8), whereas three-way interactions were negligible.




Figure 6: Standardized coefficients from Bayesian models showing how endophytes interact with climate and herbivore access to influence demographic rates of host grasses. Panels correspond to (a) survival, (b) growth, (c) flowering (inflorescence production), and (d) spikelet production. Coefficients represent standardized effect sizes for interaction terms: Endophyte $\times$ Climate, Endophyte $\times$ Herbivory, and Endophyte $\times$ Herbivory $\times$ Climate for the three species shown in the plot: *Agrostis hyemalis* (AGHY), *Elymus virginicus* (ELVI), and *Poa autumnalis* (POAU).

Acknowledgements

This research was supported by National Science Foundation Division of Environmental Biology awards.

References

- Afkhami, M. E., McIntyre, P. J., and Strauss, S. Y. (2014). Mutualist-mediated effects on species' range limits across large geographic scales. *Ecology letters*, 17(10):1265–1273.
- Agrawal, A. A., Laforsch, C., and Tollrian, R. (1999). Transgenerational induction of defences in animals and plants. *Nature*, 401(6748):60–63.
- Allsup, C. M., George, I., and Lankau, R. A. (2023). Shifting microbial communities can enhance tree tolerance to changing climates. *Science*, 380(6647):835–840.
- Amijee, F., Tinker, P., and Stribley, D. (1989). The development of endomycorrhizal root systems: Vii. a detailed study of effects of soil phosphorus on colonization. *New Phytologist*, 111(3):435–446.
- Atala, C., Acuña-Rodríguez, I. S., Torres-Díaz, C., and Molina-Montenegro, M. A. (2022). Fungal endophytes improve the performance of host plants but do not eliminate the growth/defence trade-off. *New Phytologist*, 235(2).
- Bastias, D. A., Martínez-Ghersa, M. A., Ballaré, C. L., and Gundel, P. E. (2017). Epichloë fungal endophytes and plant defenses: not just alkaloids. *Trends in Plant Science*, 22(11):939–948.
- Bennett, A. E. and Groten, K. (2022). The costs and benefits of plant–arbuscular mycorrhizal fungal interactions. *Annual Review of Plant Biology*, 73(1):649–672.
- Benning, J. W., Eckhart, V. M., Geber, M. A., and Moeller, D. A. (2019). Biotic interactions contribute to the geographic range limit of an annual plant: herbivory and phenology mediate fitness beyond a range margin. *The American Naturalist*, 193(6):786–797.
- Benning, J. W. and Moeller, D. A. (2021a). Microbes, mutualism, and range margins: testing the fitness consequences of soil microbial communities across and beyond a native plant's range. *New Phytologist*, 229(5):2886–2900.
- Benning, J. W. and Moeller, D. A. (2021b). Plant–soil interactions limit lifetime fitness outside a native plant's geographic range margin. *Ecology*, 102(3):e03254.
- Bronstein, J. L. (1994). Conditional outcomes in mutualistic interactions. *Trends in ecology & evolution*, 9(6):214–217.
- Bronstein, J. L. (2001). The exploitation of mutualisms. *Ecology letters*, 4(3):277–287.

- Clay, K., Holah, J., and Rudgers, J. A. (2005). Herbivores cause a rapid increase in hereditary symbiosis and alter plant community composition. *Proceedings of the National Academy of Sciences*, 102(35):12465–12470.
- Clay, K. and Schardl, C. (2002). Evolutionary origins and ecological consequences of endophyte symbiosis with grasses. *the american naturalist*, 160(S4):S99–S127.
- David, A. S., Quintana-Ascencio, P. F., Menges, E. S., Thapa-Magar, K. B., Afkhami, M. E., and Searcy, C. A. (2019). Soil microbiomes underlie population persistence of an endangered plant species. *The American Naturalist*, 194(4):488–494.
- Fowler, J. C., Donald, M. L., Bronstein, J. L., and Miller, T. E. (2023). The geographic footprint of mutualism: How mutualists influence species’ range limits. *Ecological Monographs*, 93(1):e1558.
- GBIF (2025a). Occurrence download for *Agrostis hyemalis*. Accessed 5 November 2025.
- GBIF (2025b). Occurrence download for *Elymus virginicus*. Accessed 5 November 2025.
- GBIF (2025c). Occurrence download for *Poa autumnalis*. Accessed 5 November 2025.
- Giauque, H., Connor, E. W., and Hawkes, C. V. (2019). Endophyte traits relevant to stress tolerance, resource use and habitat of origin predict effects on host plants. *New Phytologist*, 221(4):2239–2249.
- Gundel, P. E., Sun, P., Charlton, N. D., Young, C. A., Miller, T. E., and Rudgers, J. A. (2020). Simulated folivory increases vertical transmission of fungal endophytes that deter herbivores and alter tolerance to herbivory in *poa autumnalis*. *Annals of Botany*, 125(6):981–991.
- Hart, E. M. and Bell, K. (2015). *prism: Download data from the Oregon prism project*. R package version 0.0.6.
- Hoeksema, J. D. and Bruna, E. M. (2015). Context-dependent outcomes of mutualistic interactions. *Mutualism*, 10:181–202.
- Johnson, N. C., Graham, J. H., and Smith, F. A. (1997). Functioning of mycorrhizal associations along the mutualism–parasitism continuum. *The New Phytologist*, 135(4):575–585.
- Klironomos, J. N. (2003). Variation in plant response to native and exotic arbuscular mycorrhizal fungi. *Ecology*, 84(9):2292–2301.

- Leuchtman, A., Bacon, C. W., Schardl, C. L., White Jr, J. F., and Tadych, M. (2014). Nomenclatural realignment of neotyphodium species with genus epichloë. *Mycologia*, 106(2):202–215.
- Liu, M. (2023). Fungi canadenses no. 351: Epichloe elymi. *Canadian Journal of Plant Pathology*, 45(3):300–303.
- Louthan, A. M., Doak, D. F., and Angert, A. L. (2015). Where and when do species interactions set range limits? *Trends in ecology & evolution*, 30(12):780–792.
- Maron, J. L. and Crone, E. (2006). Herbivory: effects on plant abundance, distribution and population growth. *Proceedings of the Royal Society B: Biological Sciences*, 273(1601):2575–2584.
- Miller, T. E. X., Tenhumberg, B., and Louda, S. M. (2008). Herbivore-mediated ecological costs of reproduction shape the life history of an iteroparous plant. *The American Naturalist*, 171(2):141–149.
- R Core Team (2023). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rudgers, J. A., Afkhami, M. E., Bell-Dereske, L., Chung, Y. A., Crawford, K. M., Kivlin, S. N., Mann, M. A., and Nuñez, M. A. (2020). Climate disruption of plant-microbe interactions. *Annual review of ecology, evolution, and systematics*, 51:561–586.
- Ruiz-lazona, J. M., Azcón, R., and Palma, J. (1996). Superoxide dismutase activity in arbuscular mycorrhizal lactuca sativa plants subjected to drought stress. *New Phytologist*, 134(2):327–333.
- Shaw, R. B. (2011). *Guide to Texas grasses*. Texas A&M University Press.
- Silva, L. A. and Zanella, G. (2024). Robust leave-one-out cross-validation for high-dimensional bayesian models. *Journal of the American Statistical Association*, 119(547):2369–2381.
- Soudzilovskaia, N. A., van der Heijden, M. G., Cornelissen, J. H., Makarov, M. I., Onipchenko, V. G., Maslov, M. N., Akhmetzhanova, A. A., and van Bodegom, P. M. (2015). Quantitative assessment of the differential impacts of arbuscular and ectomycorrhiza on soil carbon cycling. *New Phytologist*, 208(1):280–293.
- Stan Development Team (2024). RStan: the R interface to Stan. R package version 2.32.6.
- Thirkell, T. J., Pastok, D., and Field, K. J. (2020). Carbon for nutrient exchange between arbuscular mycorrhizal fungi and wheat varies according to cultivar and changes in atmospheric carbon dioxide concentration. *Global change biology*, 26(3):1725–1738.

- 363 Vega, F. E. (2008). Insect pathology and fungal endophytes. *Journal of invertebrate pathology*,
364 98(3):277–279.
- 365 Vehtari, A., Gelman, A., and Gabry, J. (2017). Practical bayesian model evaluation using
366 leave-one-out cross-validation and waic. *Statistics and computing*, 27:1413–1432.
- 367 White Jr, J. F. (1994). Endophyte-host associations in grasses. xx. structural and reproductive
368 studies of *epichloë amarillans* sp. nov. and comparisons to *e. typhina*. *Mycologia*,
369 86(4):571–580.
- 370 Yule, K. M., Miller, T. E., and Rudgers, J. A. (2013). Costs, benefits, and loss of vertically
371 transmitted symbionts affect host population dynamics. *Oikos*, 122(10):1512–1520.

Supporting Information

S.0.1 Candidate models for abiotic and biotic drivers of cool-grass species

1. **Abiotic model:** includes precipitation (P) and its quadratic term (P^2) as predictors, along with endophyte status (endo) and its interactions with climate. Species-specific effects borrow strength across species:

$$\mu_i = \beta_0[\text{Sp}_i] + \beta_{\text{endo}}[\text{Sp}_i]\text{endo}_i + \beta_P[\text{Sp}_i]P_i + \beta_{P^2}[\text{Sp}_i]P_i^2 + \beta_{\text{endo} \times P}[\text{Sp}_i]\text{endo}_iP_i + \beta_{\text{endo} \times P^2}[\text{Sp}_i]\text{endo}_iP_i^2 + \phi_{\text{site}[i],\text{Sp}_i} + \omega_{\text{source}[i],\text{Sp}_i} + \rho_{\text{plot}[i],\text{Sp}_i}$$

2. **Biotic model:** includes endophyte status (endo), herbivory (herb), and their interaction, with species-specific effects:

$$\mu_i = \beta_0[\text{Sp}_i] + \beta_{\text{endo}}[\text{Sp}_i]\text{endo}_i + \beta_{\text{herb}}[\text{Sp}_i]\text{herb}_i + \beta_{\text{endo} \times \text{herb}}[\text{Sp}_i]\text{endo}_i\text{herb}_i + \phi_{\text{site}[i],\text{Sp}_i} + \omega_{\text{source}[i],\text{Sp}_i} + \rho_{\text{plot}[i],\text{Sp}_i}$$

3. **Full model:** combines abiotic and biotic predictors, including all relevant interactions up to the three-way interaction between endophyte, herbivory, and precipitation, with species-varying coefficients:

$$\mu_i = \beta_0[\text{Sp}_i] + \beta_P[\text{Sp}_i]P_i + \beta_{P^2}[\text{Sp}_i]P_i^2 + \beta_{\text{endo}}[\text{Sp}_i]\text{endo}_i + \beta_{\text{herb}}[\text{Sp}_i]\text{herb}_i + \beta_{\text{endo} \times P}[\text{Sp}_i]\text{endo}_iP_i + \beta_{\text{endo} \times P^2}[\text{Sp}_i]\text{endo}_iP_i^2 + \beta_{\text{herb} \times P}[\text{Sp}_i]\text{herb}_iP_i + \beta_{\text{herb} \times \text{endo}}[\text{Sp}_i]\text{herb}_i\text{endo}_i + \beta_{\text{endo} \times \text{herb} \times P}[\text{Sp}_i]\text{endo}_i\text{herb}_iP_i + \phi_{\text{site}[i],\text{Sp}_i} + \omega_{\text{source}[i],\text{Sp}_i} + \rho_{\text{plot}[i],\text{Sp}_i}$$

In these models, the β coefficients represent species-specific effects. The intercept $\beta_0[\text{Sp}_i]$ corresponds to the baseline value of the vital rate for species Sp_i in the absence of predictors. $\beta_{\text{endo}}[\text{Sp}_i]$ and $\beta_{\text{herb}}[\text{Sp}_i]$ describe the effects of endophyte presence and herbivory, respectively, for species Sp_i . The linear and quadratic effects of precipitation are captured by $\beta_P[\text{Sp}_i]$ and $\beta_{P^2}[\text{Sp}_i]$, while interactions between endophyte and precipitation are represented by $\beta_{\text{endo} \times P}[\text{Sp}_i]$ and $\beta_{\text{endo} \times P^2}[\text{Sp}_i]$. The interaction between endophyte and herbivory is given by $\beta_{\text{endo} \times \text{herb}}[\text{Sp}_i]$, the interaction between herbivory and precipitation by $\beta_{\text{herb} \times P}[\text{Sp}_i]$, and the three-way interaction among endophyte, herbivory, and precipitation by $\beta_{\text{endo} \times \text{herb} \times P}[\text{Sp}_i]$. All β coefficients are modeled hierarchically across species, allowing information to be shared while capturing species-specific responses. All models included a hierarchical random effects structure accounting for site-year, source population, and plot variation, with species-specific random effects $\phi_{\text{site}[i],\text{Sp}_i} \sim \mathcal{N}(0, \sigma_{\text{site}})$, $\omega_{\text{source}[i],\text{Sp}_i} \sim \mathcal{N}(0, \sigma_{\text{source}})$, and $\rho_{\text{plot}[i],\text{Sp}_i} \sim \mathcal{N}(0, \sigma_{\text{plot}})$.

S.1 Supporting Tables

Table S-1: Geographic coordinates and brief descriptions of source populations for *Agrostis hyemalis* (AGHY), *Elymus virginicus* (ELVI), and *Poa autumnalis* (POAU).

Species	Population	Latitude	Longitude	Description
AGHY	SJC	30.771969	-95.200620	Ecolab property
AGHY	COHR	29.672833	-96.567517	Columbus, TX herbarium record
AGHY	LPEF	30.085383	-97.172508	Lost Pines
POAU	LA7	30.849660	-93.276772	DeRidder, LA
POAU	LA1	32.568125	-93.932976	Jean Lafitte National Historical Park and Preserve
POAU	LA2	31.183040	-92.616969	Jean Lafitte National Historical Park and Preserve
POAU	SFAEF	31.492158	-94.772200	Stephen F. Austin Experimental Forest, TX
POAU	WB	30.399346	-95.150026	Sam Houston National Forest
ELVI	HUNT	30.741797	-95.474508	Sam Houston State University Field Station
ELVI	SHS	30.741797	-95.474508	Sam Houston State University Field Station
ELVI	RL5	30.471717	-99.783803	Texas Tech University, Junction
ELVI	PALM	29.591623	-97.584781	Palmetto State Park, TX
ELVI	JLP	29.785105	-90.114933	Jean Lafitte Park, outside of New Orleans

Table S-2: Common garden and field experiment sites used for reciprocal transplant and environmental manipulation studies.

Site	Code	Latitude	Longitude
University of Louisiana Ecology Center	LAF	30.304773	-92.012401
Center for Biological Field Studies	HUN	30.742815	-95.476881
TAMU Ecology and Natural Resource Teaching Area	COL	30.569302	-96.366324
UT Stengl Lost Pines Field Station	BAS	30.092343	-97.172442
1335 Medina Hwy, Kerrville, TX	KER	30.028437	-99.142888
Brackenridge Field Laboratory	BFL	30.284866	-97.780675
TAMU Sonora Station	SON	30.273467	-100.561463

Table S-3: Comparison of candidate models (quadratic vs. monotonic) for each vital rate based on leave-one-out cross-validation (LOOCV). ELPD difference is relative to the best model.

Vital rate	Model type	ELPD difference	SE difference
Survival	Quadratic	0	0.000
Survival	Monotonic	-0.653	0.997
Growth	Quadratic	0	0.000
Growth	Monotonic	-0.310	1.040
Flowering	Quadratic	0	0.000
Flowering	Monotonic	-0.562	1.010
Spikelet	Quadratic	0	0.078
Spikelet	Monotonic	-0.078	1.300

Table S-4: Posterior summaries of $\Delta (E^+ - E^-)$ for survival at precipitation quantiles for each species and herbivore exclusion treatment. Median, 90% credible interval, and posterior probability that $\Delta > 0$ are shown. Herbivore exclusion: Fenced = herbivores excluded, Unfenced = herbivores had access.

Species	Herbivore exclusion	Precipitation (mm)	Median Δ	Lower 90%	Upper 90%	P($\Delta > 0$)
<i>Agrostis hyemalis</i>	Unfenced	765.76	-0.0126	-0.1169	0.0719	0.3685
<i>Agrostis hyemalis</i>	Unfenced	848.28	-0.0099	-0.1011	0.0686	0.3965
<i>Agrostis hyemalis</i>	Unfenced	1113.25	0.0017	-0.0718	0.0759	0.5192
<i>Agrostis hyemalis</i>	Unfenced	1644.36	0.0258	-0.0675	0.1472	0.6943
<i>Agrostis hyemalis</i>	Unfenced	2163.28	0.0411	-0.0870	0.2249	0.7290
<i>Agrostis hyemalis</i>	Fenced	765.76	-0.0508	-0.1728	0.0156	0.0998
<i>Agrostis hyemalis</i>	Fenced	848.28	-0.0467	-0.1532	0.0189	0.1105
<i>Agrostis hyemalis</i>	Fenced	1113.25	-0.0278	-0.1131	0.0376	0.2291
<i>Agrostis hyemalis</i>	Fenced	1644.36	0.0134	-0.0899	0.1304	0.6016
<i>Agrostis hyemalis</i>	Fenced	2163.28	0.0439	-0.0888	0.2178	0.7282
<i>Elymus virginicus</i>	Unfenced	765.76	-0.0285	-0.1885	0.1340	0.3746
<i>Elymus virginicus</i>	Unfenced	848.28	-0.0174	-0.1651	0.1345	0.4170
<i>Elymus virginicus</i>	Unfenced	1113.25	0.0195	-0.1028	0.1465	0.6085
<i>Elymus virginicus</i>	Unfenced	1644.36	0.0765	-0.0642	0.2293	0.8173
<i>Elymus virginicus</i>	Unfenced	2163.28	0.1053	-0.0723	0.3098	0.8399
<i>Elymus virginicus</i>	Fenced	765.76	-0.1265	-0.2861	0.0322	0.0948
<i>Elymus virginicus</i>	Fenced	848.28	-0.0934	-0.2375	0.0457	0.1362
<i>Elymus virginicus</i>	Fenced	1113.25	-0.0019	-0.1130	0.1083	0.4883
<i>Elymus virginicus</i>	Fenced	1644.36	0.0883	-0.0246	0.2462	0.9023
<i>Elymus virginicus</i>	Fenced	2163.28	0.1131	-0.0047	0.3343	0.9435
<i>Poa autumnalis</i>	Unfenced	765.76	-0.0133	-0.1064	0.0219	0.1818
<i>Poa autumnalis</i>	Unfenced	848.28	-0.0226	-0.1354	0.0263	0.1708
<i>Poa autumnalis</i>	Unfenced	1113.25	0.0195	-0.1028	0.1465	0.1458
<i>Poa autumnalis</i>	Unfenced	1644.36	0.0769	-0.1765	0.0769	0.2885
<i>Poa autumnalis</i>	Unfenced	2163.28	0.0719	-0.1142	0.4382	0.4382
<i>Poa autumnalis</i>	Fenced	765.76	0.0060	-0.0338	0.0885	0.6755
<i>Poa autumnalis</i>	Fenced	848.28	0.0104	-0.0435	0.1079	0.6883
<i>Poa autumnalis</i>	Fenced	1113.25	0.0315	-0.0721	0.1574	0.7088
<i>Poa autumnalis</i>	Fenced	1644.36	0.0204	-0.0984	0.1579	0.6371
<i>Poa autumnalis</i>	Fenced	2163.28	0.0030	-0.0898	0.1139	0.5673

Table S-5: Posterior summaries of $\Delta (E^+ - E^-)$ for growth across precipitation quantiles for each species and herbivore treatment. Positive median values indicate beneficial effects of endophytes; negative medians indicate costly effects.

Species	Herbivore exclusion	Precipitation (mm)	Median Δ	Lower 90%	Upper 90%	P($\Delta > 0$)
<i>Agrostis hyemalis</i>	Unfenced	773.91	-0.0058	-0.5523	0.4940	0.5608
<i>Agrostis hyemalis</i>	Unfenced	1075.09	0.0555	-0.3212	0.5970	0.6812
<i>Agrostis hyemalis</i>	Unfenced	1316.25	0.0929	-0.2332	0.4248	0.7393
<i>Agrostis hyemalis</i>	Unfenced	1979.35	0.1692	-0.2486	0.5977	0.7744
<i>Agrostis hyemalis</i>	Unfenced	2163.28	0.1868	-0.2787	0.6595	0.7724
<i>Agrostis hyemalis</i>	Fenced	773.91	0.0129	-0.4028	0.5208	0.6535
<i>Agrostis hyemalis</i>	Fenced	1075.09	0.0727	-0.2255	0.3735	0.6589
<i>Agrostis hyemalis</i>	Fenced	1316.25	0.1082	-0.1698	0.3944	0.7393
<i>Agrostis hyemalis</i>	Fenced	1979.35	0.1809	-0.2083	0.5777	0.7744
<i>Agrostis hyemalis</i>	Fenced	2163.28	0.1975	-0.2318	0.6393	0.7724
<i>Elymus virginicus</i>	Unfenced	773.91	-0.1229	-0.6743	0.3539	0.3539
<i>Elymus virginicus</i>	Unfenced	1075.09	-0.0079	-0.4102	0.4873	0.4873
<i>Elymus virginicus</i>	Unfenced	1316.25	0.0626	-0.2815	0.6218	0.6218
<i>Elymus virginicus</i>	Unfenced	1979.35	0.2071	-0.1825	0.8061	0.8061
<i>Elymus virginicus</i>	Unfenced	2163.28	0.2379	-0.1845	0.8208	0.8208
<i>Elymus virginicus</i>	Fenced	773.91	-0.0684	-0.5652	0.4161	0.4161
<i>Elymus virginicus</i>	Fenced	1075.09	-0.0264	-0.3481	0.4470	0.4470
<i>Elymus virginicus</i>	Fenced	1316.25	-0.0004	-0.2469	0.4988	0.4988
<i>Elymus virginicus</i>	Fenced	1979.35	0.0524	-0.2399	0.6128	0.6128
<i>Elymus virginicus</i>	Fenced	2163.28	0.0625	-0.2673	0.6203	0.6203
<i>Poa autumnalis</i>	Unfenced	773.91	0.1916	-0.4565	0.6861	0.6861
<i>Poa autumnalis</i>	Unfenced	1075.09	0.2583	-0.1590	0.8455	0.8455
<i>Poa autumnalis</i>	Unfenced	1316.25	0.3025	0.0011	0.9508	0.9508
<i>Poa autumnalis</i>	Unfenced	1979.35	0.3888	0.1279	0.9929	0.9929
<i>Poa autumnalis</i>	Unfenced	2163.28	0.4076	0.1048	0.9875	0.9875
<i>Poa autumnalis</i>	Fenced	773.91	1.1794	0.5397	1.8184	0.9989
<i>Poa autumnalis</i>	Fenced	1075.09	0.9284	0.5167	1.3384	1.0000
<i>Poa autumnalis</i>	Fenced	1316.25	0.7718	0.4755	1.0747	1.0000
<i>Poa autumnalis</i>	Fenced	1979.35	0.4611	0.1732	0.7596	0.9958
<i>Poa autumnalis</i>	Fenced	2163.28	0.3940	0.0647	0.7341	0.9773

Table S-6: Posterior summaries of Δ ($E^+ - E^-$) for flowering output at precipitation quantiles for each species and herbivore exclusion treatment. Median, 90% credible interval, and posterior probability that $\Delta > 0$ are shown. Herbivore exclusion: Fenced = herbivores excluded, Unfenced = herbivores had access.

Species	Herbivore exclusion	Precipitation (mm)	Median Δ	Lower 90%	Upper 90%	P($\Delta > 0$)
<i>Agrostis hyemalis</i>	Unfenced	773.91	-0.0073	-0.75299	0.86870	0.4940
<i>Agrostis hyemalis</i>	Unfenced	1075.09	0.1156	-0.72258	1.03730	0.5970
<i>Agrostis hyemalis</i>	Unfenced	1316.25	0.2627	-0.72466	1.37065	0.6812
<i>Agrostis hyemalis</i>	Unfenced	1979.35	0.8796	-1.36442	4.4190	0.7445
<i>Agrostis hyemalis</i>	Unfenced	2163.28	1.0972	-1.75172	5.9933	0.7410
<i>Agrostis hyemalis</i>	Fenced	773.91	0.0223	-0.76233	0.93648	0.5208
<i>Agrostis hyemalis</i>	Fenced	1075.09	0.2099	-0.69651	1.25368	0.6589
<i>Agrostis hyemalis</i>	Fenced	1316.25	0.4266	-0.70743	1.80962	0.7393
<i>Agrostis hyemalis</i>	Fenced	1979.35	1.3034	-1.61430	5.89453	0.7744
<i>Agrostis hyemalis</i>	Fenced	2163.28	1.6062	-2.06481	7.75656	0.7724
<i>Elymus virginicus</i>	Unfenced	773.91	-0.0709	-0.44121	0.28281	0.3539
<i>Elymus virginicus</i>	Unfenced	1075.09	-0.0056	-0.34000	0.31918	0.4873
<i>Elymus virginicus</i>	Unfenced	1316.25	0.0542	-0.27674	0.39898	0.6218
<i>Elymus virginicus</i>	Unfenced	1979.35	0.2372	-0.22767	0.89597	0.8061
<i>Elymus virginicus</i>	Unfenced	2163.28	0.2921	-0.24417	1.10392	0.8208
<i>Elymus virginicus</i>	Fenced	773.91	-0.0964	-0.87499	0.79622	0.4161
<i>Elymus virginicus</i>	Fenced	1075.09	-0.0476	-0.66448	0.62745	0.4470
<i>Elymus virginicus</i>	Fenced	1316.25	-0.0006	-0.55277	0.58266	0.4988
<i>Elymus virginicus</i>	Fenced	1979.35	0.1347	-0.67562	1.11263	0.6128
<i>Elymus virginicus</i>	Fenced	2163.28	0.1640	-0.79414	1.37908	0.6203
<i>Poa autumnalis</i>	Unfenced	773.91	0.0029	-0.02154	0.07747	0.6861
<i>Poa autumnalis</i>	Unfenced	1075.09	0.0414	-0.02992	0.29571	0.8455
<i>Poa autumnalis</i>	Unfenced	1316.25	0.1812	0.00072	0.82475	0.9508
<i>Poa autumnalis</i>	Unfenced	1979.35	2.8401	0.31057	20.2205	0.9929
<i>Poa autumnalis</i>	Unfenced	2163.28	5.0522	0.37729	46.2966	0.9875
<i>Poa autumnalis</i>	Fenced	773.91	0.0744	0.00658	0.64876	0.9989
<i>Poa autumnalis</i>	Fenced	1075.09	0.3639	0.07496	1.53393	1.0000
<i>Poa autumnalis</i>	Fenced	1316.25	0.9483	0.23684	3.28046	1.0000
<i>Poa autumnalis</i>	Fenced	1979.35	5.2444	0.61659	36.1808	0.9958
<i>Poa autumnalis</i>	Fenced	2163.28	6.8435	0.35587	65.0251	0.9773

Table S-7: Posterior summaries of Δ ($E^+ - E^-$) at precipitation quantiles for each species and herbivore exclusion treatment. Median, 90% credible interval, and posterior probability that $\Delta > 0$ are shown. Herbivore exclusion: Fenced = herbivores excluded, Unfenced = herbivores had access.

Species	Herbivore exclusion	Precipitation (mm)	Median Δ	Lower 90%	Upper 90%	$P(\Delta > 0)$
<i>Elymus virginicus</i>	Unfenced	1035.41	-1.4080	-5.6588	2.5136	0.272
<i>Elymus virginicus</i>	Unfenced	1276.99	-0.6715	-4.3261	2.8829	0.371
<i>Elymus virginicus</i>	Unfenced	1644.36	0.3298	-3.1403	3.9781	0.564
<i>Elymus virginicus</i>	Unfenced	2163.28	1.5276	-2.5967	6.4739	0.737
<i>Elymus virginicus</i>	Unfenced	2182.71	1.5725	-2.5800	6.5641	0.741
<i>Elymus virginicus</i>	Fenced	1035.41	-1.0142	-4.4019	2.6090	0.306
<i>Elymus virginicus</i>	Fenced	1276.99	0.0749	-2.6481	2.9073	0.517
<i>Elymus virginicus</i>	Fenced	1644.36	1.5186	-1.0554	4.5225	0.839
<i>Elymus virginicus</i>	Fenced	2163.28	3.2830	-0.5052	8.7059	0.922
<i>Elymus virginicus</i>	Fenced	2182.71	3.3381	-0.4955	8.9005	0.923
<i>Poa autumnalis</i>	Unfenced	1035.41	-1.2269	-5.1615	2.0682	0.253
<i>Poa autumnalis</i>	Unfenced	1276.99	-0.7580	-3.6361	1.9503	0.314
<i>Poa autumnalis</i>	Unfenced	1644.36	0.1225	-2.2064	2.5035	0.538
<i>Poa autumnalis</i>	Unfenced	2163.28	1.4394	-2.0529	5.7896	0.764
<i>Poa autumnalis</i>	Unfenced	2182.71	1.4904	-2.0767	5.9711	0.766
<i>Poa autumnalis</i>	Fenced	1035.41	3.3952	-0.5175	8.5092	0.927
<i>Poa autumnalis</i>	Fenced	1276.99	3.5317	0.4555	7.5555	0.969
<i>Poa autumnalis</i>	Fenced	1644.36	3.5791	0.7899	7.3800	0.984
<i>Poa autumnalis</i>	Fenced	2163.28	3.4620	-0.8494	9.3572	0.910
<i>Poa autumnalis</i>	Fenced	2182.71	3.4513	-0.9420	9.4863	0.906

Table S-8: Posterior summaries of interaction effects (mean estimate, 95% credible intervals, and posterior probability) for survival, growth, flowering, and spikelet production across three grass species.

Fitness component	Species	Parameter	Estimate	Lower CI	Upper CI	Probability
Survival	<i>Agrostis hyemalis</i>	Endophyte × Climate	1.213	0.658	2.354	0.730
Survival	<i>Elymus virginicus</i>	Endophyte × Climate	1.364	0.666	2.802	0.806
Survival	<i>Poa autumnalis</i>	Endophyte × Climate	1.255	0.483	3.344	0.681
Survival	<i>Agrostis hyemalis</i>	Endophyte × Herbivory	0.804	0.364	1.720	0.287
Survival	<i>Elymus virginicus</i>	Endophyte × Herbivory	0.868	0.347	2.119	0.376
Survival	<i>Poa autumnalis</i>	Endophyte × Herbivory	1.941	0.642	6.026	0.879
Survival	<i>Agrostis hyemalis</i>	Endophyte × Herbivory × Climate	1.149	0.503	2.525	0.630
Survival	<i>Elymus virginicus</i>	Endophyte × Herbivory × Climate	1.450	0.560	3.880	0.774
Survival	<i>Poa autumnalis</i>	Endophyte × Herbivory × Climate	0.737	0.229	2.394	0.303
Growth	<i>Agrostis hyemalis</i>	Endophyte × Climate	-0.005	-0.340	0.329	0.488
Growth	<i>Elymus virginicus</i>	Endophyte × Climate	-0.013	-0.290	0.268	0.461
Growth	<i>Poa autumnalis</i>	Endophyte × Climate	0.208	-0.156	0.559	0.873
Growth	<i>Agrostis hyemalis</i>	Endophyte × Herbivory	-0.283	-0.734	0.158	0.107
Growth	<i>Elymus virginicus</i>	Endophyte × Herbivory	-0.208	-0.621	0.218	0.169
Growth	<i>Poa autumnalis</i>	Endophyte × Herbivory	0.365	-0.135	0.862	0.929
Growth	<i>Agrostis hyemalis</i>	Endophyte × Herbivory × Climate	0.084	-0.296	0.462	0.666
Growth	<i>Elymus virginicus</i>	Endophyte × Herbivory × Climate	0.060	-0.296	0.405	0.634
Growth	<i>Poa autumnalis</i>	Endophyte × Herbivory × Climate	-0.549	-0.966	-0.141	0.004
Flowering	<i>Agrostis hyemalis</i>	bendoclim	1.092	0.711	1.646	0.656
Flowering	<i>Elymus virginicus</i>	bendoclim	1.178	0.817	1.698	0.805
Flowering	<i>Poa autumnalis</i>	bendoclim	1.103	0.725	1.689	0.676
Flowering	<i>Agrostis hyemalis</i>	bendoherb	1.014	0.577	1.784	0.523
Flowering	<i>Elymus virginicus</i>	bendoherb	0.987	0.549	1.781	0.478
Flowering	<i>Poa autumnalis</i>	bendoherb	1.964	1.052	3.807	0.982
Flowering	<i>Agrostis hyemalis</i>	bendoherbclim	0.997	0.604	1.670	0.496
Flowering	<i>Elymus virginicus</i>	bendoherbclim	0.900	0.567	1.437	0.332
Flowering	<i>Poa autumnalis</i>	bendoherbclim	0.640	0.380	1.055	0.038
Spikelet	<i>Elymus virginicus</i>	bendoclim	1.100	0.930	1.310	0.866
Spikelet	<i>Poa autumnalis</i>	bendoclim	1.103	0.888	1.362	0.817
Spikelet	<i>Elymus virginicus</i>	bendoherb	1.029	0.760	1.395	0.573
Spikelet	<i>Poa autumnalis</i>	bendoherb	1.315	0.969	1.786	0.961
Spikelet	<i>Elymus virginicus</i>	bendoherbclim	1.023	0.806	1.293	0.573
Spikelet	<i>Poa autumnalis</i>	bendoherbclim	0.873	0.682	1.119	0.145

S.2 Supporting Figures




Figure S-1: Abiotic conditions across the seven study sites. (A) Mean annual precipitation showing the pronounced east–west aridity gradient from mesic Louisiana to arid Texas. (B) Mean annual temperature across sites, which does not differ much. Panels highlight that precipitation, rather than temperature, constitutes the primary abiotic gradient in this system.




Figure S-2: Posterior predictive checks for the full demographic model. Dark blue lines show observed data (y), and light blue lines show replicated datasets generated from posterior predictions (y_{rep}). The strong overlap across survival, growth, flowering, and spikelet production indicates adequate model fit for all vital rates.




Figure S-3: Predicted effects of endophyte presence (E^+) versus absence (E^-) on survival across precipitation gradients for three cool-season grasses. Solid lines show the median $\Delta (E^+ - E^-)$, or the posterior probability that $\Delta > 0$ under fenced and unfenced conditions. Horizontal dashed lines indicate no effect ($\Delta = 0$) or a 50% probability of benefit. Vertical dashed lines represent the precipitation values at the most westerly natural occurrence of each species, based on GBIF records.




Figure S-4: Effects of precipitation and endophyte presence on growth across three cool-season grass species. Lines show the predicted difference in growth between endophyte-present (E^+) and endophyte-absent (E^-) plants (Median Δ , top row) and the posterior probability that this difference is positive ($\Pr(\Delta > 0)$, bottom row) under fenced and unfenced treatments. Predictions were calculated at the 10th, 25th, 50th, 75th, and 90th percentiles of precipitation observed in the study. Horizontal dashed lines indicate $\Delta = 0$ (top row) and $\Pr(\Delta > 0) = 0.5$ (bottom row). Vertical dashed lines represent the precipitation values at the most westerly natural occurrence of each species, based on GBIF records.




Figure S-5: Effects of endophyte infection on flowering output across a precipitation gradient for three cool-season grass species. Lines show the median posterior difference $\Delta(E^+ - E^-)$ and the posterior probability that $\Delta > 0$ under fenced (herbivores excluded) and unfenced (herbivores present) conditions. Horizontal dashed lines indicate $\Delta = 0$ for median differences and $\Pr(\Delta > 0) = 0.5$. Endophyte effects were generally positive for *Agrostis hyemalis* and *Poa autumnalis*, but for *Elymus virginicus* benefits were context-dependent, with costs at lower precipitation




Figure S-6: Effects of precipitation and endophyte presence on spike (reproductive output) for three cool-season grass species. Median $\Delta (E^+ - E^-)$ and posterior probability $\Pr(\Delta > 0)$ are shown across precipitation quantiles. Herbivore exclusion treatments are indicated by color (Fenced = herbivores excluded, Unfenced = herbivores had access). Horizontal dashed lines indicate $\Delta = 0$ for medians and $\Pr(\Delta > 0) = 0.5$ for probabilities. Facets separate metrics and species.