



Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

Adjustment Methods for Measurement Error

Jose Pina-Sánchez

Albert Varela



Adjustments

Sensitivity Analysis

Simulations

SIMEX

Bayesian Adjustments

Adjustment Methods

- We should always aim to improve data collection processes to avoid measurement error
- When that is not possible, we can (and should) adjust its impact
 - This enhances the rigour of our research
 - And allows us to analyse data that would otherwise be too dubious



Adjustment Methods

- Sometimes we can adjust the impact of measurement error directly
 - We can do so in some simple settings, where we can anticipate its impact



Adjustment Methods

- Sometimes we can adjust the impact of measurement error directly
 - We can do so in some simple settings, where we can anticipate its impact
- Ex.1, the effect of self-reported anxiety on life satisfaction (both of them subject to classical errors)
 - the reliability ratio can be derived by repeating the interview for a subsample of participants,
 - which can then be used to adjust the expected bias (assuming a simple linear model),

$$\hat{\beta}^* = \hat{\beta} \left(\frac{\sigma_X^2}{\sigma_X^2 + \sigma_U^2} \right)$$



Adjustment Methods

- Sometimes we can adjust the impact of measurement error directly
 - We can do so in some simple settings, where we can anticipate its impact
- Ex.1, the effect of self-reported anxiety on life satisfaction (both of them subject to classical errors)
 - the reliability ratio can be derived by repeating the interview for a subsample of participants,
 - which can then be used to adjust the expected bias (assuming a simple linear model),

$$\hat{\beta}^* = \hat{\beta} \left(\frac{\sigma_X^2}{\sigma_X^2 + \sigma_U^2} \right)$$

- Ex.2, the effect of immigration on crime recorded by the police (systematic multiplicative errors)
 - the under-recording can be estimated using victimisation surveys,
 - and we can adjust the estimate of interest accordingly (assuming a linear model),

$$\hat{\beta}^* = \hat{\beta} / \bar{U}$$



Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

Adjustment Methods

- When we can't trace out the impact of measurement error algebraically we need to use adjustment methods



Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

Adjustment Methods

- When we can't trace out the impact of measurement error algebraically we need to use adjustment methods
- Most adjustment methods require additional forms of data
 - Multiple reflective indicators (latent variable models)
 - Instrumental variables (two stage processes)
 - A validation subsample (multiple imputation)
 - Repeated observations (regression calibration)



Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

Adjustment Methods

- When we can't trace out the impact of measurement error algebraically we need to use adjustment methods
- Most adjustment methods require additional forms of data
 - Multiple reflective indicators (latent variable models)
 - Instrumental variables (two stage processes)
 - A validation subsample (multiple imputation)
 - Repeated observations (regression calibration)
- **Question:** Could you use any of these methods for the measurement problems you have encountered in your research?
 - Validation and repeated observations are hard to find when you rely on secondary data



Adjustment Methods

- We will focus on methods that can be used without additional data
 - SIMEX (Cook & Stefanski, 1994)
 - Simulations (*RCME* Pina-Sánchez et al., 2022)
 - Bayesian adjustments (Gustaffson, 2003)
- All we need is an intuition of the form and prevalence of the measurement error



Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

Estimating Measurement Error

- We can estimate the form and prevalence of measurement error in a given variable using different sources
- **Question:** Any ideas?



Estimating Measurement Error

- We can estimate the form and prevalence of measurement error in a given variable using different sources
- **Question:** Any ideas?
 - More valid estimates from the same population
E.g. Comparing crime rates from police statistics to victimisation surveys we can ascertain systematic errors in the former



Estimating Measurement Error

- We can estimate the form and prevalence of measurement error in a given variable using different sources
- **Question:** Any ideas?
 - More valid estimates from the same population
E.g. Comparing crime rates from police statistics to victimisation surveys we can ascertain systematic errors in the former
 - Measurement error studies from the literature
E.g. A test-retest mental health assessment conducted in a different country (Biemer et al., 2004)



Estimating Measurement Error

- We can estimate the form and prevalence of measurement error in a given variable using different sources
- **Question:** Any ideas?
 - More valid estimates from the same population
E.g. Comparing crime rates from police statistics to victimisation surveys we can ascertain systematic errors in the former
 - Measurement error studies from the literature
E.g. A test-retest mental health assessment conducted in a different country (Biemer et al., 2004)
 - Replicate a coding exercise for a subsample of cases
E.g. Manually review a subsample of automatically classified offenders' ethnicity based on their name



Estimating Measurement Error

- We can estimate the form and prevalence of measurement error in a given variable using different sources
- **Question:** Any ideas?
 - More valid estimates from the same population
E.g. Comparing crime rates from police statistics to victimisation surveys we can ascertain systematic errors in the former
 - Measurement error studies from the literature
E.g. A test-retest mental health assessment conducted in a different country (Biemer et al., 2004)
 - Replicate a coding exercise for a subsample of cases
E.g. Manually review a subsample of automatically classified offenders' ethnicity based on their name
 - Interviews with survey interviewers, experts (e.g. practitioners), or individuals from the target population



Estimating Measurement Error

- We can estimate the form and prevalence of measurement error in a given variable using different sources
- **Question:** Any ideas?
 - More valid estimates from the same population
E.g. Comparing crime rates from police statistics to victimisation surveys we can ascertain systematic errors in the former
 - Measurement error studies from the literature
E.g. A test-retest mental health assessment conducted in a different country (Biemer et al., 2004)
 - Replicate a coding exercise for a subsample of cases
E.g. Manually review a subsample of automatically classified offenders' ethnicity based on their name
 - Interviews with survey interviewers, experts (e.g. practitioners), or individuals from the target population
 - Our own educated guess as subject experts



Sensitivity Analysis

- Such estimates should be taken as highly uncertain
 - ‘Gold standard’ measures are rarely perfect
 - Problems of transportability with studies using different samples/populations
 - Subjective nature of qualitative methods
 - Researcher bias



Sensitivity Analysis

- Such estimates should be taken as highly uncertain
 - ‘Gold standard’ measures are rarely perfect
 - Problems of transportability with studies using different samples/populations
 - Subjective nature of qualitative methods
 - Researcher bias
- We should undertake multiple adjustments
 - Using a range of plausible values, as opposed to assuming we know the form and prevalence of measurement error mechanism/s perfectly



Sensitivity Analysis

- Such estimates should be taken as highly uncertain
 - ‘Gold standard’ measures are rarely perfect
 - Problems of transportability with studies using different samples/populations
 - Subjective nature of qualitative methods
 - Researcher bias
- We should undertake multiple adjustments
 - Using a range of plausible values, as opposed to assuming we know the form and prevalence of measurement error mechanism/s perfectly
- We will not obtain a single ‘adjusted’ finding
 - Rather, we will seek to assess how ‘sensitive’ or robust our findings are under different scenarios
 - This is known as sensitivity analysis



Simulations

- The idea is to use our understanding of the measurement error process to recreate the original variable
- Then repeat the analysis using the ‘adjusted’ variable
 - Ideally for a range of measurement error scenarios
- Examples:
 - The reporting rate of burglaries has fluctuated between 40% to 60% in England and Wales (Pina-Sánchez et al., 2022)
 - Men report an average 14 lifetime opposite-sex partners, women report 7 (Mitchell et al., 2019)



Adjustments


Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

Simulations: Under-recorded Crime



- We formalise the above intuition into a measurement model
 - $X^* = X \cdot U$ with $U \sim N(0.5, \sigma_U)$
- We rearrange the measurement model and substitute to adjust the error-prone variable
 - $\hat{X} = X^* / 0.5$



Adjustments

Sensitivity
Analysis


Simulations

SIMEX

Bayesian
Adjustments

Simulations: Underrecorded Crime

Police recorded burglaries in London (2011)





Adjustments

Sensitivity
Analysis


Simulations

SIMEX


Bayesian
Adjustments

Simulations: Underrecorded Crime

Police recorded burglaries in London (2011)



Estimated burglaries in London (2011)





Simulations: Lifetime Partners

- A slightly more complex measurement error mechanism
- If we assume the true number of partners is in the middle (i.e. men overreport as much as women underreport)
 - We have the following measurement error model

$$\begin{cases} X^* = X \cdot U_1; & \text{if } Z=\text{man} \\ X^* = X \cdot U_2; & \text{if } Z=\text{woman} \end{cases}$$
 - And the adjusted variable

$$\begin{cases} \hat{X} = X^*/1.33; & \text{if } Z=\text{man} \\ \hat{X} = X^*/0.66; & \text{if } Z=\text{woman} \end{cases}$$
 - With the 33% worked out for men as: $14/(14 - (7/2)) = 1.33$
and similarly for women: $7/(7 + (7/2)) = 0.66$



Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

Simulations

- Simulations represent a direct and simple approach to adjusting measurement error
 - Making them an intuitive, parsimonious and transparent method



Simulations

- Simulations represent a direct and simple approach to adjusting measurement error
 - Making them an intuitive, parsimonious and transparent method
- They can be applied to any kind of analysis
 - Focus on adjusting the error-prone variable, which can then be used anywhere we want
 - Many other adjustment methods can only be used in specific outcome models, or estimation methods



Simulations

- Simulations represent a direct and simple approach to adjusting measurement error
 - Making them an intuitive, parsimonious and transparent method
- They can be applied to any kind of analysis
 - Focus on adjusting the error-prone variable, which can then be used anywhere we want
 - Many other adjustment methods can only be used in specific outcome models, or estimation methods
- They are also remarkably flexible in that they can mimic a wide range of forms of measurement error and misclassification
 - Gallop & Weschle, 2019
- One exception being random errors
 - Even if we know the magnitude of the error mechanism, we will not be able to estimate each true value



Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

SIMEX

- A simulation-based, but indirect, approach to adjusting for measurement error
 - Simulates increasing layers of measurement error, to trace out its impact
 - Then extrapolates to retrieve the true finding, when no measurement error is present



SIMEX

- A simulation-based, but indirect, approach to adjusting for measurement error
 - Simulates increasing layers of measurement error, to trace out its impact
 - Then extrapolates to retrieve the true finding, when no measurement error is present
- The SIMulation-EXtrapolation algorithm
 - Assuming $Y = \alpha + \beta X^* + \epsilon$, and $X^* = X + U$



SIMEX

- A simulation-based, but indirect, approach to adjusting for measurement error
 - Simulates increasing layers of measurement error, to trace out its impact
 - Then extrapolates to retrieve the true finding, when no measurement error is present
- The SIMulation-EXtrapolation algorithm
 - Assuming $Y = \alpha + \beta X^* + \epsilon$, and $X^* = X + U$
 - ① Generate new variables with increasing levels of measurement error, $X_k^*(\lambda_k) = X^* + \sqrt{(\lambda_k)}U$, with $\lambda_k = (0.5, 1, 1.5, 2)$



SIMEX

- A simulation-based, but indirect, approach to adjusting for measurement error
 - Simulates increasing layers of measurement error, to trace out its impact
 - Then extrapolates to retrieve the true finding, when no measurement error is present
- The SIMulation-EXtrapolation algorithm
 - Assuming $Y = \alpha + \beta X^* + \epsilon$, and $X^* = X + U$
 - ① Generate new variables with increasing levels of measurement error, $X_k^*(\lambda_k) = X^* + \sqrt{(\lambda_k)}U$, with $\lambda_k = (0.5, 1, 1.5, 2)$
 - ② Re-estimate the outcome model using X_k^* , which produces increasingly biased estimates, $\hat{\beta}_k^* = \beta_1 \frac{S_X^2}{S_X^2 + (1 + \lambda_k)S_U^2}$



SIMEX

- A simulation-based, but indirect, approach to adjusting for measurement error
 - Simulates increasing layers of measurement error, to trace out its impact
 - Then extrapolates to retrieve the true finding, when no measurement error is present
- The SIMulation-EXtrapolation algorithm
 - Assuming $Y = \alpha + \beta X^* + \epsilon$, and $X^* = X + U$
 - ① Generate new variables with increasing levels of measurement error, $X_k^*(\lambda_k) = X^* + \sqrt{(\lambda_k)}U$, with $\lambda_k = (0.5, 1, 1.5, 2)$
 - ② Re-estimate the outcome model using X_k^* , which produces increasingly biased estimates, $\hat{\beta}_k^* = \beta_1 \frac{S_X^2}{S_X^2 + (1 + \lambda_k)S_U^2}$
 - ③ Steps 1 and 2 are repeated to obtain $\overline{\hat{\beta}_k^*}$, reduce simulation error



SIMEX

- A simulation-based, but indirect, approach to adjusting for measurement error
 - Simulates increasing layers of measurement error, to trace out its impact
 - Then extrapolates to retrieve the true finding, when no measurement error is present
- The SIMulation-EXtrapolation algorithm
 - Assuming $Y = \alpha + \beta X^* + \epsilon$, and $X^* = X + U$
 - ① Generate new variables with increasing levels of measurement error, $X_k^*(\lambda_k) = X^* + \sqrt{(\lambda_k)}U$, with $\lambda_k = (0.5, 1, 1.5, 2)$
 - ② Re-estimate the outcome model using X_k^* , which produces increasingly biased estimates, $\hat{\beta}_k^* = \beta_1 \frac{S_X^2}{S_X^2 + (1 + \lambda_k)S_U^2}$
 - ③ Steps 1 and 2 are repeated to obtain $\overline{\hat{\beta}_k^*}$, reduce simulation error
 - ④ $\overline{\hat{\beta}_k^*}$ and λ_k can now be paired and their relationship estimated



SIMEX

- A simulation-based, but indirect, approach to adjusting for measurement error
 - Simulates increasing layers of measurement error, to trace out its impact
 - Then extrapolates to retrieve the true finding, when no measurement error is present
- The SIMulation-EXtrapolation algorithm
 - Assuming $Y = \alpha + \beta X^* + \epsilon$, and $X^* = X + U$
 - ① Generate new variables with increasing levels of measurement error, $X_k^*(\lambda_k) = X^* + \sqrt{(\lambda_k)}U$, with $\lambda_k = (0.5, 1, 1.5, 2)$
 - ② Re-estimate the outcome model using X_k^* , which produces increasingly biased estimates, $\hat{\beta}_k^* = \beta_1 \frac{S_X^2}{S_X^2 + (1 + \lambda_k)S_U^2}$
 - ③ Steps 1 and 2 are repeated to obtain $\overline{\hat{\beta}_k^*}$, reduce simulation error
 - ④ $\overline{\hat{\beta}_k^*}$ and λ_k can now be paired and their relationship estimated
 - ⑤ $\hat{\beta}_{SIMEX}$ can now be calculated by extrapolating to $\lambda_k = -1$



Adjustments


Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

SIMEX





Adjustments


Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

SIMEX





Adjustments


Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

SIMEX





Adjustments


Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

SIMEX





Adjustments


Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

SIMEX





Adjustments


Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

SIMEX





Adjustments


Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

SIMEX





Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

SIMEX

- The quality of the adjustment depends on:
 - The accuracy with which we define the measurement error mechanism
 - Choosing the right extrapolation function



SIMEX

- The quality of the adjustment depends on:
 - The accuracy with which we define the measurement error mechanism
 - Choosing the right extrapolation function
- A very flexible approach
 - Works for all kinds of outcome models
 - An R package ([simex](#)) with built-in commands to explore general cases (e.g. classical errors, misclassification)
 - New packages exploring other measurement error forms (e.g. multiplicative errors)
 - Not perfectly flexible though, we can only explore pre-established measurement error forms
 - And explore the impact of measurement error when the variable affected is the predictor of interest



Bayesian Adjustments

- The most flexible approach
 - Can be used in any outcome model to adjust for any form of measurement error
 - Overcomes the limitations of simulations-based approaches (e.g. simulating classical errors, or multiplicative errors affecting count-duration data)



Bayesian Adjustments

- The most flexible approach
 - Can be used in any outcome model to adjust for any form of measurement error
 - Overcomes the limitations of simulations-based approaches (e.g. simulating classical errors, or multiplicative errors affecting count-duration data)
- We specify both an outcome and a measurement model
 - The former reflects the substantive relationship that we want to estimate
 - The latter can reflect any form of measurement error that we can express algebraically



Bayesian Adjustments

- The most flexible approach
 - Can be used in any outcome model to adjust for any form of measurement error
 - Overcomes the limitations of simulations-based approaches (e.g. simulating classical errors, or multiplicative errors affecting count-duration data)
- We specify both an outcome and a measurement model
 - The former reflects the substantive relationship that we want to estimate
 - The latter can reflect any form of measurement error that we can express algebraically
- These two (or more) models are estimated simultaneously
 - Using Markov chain Monte Carlo (MCMC) methods
 - We obtain a ‘posterior distribution’ for each estimate included in our models
 - This reflects the probability distribution of an estimate given the models we are using, the data that we observe, and any prior knowledge we might want to include



Adjustments


Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

Bayesian Adjustments





Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

Bayesian Adjustments

- Can be expanded in many different ways
 - Informative priors, we can incorporate any subjective knowledge we possess about any of the parameters to be estimated
 - Can adjust for measurement error and missing data simultaneously



Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

Bayesian Adjustments

- Can be expanded in many different ways
 - Informative priors, we can incorporate any subjective knowledge we possess about any of the parameters to be estimated
 - Can adjust for measurement error and missing data simultaneously
- Probably the method with the steepest learning curve
 - To really exploit the full flexibility of Bayesian methods we need to use Bayesian software (e.g. Stan, JAGS)
 - And learn more about Bayesian inference



Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

NCRM NATIONAL CENTRE FOR
RESEARCH METHODS

comprehensive training in research methods

www.ncrm.ac.uk





Adjustments

Sensitivity
Analysis

Simulations

SIMEX

Bayesian
Adjustments

Evaluation Form

www.ncrm.ac.uk/surveys/hub

