

# Reactor Kinetics

Justin Pounders

April 25, 2016

## PKE

$$\frac{dP}{dt} = \frac{\rho - \bar{\beta}}{\Lambda} P + \sum_{i=1}^6 \lambda_i C_i$$
$$\frac{dC_i}{dt} = \frac{\bar{\beta}_i}{\Lambda} P - \lambda_i C_i$$

# Inhour Equation

Assume constant reactivity and a solution of the form

$$P(t) = P_0 e^{\omega t}$$

$$C_i(t) = C_{i,0} e^{\omega t}$$

# Inhour Equation

Assume constant reactivity and a solution of the form

$$P(t) = P_0 e^{\omega t}$$

$$C_i(t) = C_{i,0} e^{\omega t}$$

Plugging into the PKE leads to

$$\omega P_0 e^{\omega t} = \frac{\rho - \bar{\beta}}{\Lambda} P_0 e^{\omega t} + \sum_{i=1}^6 \lambda_i C_{i,0} e^{\omega t}$$

$$\omega C_{i,0} e^{\omega t} = \frac{\bar{\beta}_i}{\Lambda} P_0 e^{\omega t} - \lambda_i C_{i,0} e^{\omega t}$$

# Inhour Equation

Assume constant reactivity and a solution of the form

$$P(t) = P_0 e^{\omega t}$$

$$C_i(t) = C_{i,0} e^{\omega t}$$

Plugging into the PKE leads to

$$\omega P_0 e^{\omega t} = \frac{\rho - \bar{\beta}}{\Lambda} P_0 e^{\omega t} + \sum_{i=1}^6 \lambda_i C_{i,0} e^{\omega t}$$

$$\omega C_{i,0} e^{\omega t} = \frac{\bar{\beta}_i}{\Lambda} P_0 e^{\omega t} - \lambda_i C_{i,0} e^{\omega t}$$

Combining the two equations results in the *inhour* equation

$$\rho = \Lambda \omega + \sum_{i=1}^6 \frac{\bar{\beta}_i \omega}{\omega + \lambda_i}$$

# Roots of the Inhour Equation

There are 7 roots of the inhour equation (they are the eigenvalues of the PKE *system matrix*):

$$\rho = \Lambda\omega_j + \sum_{i=1}^6 \frac{\bar{\beta}_i\omega_j}{\omega_j + \lambda_i}, \quad j = 1, 2, \dots, 7$$

# Roots of the Inhour Equation

There are 7 roots of the inhour equation (they are the eigenvalues of the PKE *system matrix*):

$$\rho = \Lambda\omega_j + \sum_{i=1}^6 \frac{\bar{\beta}_i\omega_j}{\omega_j + \lambda_i}, \quad j = 1, 2, \dots, 7$$

This means we have a solution of the form

$$P(t) = \sum_{j=0}^6 a_j e^{\omega_j t}$$

Six of the roots will always be negative

$$\omega_j < -\lambda_j, \quad j = 1, 2, \dots, 6$$

The root  $\omega_0$  may be positive or negative.

$$P(t) = a_0 e^{\omega_0 t} + \sum_{j=1}^6 a_j e^{-|\omega_j|t}$$

Asymptotically ( $t \gg 0$ ) we have

$$P(t) \approx a_0 e^{\omega_0 t} \sim e^{t/T}$$

where the (asymptotic) period is defined as

$$T = \omega_0^{-1}$$



As  $\rho \rightarrow -\infty$

$$T \rightarrow \lambda_1^{-1} \approx 80 \text{ seconds (U-235)}$$

# Supercritical (I)

For  $0 \leq \rho \leq \bar{\beta}/2$ ,

$$\omega_0 + \lambda_i \approx \lambda_i$$

# Supercritical (I)

For  $0 \leq \rho \leq \bar{\beta}/2$ ,

$$\omega_0 + \lambda_i \approx \lambda_i$$

so

$$\begin{aligned}\rho &= \omega_0 \left[ \Lambda + \sum_{i=1}^6 \frac{\bar{\beta}_i}{\lambda_i} \right] \\ &\approx \omega_0 \times 0.1 \text{ seconds}\end{aligned}$$

# Supercritical (I)

For  $0 \leq \rho \leq \bar{\beta}/2$ ,

$$\omega_0 + \lambda_i \approx \lambda_i$$

so

$$\begin{aligned}\rho &= \omega_0 \left[ \Lambda + \sum_{i=1}^6 \frac{\bar{\beta}_i}{\lambda_i} \right] \\ &\approx \omega_0 \times 0.1 \text{ seconds}\end{aligned}$$

This means that

$$T = 0.1\rho^{-1} \text{ seconds} \approx \frac{1}{4} \frac{\bar{\beta}}{\rho} \text{ minutes}$$

# Supercritical (II)

For  $\rho > \bar{\beta}$ ,

$$\omega_0 + \lambda_i \approx \omega_0$$

# Supercritical (II)

For  $\rho > \bar{\beta}$ ,

$$\omega_0 + \lambda_i \approx \omega_0$$

so

$$\rho = \Lambda \omega_0 + \bar{\beta}$$

# Supercritical (II)

For  $\rho > \bar{\beta}$ ,

$$\omega_0 + \lambda_i \approx \omega_0$$

so

$$\rho = \Lambda \omega_0 + \bar{\beta}$$

This means that

$$T = \frac{\Lambda}{\rho - \bar{\beta}} \approx 10^{-2} \left( \frac{\rho}{\bar{\beta}} - 1 \right)^{-1}$$