

**Problem 1:** Write down the gradient descent update rule for the objective function

$$J(\theta) = \theta^4 - 6\theta^3 + 11\theta^2 - 7\theta + 4$$

from class. Suppose the learning rate is  $\alpha$ .

**Problem 2:** Consider the affine objective function

$$J(\theta) = m\theta + b,$$

for some parameters  $m \neq 0$  and  $b \in \mathbb{R}$ .

(a) Write down the gradient descent update rule. Suppose the learning rate is  $\alpha$ .

(b) Find a closed form expression for  $\theta_t$ .

(c) Using your answer to (b), discuss convergence of the gradient descent algorithm.

**Problem 3:** Consider the quadratic objective function

$$J(\theta) = \theta^2.$$

(a) Write down the gradient descent update rule. Suppose the learning rate is  $\alpha$ .

(b) Find a closed form expression for  $\theta_t$ .

(c) Using your answer to (b), discuss convergence of the gradient descent algorithm.

**Problem 4:** Consider again the affine objective function

$$J(\theta) = m\theta + b,$$

for some parameters  $m \neq 0$  and  $b \in \mathbb{R}$ .

(a) Write down the gradient descent update rule with learning rate  $\alpha$  and decay rate  $\beta$ .

(b) Find a closed form expression for  $\theta_t$ .

(c) Using your answer to (b), discuss convergence of the gradient descent algorithm.

**Problem 5:** Consider the function

$$J : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad J(\boldsymbol{\theta}) = 2\theta_1^2 + 4\theta_1\theta_2 + 3\theta_2^2 - 8\theta_1 - 10\theta_2 + 9.$$

(a) Compute the directional first derivative  $J'_{\mathbf{v}}(1, 1)$  where  $\mathbf{v}^\top = (1, 0)$ .

(b) Compute the directional second derivative  $J''_{\mathbf{v}}(1, 1)$  where  $\mathbf{v}^\top = (1, 0)$ .

**Problem 6:** Re-do the previous problem, but use the relationship between directional first and second derivatives and gradient vectors and Hessian matrices.

**Problem 7:** Consider again the function  $J$  from Problem 5. Here it is, for reference:

$$J : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad J(\boldsymbol{\theta}) = 2\theta_1^2 + 4\theta_1\theta_2 + 3\theta_2^2 - 8\theta_1 - 10\theta_2 + 9.$$

- (a) Find the direction of maximum rate of change of  $J$  at the point  $\boldsymbol{\theta}^\top = (0, 0)$ . What is the rate of change in this direction?

- (b) Find the direction of minimum rate of change of  $J$  at the point  $\boldsymbol{\theta}^\top = (0, 0)$ . What is the rate of change in this direction?

**Problem 8:** Yet again, consider the function  $J$  from Problem 5:

$$J : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad J(\boldsymbol{\theta}) = 2\theta_1^2 + 4\theta_1\theta_2 + 3\theta_2^2 - 8\theta_1 - 10\theta_2 + 9.$$

Find and classify all extremizers of  $J$ .

**Problem 9:** For the fifth time, consider the function  $J$  from Problem 5:

$$J : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad J(\boldsymbol{\theta}) = 2\theta_1^2 + 4\theta_1\theta_2 + 3\theta_2^2 - 8\theta_1 - 10\theta_2 + 9.$$

Find the directions of extreme curvature at the point  $\boldsymbol{\theta} = (0, 0)$ . What are the curvatures in these directions?

**Problem 10:** One last time, consider the function  $J$  from Problem 5:

$$J : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad J(\boldsymbol{\theta}) = 2\theta_1^2 + 4\theta_1\theta_2 + 3\theta_2^2 - 8\theta_1 - 10\theta_2 + 9.$$

Set  $\mathbf{H} = \nabla^2 J(0, 0)$ . Compute the spectrum  $\sigma(\mathbf{H})$ , the spectral radius  $\rho(\mathbf{H})$ , and the condition number  $\kappa(\mathbf{H})$ .

**Problem 11:** Consider the objective function

$$J : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad J(\boldsymbol{\theta}) = 2\theta_1^2 + 4\theta_2^2.$$

- (a) Write down the gradient descent update rule. Suppose the learning rate is  $\alpha$ , while the decay rate is  $\beta = 0$ .

- (b) Find a closed form expression for  $\boldsymbol{\theta}_t$ .

- (c) Using your answer to (b), discuss convergence of the gradient descent algorithm.

**Problem 12:** Consider the stochastic objective function

$$J : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad J(\boldsymbol{\theta}) = \frac{1}{m} \sum_{i=1}^m \frac{1}{2} |\mathbf{x}_i - \boldsymbol{\theta}|^2$$

from class, where  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_m \in \mathbb{R}^2$  is an observed dataset.

- (a) Compute the update rule in the batch gradient descent algorithm with learning rate  $\alpha$  and decay rate  $\beta$ .

- (b) Assuming  $\beta = 0$ , discuss convergence of the batch gradient descent algorithm.