

Animal manure application and soil organic carbon stocks: a meta-analysis

ÉMILIE MAILLARD*† and DENIS A. ANGERS*

*Soils and Crops Research and Development Centre, Agriculture and Agri-Food Canada, 2560 Hochelaga Boulevard, Québec, QC G1V 2J3, Canada, †Faculté des sciences de l'agriculture et de l'alimentation, Pavillon Paul Comtois, Université Laval, 2425 rue de l'agriculture, Québec, QC G1V 0A6, Canada

Abstract

The impact of animal manure application on soil organic carbon (SOC) stock changes is of interest for both agronomic and environmental purposes. There is a specific need to quantify SOC change for use in national greenhouse gas (GHG) emission inventories. We quantified the response of SOC stocks to manure application from a large world-wide pool of individual studies and determined the impact of explanatory factors such as climate, soil properties, land use and manure characteristics. Our study is based on a meta-analysis of 42 research articles totaling 49 sites and 130 observations in the world. A dominant effect of cumulative manure-C input on SOC response was observed as this factor explained at least 53% of the variability in SOC stock differences compared to mineral fertilized or unfertilized reference treatments. However, the effects of other determining factors were not evident from our data set. From the linear regression relating cumulative C inputs and SOC stock difference, a global manure-C retention coefficient of $12\% \pm 4$ (95% Confidence Interval, CI) could be estimated for an average study duration of 18 years. Following an approach comparable to the Intergovernmental Panel on Climate Change, we estimated a relative SOC change factor of 1.26 ± 0.14 (95% CI) which was also related to cumulative manure-C input. Our results offer some scope for the refinement of manure retention coefficients used in crop management guidelines and for the improvement of SOC change factors for national GHG inventories by taking into account manure-C input. Finally, this study emphasizes the need to further document the long-term impact of manure characteristics such as animal species, especially pig and poultry, and manure management systems, in particular liquid vs. solid storage.

Keywords: animal manure, manure carbon input, manure retention coefficient, meta-analysis, soil carbon change factor, soil organic carbon

Received 17 May 2013 and accepted 22 September 2013

Introduction

As animal manure contains organic matter, an increase in soil organic carbon (SOC) content is generally expected following its land application, as reported in many individual studies. For example, in the southeastern United States, Sainju *et al.* (2008a) observed an increase of soil surface (0–20 cm) SOC stock of about 3.2 Mg C ha^{-1} after 10 years of poultry litter application in comparison to mineral fertilizer plots. In Nepal, after 25 annual cattle manure applications, surface (0–30 cm) SOC stocks were higher by about $19.1 \text{ Mg C ha}^{-1}$ than control (unfertilized) plots (Gami *et al.*, 2009). In China, after 22 years of pig manure application, the surface soil layer (0–15 cm) accumulated 3.8 Mg C ha^{-1} more than mineral fertilizer alone (Huang *et al.*, 2010a). However, some studies report no significant or even negative change of SOC stocks

following manure application (Franzluebbers *et al.*, 2001; Angers *et al.*, 2010). Obviously, there is great variability in the magnitude of SOC stocks change after manure application.

Individual studies and a few review articles have attempted to relate the variability in the magnitude of SOC change to various explanatory factors such as climate (Triberti *et al.*, 2008), manure application rate (Franzluebbers *et al.*, 2001), manure management system (farmyard manure/slurry) (Grignani *et al.*, 2007), soil texture (Gami *et al.*, 2009), initial SOC concentration (Dersch & Bohm, 2001), land use, and time of application. Few global studies have attempted to quantify the effects of these explanatory factors from all the available literature. Ogle *et al.* (2005) and Franzluebbers & Doraiswamy (2007) focused solely on the impact of climatic zone on SOC change after manure application. The former study used probably <5 experimental sites, and the latter probably <15. Moreover, the data presented suggest that the effect of climatic zone on SOC change after manure application was not statistically significant.

Correspondence: Denis A. Angers, tel. +418 210 5022,

fax +418 648 2402, e-mail: denis.angers@agr.gc.ca

Reproduced with the permission of the Minister of Agriculture and Agri-Food Canada

Due to the close relationship between SOC and soil quality (Gregorich *et al.*, 1994), there is strong interest in developing tools to advise farmers on the effect of manure on SOC. For that purpose, some fertilizer and crop management guidelines report retention coefficients for manures of different animal species or decomposition degree (e.g., Soltner, 2000; Clément *et al.*, 2010). The retention or isohumic coefficient is defined as the fraction of applied organic matter which is 'transformed' into soil organic matter (Hénin & Dupuis, 1945). The determination of manure C retention coefficient is also of interest for C modeling as C retention coefficient for crop residues is a substrate quality parameter used in most SOC models (Kätterer *et al.*, 2011).

The global quantification of SOC change after manure application is also relevant for national greenhouse gas (GHG) emission inventories. The Intergovernmental Panel on Climate Change (IPCC) provides default SOC stock change factors for manure application for C accounting for national GHG inventories (IPCC, 2006). These factors represent the effect on SOC stocks after regular addition of animal manure for a period of at least 20 years (IPCC, 2006). We understand that these estimates were generated from probably five individual studies, and the amount of manure-C input and management activity categories such as animal species or manure management system were not taken into account. Such considerations would represent a major improvement in estimate accuracy. Emission factors for CH₄ and N₂O are already proposed for specific animal species and manure management systems (IPCC, 2006).

There is therefore a need to refine global quantification of SOC changes following manure application from both agronomic and environmental perspectives. The objective of our work was to quantify the response of SOC stocks to manure application from a large worldwide pool of individual studies, and to assess the impact of explanatory factors such as climate, soil properties (texture, initial SOC concentration), land use and manure characteristics (C input, animal species, manure management system).

Materials and methods

Literature search and study selection

We searched literature published up to the end of 2011 using two bibliographic databases: CAB Abstracts and Scopus. Specific keywords describing animal category (*animal, pig, cattle, hog, poultry, sheep, horse, livestock*), manure (*compost, mud, sludge, ooze, effluent, waste, manure, dung, slurry, muck slurry, farmyard manure*), and soil carbon (*soil carbon sequestration, soil carbon accumulation, soil carbon content, soil carbon quantity, soil*

carbon concentration, soil carbon density, soil carbon stocks) were combined. From >1000 articles containing these keywords, we selected those which met the following criteria:

- Agronomic field experiments on cropland or grassland were included, but greenhouse, forest and mine soil reclamation experiments were excluded;
- The experimental design should be replicated;
- The soil C stocks (Mg C ha⁻¹) should be available or computable from soil C concentrations (g C kg⁻¹ or %) and bulk density values. In studies which reported soil organic matter concentration instead of C, we estimated C as 58% of the organic matter;
- Experiments should include at least one treatment with animal manure and a reference treatment which could be a control (without any fertilization) and/or a mineral fertilization treatment. Based on this criterion, two data sets (REF-zero; REF-min) were computed to evaluate SOC response to manure application in comparison to an unfertilized control and a mineral fertilization treatment, respectively. Studies could be part of one or both data set(s). Mineral fertilization had to include nitrogen (N). For the REF-min data set, manure could be applied alone or combined with mineral fertilization to carry additional N, P, and K to approach mineral supply of the mineral fertilization treatment.
- Study duration should be at least 3 years. When there were more than one article on the same experiment, the latest was considered if soil C stocks (Mg ha⁻¹) were reported in both articles. If only one reported soil C stocks, this article was selected even if it was not the latest.

Soil C stocks were recorded or computed to 30 cm depth when possible. Stocks for soil depths of <15 cm were not retained. For one site, soil C stocks were given for a whole profile of 100 cm (Srinivasarao *et al.*, 2011). Only two sites reported soil C stocks on an equivalent mass basis (Kätterer *et al.*, 2011; Viaud *et al.*, 2011). Standard deviations for each treatment were collected or evaluated from standard errors or critical value of comparison test (e.g., LSD). The sample size was also collected. The DataThief software (Tummers & Van Der Laan, 2006) was used to estimate data from graphs. In addition, for each experimental site, we compiled metadata that would be used as explanatory factors of the effect of manure application, that is, climatic data, soil properties (texture, initial SOC concentration), land use, manure properties (application rate, C concentration, C input, animal species, management system: liquid or solid storage), and study duration. If the data were not available in the article, the authors were requested to provide missing information. IPCC climate zone was determined for each experimental site from geographic coordinates and the world map of IPCC climate zones (European Commission, 2012). We also considered precipitation and temperature data in the articles to confirm the climate zone given by the world map. For two sites, cumulative manure-C input was known for a shorter study duration and was extended for the whole period of application (Singh *et al.*, 1998; Holeplass *et al.*, 2004; Bandyopadhyay *et al.*, 2011; Nayak *et al.*, 2012). For two sites, manure-C input was

evaluated with manure-C concentration which was available only for 1 year (Grandy *et al.*, 2002; Gami *et al.*, 2009).

Choice of effect size index

An effect size is a value reflecting the magnitude of the treatment effect in comparison to a reference treatment (Borenstein *et al.*, 2009). For each observation (comparison between manure and reference treatment), we calculated two effect size indices to evaluate SOC response to manure application. A first index corresponded to the difference between the SOC stock in the plot receiving manure and the SOC stock in the reference plot (Mg C ha^{-1}) thereafter referred to as *SOC stock difference*. As mentioned earlier, two data sets were created according to the reference plot, that is, a REF-zero data set when the reference plot was a control and a REF-min data set when the reference plot was a plot receiving mineral fertilization. Finally, a second index corresponded to the ratio of the SOC stock in the plot receiving manure to the SOC stock of the plot receiving mineral fertilization thereafter referred to as *relative SOC change*. This index would be comparable to the default relative SOC change factors provided by IPCC (2006) and to factors determined by Ogle *et al.* (2005) from experiments which lasted at least 20 years. This index was estimated on a reduced REF-min data set including only studies with duration of 20 years or more to be consistent and comparable with the IPCC approach.

Statistical analysis

Three categorical factors (climate, animal species, and land use) and three continuous factors [cumulative manure-C input, clay and silt content, and initial SOC concentration (g kg^{-1})] were retained for the analysis. Three climate classes were constituted from the IPCC climate classification: cool temperate, warm temperate, and tropical. Three land use classes were tested: annual crops, perennial crops, and rice paddies. For animal species, cattle and poultry manure had sufficient observations for statistical analysis of both data sets. The effect of pig manure could only be determined for the REF-zero data set and the effect of goat manure could not be tested due to lack of observations. The effect of manure management system (solid vs. liquid) could not be tested due to too few observations for liquid manure.

A mixed model was used by including a random variable for the site, to take into account the dependency of several effect sizes at a same site. Since standard deviations were rarely available to weight by the inverse of variance, the analysis was weighted by sample size. Statistical significance of each explanatory factor (cumulative manure-C input, animal species, clay and silt content, initial SOC concentration, climate, and land use) was tested independently from the others for the two indices as data sets were not complete for all explanatory factors. The assumptions of homogeneity of variance and normality of the residuals were verified graphically. For normality, Shapiro-Wilk and Kolmogorov-Smirnov tests were also verified. Heterogeneity was corrected either by modeling a different variance for each level of a categorical

factor or by modeling the variance as a power of the mean. Log transformations were carried out for some cases. The statistical significance between means of the different levels of categorical factors was given by least square differences. For continuous factors, R-Squared was calculated using a method recommended for mixed models (Edwards *et al.*, 2008). As data sets were not complete for all categorical and continuous factors, the number of observations was specified on the figures. A Student's *t*-test was used to test the statistical significance of global means for both indices. All statistical analyses were performed using the SAS software, version 9.2 (SAS Institute, Cary, NC, USA). Statistical results were considered to be significant at the 0.05 α level.

Results and discussion

General results

By applying our selection criteria, 42 articles were retained for 49 sites in the world (Table 1). Among these 49 replicated sites, 40 were randomized, eight were presumed to be randomized based on the description of the experimental design, and one was not randomized. The soil depth considered varied from 15 to 100 cm, with an average of 26 cm. Duration of the studies ranged from 3 to 82 years, with an average of 18 years (18 years also for studies with known cumulative manure-C input).

The 130 observations from the 49 sites were separated into REF-zero and REF-min data sets to evaluate the SOC response to manure application in comparison to an unfertilized control and to mineral fertilization, respectively. The *SOC stock difference* (1st index) was calculated for every observation of both data sets. In total for this effect size, 36 sites (57 observations) were available for the REF-zero data set and 43 (73 observations) for the REF-min data set. Among the 43 sites included in the REF-min data set, 22 (28 observations) had duration of ≥ 20 years and were used to analyze the *relative SOC change* (2nd index).

Overall effect of manure application on SOC stocks compared to a reference

Overall, as expected, the positive response in the *SOC stock difference* (first index) indicated a significantly larger SOC stock in plots receiving animal manure than in both reference treatments (Table 2). The higher SOC stock in manured treatments could be due to direct C input by the manure itself and indirect C input through increased net primary production (including roots and crop residues) (Aoyama *et al.*, 1999; Whalen & Chang, 2002; Bhattacharyya *et al.*, 2010). The average SOC response to manure application was larger when using control plots as the reference (9.4 Mg C ha^{-1}) than

Table 1 Summary of data for the sites included in the analysis

Location	Data set	<i>n</i> *	Sampling depth (cm)	Duration (years)	Cumulative manure-C input (Mg C ha ⁻¹)	Animal species	IPCC climatic zone	Soil clay (%)	Soil silt (%)	Initial SOC (g kg ⁻¹)	Land use	Reference†
Saint-Lambert de Lévis, Qc, Canada	REF-zero	2	30	20	NA	pig	cool temperate	34	56	NA	perennial	(Angers <i>et al.</i> , 2010)
Ludhiana, Punjab, India	REF-zero, REF-min	4	15	7	28	NA	tropical	17	24	5	paddy	(Benbi & Senapati, 2010)
Almora, Uttarakhand, India	REF-zero, REF-min	2	30	8	13	cattle	warm temperate	34	56	7	annual	(Bhattacharyya <i>et al.</i> , 2007, 2009); pers. comm.
Meden Vale, Nottinghamshire, UK	REF-zero	5	30	4	6;13;19;25;32	poultry	cool temperate	6	8	11	annual	(Bhagal <i>et al.</i> , 2009); (Bhagal & Shepherd, 1997)
Foulum, Denmark	REF-zero	1	30	11	2	pig	cool temperate	9	13	23	annual	(Chirinda <i>et al.</i> , 2010)
Koraput, Orissa, India	REF-min	4	15	3	NA	cattle	tropical	19	24	6	annual	(Dass <i>et al.</i> , 2008)
Waldviertel, Austria	REF-zero, REF-min	2	25	21	NA	NA	cool temperate	8	13	NA	annual	(Dersch & Bohm, 2001)
Alpenvorland, Austria	REF-zero, REF-min	2	25	21	NA	NA	cool temperate	50	40	NA	annual	(Dersch & Bohm, 2001)
Pingliang, Gansu, China	REF-zero, REF-min	2	20	26	20	cattle	warm temperate	34	43	5	annual	(Fan <i>et al.</i> , 2008); pers. comm.
Farmington, Georgia, USA	REF-min	4	20	12	29	poultry	warm temperate	13	18	NA	perennial	(Franzluebbers & Stuedemann, 2010); (Franzluebbers & Stuedemann, 2009); pers. comm.
Bhairahawa, Nepal	REF-zero, REF-min	2	30	25	NA	cattle	tropical	18	71	NA	paddy	(Gami <i>et al.</i> , 2009); pers. comm.
Tarahara, Nepal	REF-zero, REF-min	2	30	25	NA	cattle	tropical	11	58	NA	paddy	(Gami <i>et al.</i> , 2009); pers. comm.
Parwanipur, Nepal	REF-zero, REF-min	3	30	23	22	cattle	tropical	14	58	NA	paddy	(Gami <i>et al.</i> , 2009); pers. comm.
Presque Isle, Maine, USA	REF-min	1	15	6	26	cattle	cool temperate	NA	NA	NA	annual	(Grandy <i>et al.</i> , 2002)
Isfahan, Isfahan, Iran	REF-zero, REF-min	6	20	7	44;87;175	cattle	warm temperate	34	56	5	annual	(Hemmat <i>et al.</i> , 2010)

Table 1 (Continued)

Location	Data set	<i>n</i> *	Sampling depth (cm)	Duration (years)	Cumulative manure-C input (Mg C ha ⁻¹)	Animal species	IPCC climatic zone	Soil clay (%)	Soil silt (%)	Initial SOC (g kg ⁻¹)	Land use	Reference†
Akershus, Norway	REF-min	3	25	48	29	cattle	cool temperate	20	40	40	annual; perennial	(Holeplant <i>et al.</i> , 2004); (Singh <i>et al.</i> , 1998)
Nanchang, Jiangxi, China	REF-zero, REF-min	2	15	22	67	pig	warm temperate	26	NA	9	annual	(Huang <i>et al.</i> , 2010a); (Huang <i>et al.</i> , 2010b); pers. comm.
Prairie du Sac, Wisconsin, USA	REF-min	1	30	4	NA	cattle	cool temperate	15	60	NA	annual	(Jokela <i>et al.</i> , 2009)
Nairobi, Kenya	REF-zero, REF-min	4	20	19	20;39	cattle	warm temperate	67	22	20	annual	(Kamoni <i>et al.</i> , 2007); (Kapkiyai <i>et al.</i> , 1999); (Kibunja <i>et al.</i> , 2010)
Embu, Kenya	REF-zero	2	20	13	20;41	goat	tropical	31	13	NA	annual	(Kamoni <i>et al.</i> , 2007); (Kihanda <i>et al.</i> , 2005)
Uppsala, Sweden	REF-zero, REF-min	2	20‡	53	101	cattle	cool temperate	37	41	15	annual	(Kätterer <i>et al.</i> , 2011); (Kirchmann <i>et al.</i> , 2004)
Taipei, Taiwan	REF-zero	2	20	3	NA	pig	tropical	14	66	14	annual	(Lee <i>et al.</i> , 2006)
Brookings, South Dakota, USA	REF-zero, REF-min	4	30	4	12;23	cattle	cool temperate	32	62	NA	perennial	(Lee <i>et al.</i> , 2007)
Therwil, Switzerland	REF-zero, REF-min	2	20	27	24;32	cattle	cool temperate	16	72	15	annual	(Leifeld <i>et al.</i> , 2009); (Fliessbach <i>et al.</i> , 2007)
Yangling, Shaanxi, China	REF-min	1	20	17	NA	cattle	warm temperate	17	52	6	annual	(Liang <i>et al.</i> , 2011); (Zhao <i>et al.</i> , 2010)
Shenyang, Liaoning, China	REF-zero, REF-min	3	20	22	44;22	pig	cool temperate	25	58	9	annual	(Lou <i>et al.</i> , 2011)
Padova, Italy	REF-zero, REF-min	4	30	36	155;129	cattle	warm temperate	15	38	12	annual	(Morari <i>et al.</i> , 2006); pers. comm.
Parbhani, Maharashtra, India	REF-min	2	15	7	NA	NA	tropical	53	16	5	annual	(More & Hangarge, 2003)

Table 1 (Continued)

Location	Data set	<i>n</i> *	Sampling depth (cm)	Duration (years)	Cumulative manure-C input (Mg C ha ⁻¹)	Animal species	IPCC climatic zone	Soil clay (%)	Soil silt (%)	Initial SOC (g kg ⁻¹)	Land use	Reference†
Chinon, France	REF-zero	2	30	28	23;46	cattle	warm temperate	9	5	8	perennial	(Morlat & Chaussod, 2008); pers. comm.
Cuttack, Orissa, India	REF-zero, REF-min	2	15	35	32	cattle	tropical	20	14	7	paddy	(Nayak <i>et al.</i> , 2009)
Kanpur, Uttar Pradesh, India	REF-min	1	30	25	NA	NA	tropical	18	35	3	paddy	(Nayak <i>et al.</i> , 2012)
Sabour, Bihar, India	REF-min	1	30	25	NA	NA	tropical	28	22	5	paddy	(Nayak <i>et al.</i> , 2012)
Kalyani, West Bengal, India	REF-min	1	30	23	12	cattle	tropical	50	30	9	paddy	(Nayak <i>et al.</i> , 2012); (Bandyopadhyay <i>et al.</i> , 2011); pers. comm.
Ludhiana, Punjab, India	REF-min	1	30	26	NA	cattle	tropical	18	28	3	paddy	(Nayak <i>et al.</i> , 2012); (Walia <i>et al.</i> , 2010)
Pendleton, Oregon, USA	REF-zero, REF-min	2	30	56	37	cattle	warm temperate	18	70	NA	annual	(Rasmussen & Parton, 1994)
Ludhiana, Punjab, India	REF-zero, REF-min	2	30	32	NA	NA	tropical	7	33	3	paddy	(Rasool <i>et al.</i> , 2007); (Kukul <i>et al.</i> , 2009)
Ludhiana, Punjab, India	REF-zero, REF-min	2	30	32	NA	NA	tropical	16	24	3	annual	(Rasool <i>et al.</i> , 2008); (Kukul <i>et al.</i> , 2009)
Almora, Uttarakhand, India	REF-zero, REF-min	8	15	3	34;27;20;14;22	cattle	warm temperate	34	56	11	annual	(Saha <i>et al.</i> , 2008)
Bhubaneswar, Orissa, India	REF-zero, REF-min	6	30	3	NA	cattle	tropical	10	25	NA	perennial	(Saha <i>et al.</i> , 2010b)
Umiam, Meghalaya, India	REF-min	1	30	5	NA	NA	tropical	34	34	NA	annual	(Saha <i>et al.</i> , 2010a)
Belle Mina, Alabama, USA	REF-zero, REF-min	4	20	10	11	poultry	warm temperate	27	58	12	annual	(Sainju <i>et al.</i> , 2008a); (Sainju <i>et al.</i> , 2010); (Sainju <i>et al.</i> , 2008b)
Varanasi, Uttar Pradesh, India	REF-zero, REF-min	3	15	9	22;11	NA	tropical	62	22	1	annual	(Sarkar <i>et al.</i> , 2003)
Akola, Maharashtra, India	REF-zero, REF-min	3	15	19	NA	NA	tropical	NA	NA	NA	annual	(Sharma <i>et al.</i> , 2011)

Table 1 (Continued)

Location	Data set	n*	Sampling depth (cm)	Duration (years)	Cumulative manure-C input (Mg C ha ⁻¹)	Animal species	IPCC climatic zone	Soil clay (%)	Soil silt (%)	Initial SOC (g kg ⁻¹)	Land use	Reference†
Patna, Bihar, India	REF-zero, REF-min	3	15	4	NA	NA	tropical	74	9	5	paddy	(Singh <i>et al.</i> , 2001)
Sardar Krushinagar, Gujarat, India	REF-zero, REF-min	5	100	18	27;10	cattle	tropical	11	4	2	annual	(Srinivasarao <i>et al.</i> , 2011); pers. comm.
Zhangye, Gansu, China	REF-zero, REF-min	2	20	23	52	pig-cattle-sheep	cool temperate	20	52	12	annual	(Su <i>et al.</i> , 2006); (Hai <i>et al.</i> , 2010)
Lucknow, Uttar Pradesh, India	REF-zero, REF-min	2	30	5	23	NA	tropical	13	25	3	perennial	(Suman <i>et al.</i> , 2009); (Singh <i>et al.</i> , 2007)
Bignan, France	REF-min	1	30‡	8	18	poultry	warm temperate	17	42	25	annual	(Viaud <i>et al.</i> , 2011); pers. comm.
Moystad, Norway	REF-zero, REF-min	4	20	82	NA	cattle	cool temperate	14	34	24	annual	(Yang <i>et al.</i> , 2007)

*n = number of observations.

†for each site, the first cited reference included soil C stocks. The other references contained unavailable information in the first one.

‡Equivalent topsoil depth of the study reference treatment.

Table 2 Global means of the two effect size indices with the 95% Confidence Interval (CI) and probability level corresponding to the Student's *t*-test

Index	REF-zero data set	REF-min data set
First: SOC stock difference (Mg C ha ⁻¹)	9.4 ± 4.1 (95% CI); <i>P</i> < 0.0001	5.6 ± 2.8 (95% CI); <i>P</i> < 0.0001
Second: relative SOC change		1.26 ± 0.14 (95% CI); <i>P</i> < 0.0001

mineral fertilizer (5.6 Mg C ha⁻¹). If we assume that in most studies of the REF-min data set the amount of C added as roots and crop residues was similar in the manure treatment and in the mineral fertilizer reference treatment, thus the positive SOC response following manure application can probably be mainly attributed to direct C input by manure.

Factors influencing the SOC stock difference

Our meta-analysis showed a significant linear relationship between the *SOC stock difference* (first index; Mg C ha⁻¹) and cumulative manure-C input (Mg C ha⁻¹) for the REF-zero (*P* < 0.0001; *R*² = 0.53) and the REF-min (*P* < 0.0001; *R*² = 0.59) data sets (Fig. 1a, b). In this analysis, cumulative manure-C input is a combination of annual manure-C application rate and study duration. Positive linear relationships between cumulative amount of added organic matter and SOC stocks or stock changes associated with treatments involving various amounts of manure or other organic C sources have often been observed in individual studies (e.g., Thomsen & Christensen, 2004; Kong *et al.*, 2005; Majumder *et al.*, 2008). As far as we know, our study is the first one reporting such a relationship between *SOC stock difference* compared to a reference treatment and cumulative animal manure-C input from a worldwide set of studies. **It is noteworthy that despite widely variable sources of data (soil, climate, study duration, animal species, land use, etc.), we can explain at least 53% of the variability in SOC stock difference by cumulative manure-C input.** For comparison purposes, we added eight individual observations (not included in our meta-analysis due to lack of replications) from seven long-term sites in the United Kingdom (Jenkinson & Johnston, 1977; Jenkinson *et al.*, 1990), Germany (Ludwig *et al.*, 2011), Canada (Grant *et al.*, 2001), and China (Guo *et al.*, 2007) (Fig. 1a, b). These points clearly fit in the same general linear relationship as the data retained in our meta-analysis. Our results corroborate linear increases in SOC stocks with increasing organic C inputs shown by most current models of SOC dynamics




Fig. 1 Relationship between SOC stock difference and cumulative manure-C input for the REF-zero (a) and REF-min (b) data sets (•), and with additional observations from non-replicated sites (×).




Fig. 2 Effect of climatic zone on the SOC stock difference due to manure application for the REF-zero (a) and REF-min (b) data sets [number of observations in each climate class in brackets, means and 95% Confidence Interval (CI)].

[e.g., RothC (Coleman & Jenkinson, 1995), Century (Parton *et al.*, 1987)]. Indeed, the relationship between C input and SOC stock difference is linear up to very high levels of cumulative C input.

The group analysis indicated a strong trend toward an effect of climatic zone on the SOC stock difference for both the REF-zero ($P = 0.062$) and the REF-min ($P = 0.060$) data sets (Fig. 2a,b). The SOC stock difference was lower in tropical climate than in warm temperate climate for the REF-zero ($P = 0.027$) and REF-min ($P = 0.052$) data sets. In addition, for the REF-min data set, the SOC stock difference tended to be lower in tropical climate than in cool temperate climate ($P = 0.078$) (Fig. 2b). This observation is consistent with the general understanding that soils in warmer climates, where decomposition is faster, may accumulate C slower than soils in colder climates (Freibauer *et al.*, 2004). However, it should be noted that for tropical climate, a significant number of observations (10 of 21 for the

REF-zero data set and 12 of the 33 for the REF-min data set) actually had a cumulative manure-C input <45 Mg C ha⁻¹ and that cumulative manure-C input was unknown for the other observations. Therefore, from our meta-analysis, we cannot rule out that the lower SOC stocks under tropical climate may simply be due to relatively lower cumulative manure-C inputs. This clearly emphasizes the importance of documenting C inputs in agronomic studies reporting SOC change.

The effect of animal species on the SOC stock difference was not statistically significant (Fig. 3a,b). However, in both REF-zero and REF-min data sets, cattle manure showed a slight trend to higher values than the others, and confidence intervals suggested a significant positive SOC stock difference after cattle manure application but not after pig and poultry manure application. This significant response of SOC after cattle manure application would be consistent with the idea that its organic




Fig. 3 Effect of animal species on the SOC stock difference due to manure application for the REF-zero (a) and REF-min (b) data sets [number of observations in each species class in brackets, means and 95% Confidence Interval (CI)].

matter is more stable than that from poultry and pig manure (Velthof *et al.*, 2000). However, it should be underlined that compared to cattle, the number of observations was very low and the variability very high for pig and poultry manures, which limits the interpretation. In addition, similar to climate, we cannot rule out that the low values for pig and poultry manure may also be related to a relatively low cumulative manure-C input associated with these studies (<45 Mg C ha⁻¹, for all relevant studies except one). Clearly, more studies on the impact of pig and poultry manure are required, and these should include C input data.

We found no effects of land use, clay and silt content, or initial SOC concentration on the SOC stock difference. This is partly due to missing observations in the case of initial SOC concentration (Table 1) or to the limited number of studies involving 'perennial' and 'paddy' cropping systems in the case of land use. In addition, these explanatory factors may be of minor significance compared to the overwhelming effect of cumulative manure-C input.

Estimation of manure-C retention coefficient

Overall, our global analysis showed that cumulative manure-C input was the main factor explaining SOC response to manure application. Many studies have estimated C retention coefficients from the slope of linear relationships between cumulative C input and SOC stocks or stock changes associated with experimental treatments including various amounts of manure or crop residues (Campbell *et al.*, 1991; Bhogal & Shepherd, 1997; Buyanovsky & Wagner, 1998; Thomsen & Christensen, 2004; Kong *et al.*, 2005; Mandal *et al.*, 2007; Benbi & Senapati, 2010). We outlined previously that in the case of the REF-min data set, the positive SOC response to manure application can probably be

mainly attributed to C input by manure. Thus, the slope of the linear relationship between the cumulative manure-C input (Mg C ha⁻¹) and the SOC stock difference (1st index; Mg C ha⁻¹) for the REF-min data set (Fig. 1b) can be considered an approximation of the manure-C retention coefficient, which represents the average proportion of manure-C remaining in soil. Our estimated global manure-C retention coefficient is 12% ± 4 (95% Confidence Interval, CI) for an average study duration of 18 years. As described earlier, by adding eight individual observations from seven long-term sites (without replicates) in the United Kingdom (Jenkinson & Johnston, 1977; Jenkinson *et al.*, 1990), Germany (Ludwig *et al.*, 2011), Canada (Grant *et al.*, 2001), and China (Guo *et al.*, 2007), the slope remains at 12% ± 2 (95% CI).

To the best of our knowledge, this is the first time that a C retention coefficient of animal manure can be deduced from a worldwide set of studies offering a wide range of cumulative C input levels (10–175 Mg C ha⁻¹ for observations included in the meta-analysis; 3–402 Mg C ha⁻¹ with additional observations from unreplicated designs). Indeed, all of the studies cited previously involved individual sites. Moreover, the C input estimates in these studies included either crop residues only or a mix of crop residues and manure. Using 14 long-term field experiments in Europe, Smith *et al.* (1997) established a linear relationship between yearly relative changes in SOC stock compared to a reference treatment and the amount of fresh organic manure added annually. However, the absence of manure dry matter and C contents did not allow us to calculate the cumulative manure-C input and consequently a C retention coefficient from their data.

Our estimated manure-C retention coefficient is lower than the value of 23% reported for animal manure in a study from seven different sites in the United Kingdom (Bhogal *et al.*, 2007). However, their 95% CI

was 9 to 37 which overlaps with ours. In addition, Bhojal *et al.* (2007) pointed out that the average increase of SOC change representing 23% of manure-C could only be regarded as the initial rate of SOC increase, as SOC accumulation rates decline over time, and consequently their retention coefficient could be overestimated. Moreover, their value was measured by calculating the difference in SOC between manured and unmanured treatments without specifying if the unmanured treatment included mineral fertilization or not. Thus, their manure-C retention coefficient could also be overestimated if some C in soils originated from crop residues rather than manure.

Our manure-C retention coefficient can be compared to those estimated for crop residues. After 18 years of annual straw addition, Thomsen & Christensen (2004) observed a retention coefficient of 14% for straw C. Campbell *et al.* (1991) reported a C retention coefficient of 6% for crop residues on a 30-year old site. After 10 years, Kong *et al.* (2005) observed a value of 8% across ten different cropping systems. Manure was also applied in one of their cropping systems. From Buyanovsky & Wagner (1998), we calculated that 6% and 7% of C applied as crop residues, and additionally from manure in some plots, were retained in soil during a very long period of 'humification' extending to 100 years. Therefore, our value of manure-C retention coefficient ($12\% \pm 4$), estimated for an average study duration of 18 years, is comparable to those for crop residues determined in the medium to long-term (10–100 years), and supports Buyanovsky & Wagner (1998) who argued that there was no reason to consider that manure was more effective for SOM enhancement than plant residues. However, retention of C added as manure may be greater than crop residues in the short term. For instance, in an *in situ* decomposition study, Thomsen & Christensen (2010) observed a higher retention of C added in the form of sheep feces (30%) than C directly applied as crop residues (19%) after nine annual additions. Likewise, in a laboratory incubation, 14% of C applied in feed was retained in soil after 1 to 2 years, against 48% of C applied as feces (Thomsen *et al.*, 2013). Taken together, our results and the above-cited literature suggest that despite apparently faster early decomposition of crop residues compared to manure, longer term stabilization may be relatively similar for both.

Estimation of relative SOC change after manure application

The second effect size index is defined as the ratio of SOC stock in plots receiving manure to SOC stock in plots receiving mineral fertilization. It represents a

relative SOC change factor such as used by IPCC (2006) and Ogle *et al.* (2005), and estimated from experiments that lasted at least 20 years. The average value for the relative SOC change (2nd index) was 1.26 ± 0.14 (95% CI) for experiments that lasted at least 20 years (Table 2). This ratio was lower than the average of IPCC relative SOC change factors (1.41) proposed for high inputs with manure (IPCC, 2006) and the average of high input factors with amendments (1.36) proposed by Ogle *et al.* (2005). Nevertheless, our estimation was based on a larger number of sites (22 sites for 28 observations) against probably five studies for IPCC (2006) and less than five for Ogle *et al.* (2005).

A significant linear relationship was also observed between the cumulative manure-C input (Mg C ha^{-1}) and the relative SOC change ($P = 0.026$; $R^2 = 0.31$) (Fig. 4). Climatic zone did not affect the relative SOC change (Fig. 5). Similar to the SOC stock difference (1st index), no effect of land use, clay and silt content or initial SOC concentration was observed. The reduced REF-min data set was limited by the low number of observations for cumulative manure-C input, and some climate and animal species classes. Clearly, as mentioned previously, more data are necessary to elucidate the global impact of these explanatory factors.

The IPCC Guidelines for National Greenhouse Gas Inventories provide different values of soil C stock change after animal manure application in different climatic zones: 1.37 for temperate, boreal, and tropical dry climates; 1.44 for temperate, boreal, and tropical moist/wet climates; and 1.41 for tropical montane climates (IPCC, 2006). From a statistical standpoint, these values




Fig. 4 Relationship between the relative SOC change and cumulative manure-C input for the reduced REF-min data set (●), and with additional observations from non-replicated sites (x).




Fig. 5 Effect of climatic zone on the *relative SOC change* for the reduced REF-min data set [number of observation in each climate class in brackets, means and 95% Confidence Interval (CI)].

are not significantly different from each other. Moreover, the amount of C input by manure is not taken into account in these estimates. Our results offer some scope for refinement of these change factors by taking into account the amount of manure applied. However, the relationship between the cumulative manure-C input (Mg C ha^{-1}) and the *relative SOC change* probably needs to be confirmed with more data to propose a new operational set of IPCC factors.

Limits and outlook

To the best of our knowledge, this study is the first global analysis of factors determining SOC response to manure application which includes a significant number of studies. For this meta-analysis, we chose to work with high-quality data (replicated randomized experiments, values expressed as SOC stocks, C input from manure, etc.). Consequently, our conclusions were constrained by missing data and the meta-analysis raises many points that need to be improved in future research.

First, SOC expressed as stock or bulk density data necessary to compute SOC stock were missing in many articles that were scanned. This information is essential to examine the relationship between cumulative manure-C input and SOC change. Ideally, SOC stocks should also be expressed on an equivalent mass basis to avoid any bias in the estimation of changes in SOC stocks when the entire soil profile is not sampled (in case there would be a significant amount of SOC beneath the maximum depth of sampling) (Vandenbygaert & Angers, 2006). In the studies considered in our analysis, only two reported SOC stocks on an equivalent mass (Kätterer *et al.*, 2011; Viaud *et al.*, 2011).

Another important point that limited our analysis was detailed information allowing estimation of

manure-C input (C concentration, application rate, dry matter content) which was absent from many articles reporting SOC stocks. Consequently, manure-C input could be estimated in only 29 of the 49 sites, representing a total of 86 observations in both data sets. This information is necessary to analyze the relationship between SOC change and cumulative manure-C input and refine the estimation of our manure-C retention coefficient. Indeed, there is uncertainty associated with this relationship due to the substantial lack of observations with cumulative manure-C input greater than 75 Mg C ha^{-1} (Fig. 1a,b and Fig. 4). In addition, there is significant variability in our data for cumulative manure-C input $<75 \text{ Mg C ha}^{-1}$. However, these data represent the current state of available information for studies on SOC response to manure application.

Other relevant information for our analysis such as animal species and initial SOC concentration was often missing in the scanned articles (Table 1). In addition, some classes of explanatory factors had a low number of observations compared to others. For example, fewer studies used poultry or pig manure compared to cattle manure. Similarly, we did not find enough studies involving liquid manure compared to solid. Perennial cropping systems were less represented than annual systems. Another limitation was the lack of tropical sites outside of India. So extrapolating our results to other tropical regions should be done with caution. Most sites were located in Europe, Asia, or North America with South America, Africa and Oceania being under- or not represented.

The results of this meta-analysis show that at a global scale cumulative manure-C input has a dominant effect on SOC change following manure application. When more data become available, the effects of explanatory factors like climate, animal manure characteristics, soil clay and silt content, initial SOC concentration and land use could be studied further by determining their effect on the slope of the relationship between cumulative C input and SOC stock change using meta-regression with multiple explanatory factors.

In accordance with our selection criteria, many short-term studies (<3 years) addressing manure impact on SOC were omitted from the meta-analysis (Lynch *et al.*, 2005; Agbede, 2010; Bilalis & Karamanos, 2010; Mellek *et al.*, 2010; Islam *et al.*, 2011). Most of these studies mentioned specific information on manure characteristics including manure-C input (Lynch *et al.*, 2005; Agbede, 2010; Mellek *et al.*, 2010) which is encouraging for future meta-analyses, and we can only hope that these studies will be maintained.

In summary, our findings suggest that at a global scale, above animal species, land use or soil properties, the amount of applied manure-C was the dominant

driver determining the extent of SOC increase following manure application. We also observed a strong tendency for *SOC stock difference* to be lower in tropical than other climatic zones, but this effect could not be decoupled from that of manure-C input. Our results for the relative change factors, but also for manure-C retention coefficient, offer some scope for the improvement of national GHG inventory methodologies. There were obvious gaps in the data especially with regards to animal species (especially pig and poultry manure), manure management system (e.g., liquid vs. solid storage), and under-representation of some geographic regions. We emphasize that long-term field studies covering these gaps either be initiated or continued, and that these studies document important explanatory factors such as manure application rate and C concentration which were very often absent from research articles.

Acknowledgements

We thank authors who shared additional information on studies chosen for the meta-analysis. We thank Léon-Étienne Parent for advice on meta-analysis methods, Martin Chantigny, and Philippe Rochette for helpful discussions, Julien Beguin and Anne-Sophie Julien for statistical assistance, and Anaïs Charles and Caroline Hudon for help with the literature search. We are grateful to Jérôme Laganière and Frank Larney who provided comments on a draft version of the manuscript. We acknowledge the support of Agriculture and Agri-Food Canada and the Fonds Québécois de la Recherche sur la Nature et les Technologies (FQRNT) for a post-graduate scholarship to one of us (É.M.).

References

- Agbede TM (2010) Tillage and fertilizer effects on some soil properties, leaf nutrient concentrations, growth and sweet potato yield on an Alfisol in southwestern Nigeria. *Soil and Tillage Research*, **110**, 25–32.
- Angers D, Chantigny M, Macdonald J, Rochette P, Côté D (2010) Differential retention of carbon, nitrogen and phosphorus in grassland soil profiles with long-term manure application. *Nutrient Cycling in Agroecosystems*, **86**, 225–229.
- Aoyama M, Angers DA, N'dayegamiye A (1999) Particulate and mineral-associated organic matter in water-stable aggregates as affected by mineral fertilizer and manure applications. *Canadian Journal of Soil Science*, **79**, 295–302.
- Bandyopadhyay PK, Saha S, Mallick S (2011) Comparison of soil physical properties between a permanent fallow and a long-term rice-wheat cropping with inorganic and organic inputs in the humid subtropics of eastern India. *Communications in Soil Science and Plant Analysis*, **42**, 435–449.
- Benbi DK, Senapati N (2010) Soil aggregation and carbon and nitrogen stabilization in relation to residue and manure application in rice-wheat systems in northwest India. *Nutrient Cycling in Agroecosystems*, **87**, 233–247.
- Bhattacharyya R, Chandra S, Singh RD, Kundu S, Srivastva AK, Gupta HS (2007) Long-term farmyard manure application effects on properties of a silty clay loam soil under irrigated wheat-soybean rotation. *Soil & Tillage Research*, **94**, 386–396.
- Bhattacharyya R, Prakash V, Kundu S, Pandey SC, Srivastva AK, Gupta HS (2009) Effect of fertilisation on carbon sequestration in soybean-wheat rotation under two contrasting soils and management practices in the Indian Himalayas. *Australian Journal of Soil Research*, **47**, 592–601.
- Bhattacharyya R, Prakash V, Kundu S, Srivastva AK, Gupta HS, Mitra S (2010) Long term effects of fertilization on carbon and nitrogen sequestration and aggregate associated carbon and nitrogen in the Indian sub-Himalayas. *Nutrient Cycling in Agroecosystems*, **86**, 1–16.
- Bhogal A, Shepherd M (1997) Effect of poultry manure on the leaching of carbon from a sandy soil as a potential substrate for denitrification in the subsoil. *Journal of the Science of Food and Agriculture*, **74**, 313–322.
- Bhogal A, Chambers BJ, Whitmore AP, Powlson DS (2007) The effects of reduced tillage practices and organic material additions on the carbon content of arable soils. Scientific report for Defra Project, SP0561, ADAS UK Ltd., 48 pp. Available at: <http://randd.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None&Completed=2&ProjectID=15162> (accessed 6 February 2013).
- Bhogal A, Nicholson FA, Chambers BJ (2009) Organic carbon additions: Effects on soil bio-physical and physico-chemical properties. *European Journal of Soil Science*, **60**, 276–286.
- Bilalis DJ, Karamanos AJ (2010) Organic maize growth and mycorrhizal root colonization response to tillage and organic fertilization. *Journal of Sustainable Agriculture*, **34**, 836–849.
- Borenstein M, Hedges LV, Higgins JPT, Rothstein HR (2009) *Introduction to Meta-Analysis*. John Wiley & Sons Ltd, Chichester, West Sussex.
- Buyanovsky GA, Wagner GH (1998) Carbon cycling in cultivated land and its global significance. *Global Change Biology*, **4**, 131–141.
- Campbell CA, Zentner RP, Bowren KE, Townley-Smith L, Schnitzer M (1991) Effect of crop rotations and fertilization on soil organic matter and some biochemical properties of a thick Black Chernozem. *Canadian Journal of Soil Science*, **71**, 377–387.
- Chirinda N, Olesen JE, Porter JR, Schjonning P (2010) Soil properties, crop production and greenhouse gas emissions from organic and inorganic fertilizer-based arable cropping systems. *Agriculture, Ecosystems & Environment*, **139**, 584–594.
- Clément MF, Angers D, Bolinder MA, N'dayegamiye A, Parent LE (2010) La gestion de la matière organique. In: *Guide de référence en fertilisation - 2ème édition* (eds Parent LE, Gagné G), pp. 55–70. Centre de référence en agriculture et agroalimentaire du Québec, Québec.
- Coleman K, Jenkinson DS (1995) *RothC-26.3 - A model for the turnover of carbon in soil: Model description and users guide*, IACR Rothamsted, Harpenden.
- Dass A, Lenka NK, Patnaik US, Sudhishri S (2008) Integrated nutrient management for production, economics, and soil improvement in winter vegetables. *International Journal of Vegetable Science*, **14**, 104–120.
- Dersch G, Böhm K (2001) Effects of agronomic practices on the soil carbon storage potential in arable farming in Austria. *Nutrient Cycling in Agroecosystems*, **60**, 49–55.
- Edwards LJ, Muller KE, Wolfinger RD, Qaqish BF, Schabenberger O (2008) An R^2 statistic for fixed effects in the linear mixed model. *Statistics in Medicine*, **27**, 6137–6157.
- European Commission (2012) Soil projects - support to renewable energy directive. Available at: <http://eusoils.jrc.ec.europa.eu/projects/RenewableEnergy/> (accessed 3 February 2012).
- Fan T, Xu M, Song S, Zhou G, Ding L (2008) Trends in grain yields and soil organic C in a long-term fertilization experiment in the China Loess Plateau. *Journal of Plant Nutrition and Soil Science*, **171**, 448–457.
- Fliessbach A, Oberholzer HR, Gunst L, Mäder P (2007) Soil organic matter and biological soil quality indicators after 21 years of organic and conventional farming. *Agriculture, Ecosystems & Environment*, **118**, 273–284.
- Franzuebbers AJ, Doraiswamy PC (2007) Chapter 18 - Carbon sequestration and land degradation. In: *Climate and land degradation* (eds Sivakumar MVK, Ndiang'ui N), pp. 343–358. Springer, Berlin.
- Franzuebbers AJ, Stuedemann JA (2009) Soil-profile organic carbon and total nitrogen during 12 years of pasture management in the Southern Piedmont USA. *Agriculture, Ecosystems & Environment*, **129**, 28–36.
- Franzuebbers AJ, Stuedemann JA (2010) Surface soil changes during twelve years of pasture management in the Southern Piedmont USA. *Soil Science Society of America Journal*, **74**, 2131–2141.
- Franzuebbers AJ, Stuedemann JA, Wilkinson SR (2001) Bermudagrass management in the Southern Piedmont USA: I. Soil and surface residue carbon and sulfur. *Soil Science Society of America Journal*, **65**, 834–841.
- Freibauer A, Rounsevell MDA, Smith P, Verhagen J (2004) Carbon sequestration in the agricultural soils of Europe. *Geoderma*, **122**, 1–23.
- Gami SK, Lauren JG, Duxbury JM (2009) Soil organic carbon and nitrogen stocks in Nepal long-term soil fertility experiments. *Soil & Tillage Research*, **106**, 95–103.
- Grandy AS, Porter GA, Erich MS (2002) Organic Amendment and Rotation Crop Effects on the Recovery of Soil Organic Matter and Aggregation in Potato Cropping Systems. *Soil Science Society of America Journal*, **66**, 1311–1319.
- Grant RF, Juma NG, Robertson JA, Izaurralde RC, McGill WB (2001) Long-term changes in soil carbon under different fertilizer, manure, and rotation: testing the

- mathematical model ecosystems with data from the Breton plots. *Soil Science Society of America Journal*, **65**, 205–214.
- Gregorich EG, Carter MR, Angers DA, Monreall CM, Ellert BH (1994) Towards a minimum data set to assess soil organic matter quality in agricultural soils. *Canadian Journal of Soil Science*, **74**, 367–385.
- Grignani C, Zavattaro L, Sacco D, Monaco S (2007) Production, nitrogen and carbon balance of maize-based forage systems. *European Journal of Agronomy*, **26**, 442–453.
- Guo L, Falloon P, Coleman K, Zhou B, Li Y, Lin E, Zhang F (2007) Application of the RothC model to the results of long-term experiments on typical upland soils in northern China. *Soil Use and Management*, **23**, 63–70.
- Hai L, Li XG, Li FM, Suo DR, Guggenberger G (2010) Long-term fertilization and manuring effects on physically-separated soil organic matter pools under a wheat-wheat-maize cropping system in an arid region of China. *Soil Biology and Biochemistry*, **42**, 253–259.
- Hemmat A, Aghilinategh N, Rezaiejad Y, Sadeghi M (2010) Long-term impacts of municipal solid waste compost, sewage sludge and farmyard manure application on organic carbon, bulk density and consistency limits of a calcareous soil in central Iran. *Soil and Tillage Research*, **108**, 43–50.
- Hénin S, Dupuis M (1945) Essai de bilan de la matière organique du sol. *Annales Agronomiques*, **15**, 17–29.
- Holeppass H, Singh BR, Lal R (2004) Carbon sequestration in soil aggregates under different crop rotations and nitrogen fertilization in an inceptisol in southeastern Norway. *Nutrient Cycling in Agroecosystems*, **70**, 167–177.
- Huang S, Peng X, Huang Q, Zhang W (2010a) Soil aggregation and organic carbon fractions affected by long-term fertilization in a red soil of subtropical China. *Geoderma*, **154**, 364–369.
- Huang S, Zhang W, Yu X, Huang Q (2010b) Effects of long-term fertilization on corn productivity and its sustainability in an Ultisol of southern China. *Agriculture, Ecosystems & Environment*, **138**, 44–50.
- IPCC (2006) *2006 IPCC Guidelines for national greenhouse gas inventories*. IGES, Japan.
- Islam MM, Majid NM, Karim AJMS, Jahiruddin M, Islam MS, Hakim MA (2011) Integrated nutrient management for tomato-okra-stem amaranth cropping pattern in homestead area. *Journal of Food, Agriculture and Environment*, **9**, 438–445.
- Jenkinson DS, Johnston AE (1977) *Soil organic matter in the Hoosfield continuous barley experiment*. Rothamsted Report for 1976, Part 2. Rothamsted Experimental Station, Harpenden.
- Jenkinson DS, Andrew SPS, Lynch JM, Goss MJ, Tinker PB (1990) The turnover of organic carbon and nitrogen in soil. *Philosophical Transactions of the Royal Society of London, B*, **329**, 361–368.
- Jokela WE, Grabber JH, Karlen DL, Balser TC, Palmquist DE (2009) Cover crop and liquid manure effects on soil quality indicators in a corn silage system. *Agronomy Journal*, **101**, 727–737.
- Kamoni PT, Gicheru PT, Wokabi SM *et al.* (2007) Evaluation of two soil carbon models using two Kenyan long term experimental datasets. *Agriculture, Ecosystems & Environment*, **122**, 95–104.
- Kapkiyai JJ, Karanja NK, Qureshi JN, Smithson PC, Woormer PL (1999) Soil organic matter and nutrient dynamics in a Kenyan nitisol under long-term fertilizer and organic input management. *Soil Biology and Biochemistry*, **31**, 1773–1782.
- Kätterer T, Bolinder MA, André O, Kirchmann H, Menichetti L (2011) Roots contribute more to refractory soil organic matter than above-ground crop residues, as revealed by a long-term field experiment. *Agriculture, Ecosystems and Environment*, **141**, 184–192.
- Kibunja CN, Mwaura FB, Mugendi DN (2010) Long-term land management effects on soil properties and microbial populations in a maize-bean rotation at Kabete, Kenya. *African Journal of Agricultural Research*, **5**, 108–113.
- Kihanda FM, Warren GP, Micheni AN (2005) Effects of manure and fertilizer on grain yield, soil carbon and phosphorus in a 13-year field trial in semi-arid Kenya. *Experimental Agriculture*, **41**, 389–412.
- Kirchmann H, Haberer G, Kandeler E, Sessitsch A, Gerzabek MH (2004) Effects of level and quality of organic matter input on carbon storage and biological activity in soil: synthesis of a long-term experiment. *Global Biogeochemical Cycles*, **18**, GB4011.
- Kong AYY, Six J, Bryant DC, Denison RF, Van Kessel C (2005) The relationship between carbon input, aggregation, and soil organic carbon stabilization in sustainable cropping systems. *Soil Science Society of America Journal*, **69**, 1078–1085.
- Kukul SS, Rehana R, Benbi DK (2009) Soil organic carbon sequestration in relation to organic and inorganic fertilization in rice-wheat and maize-wheat systems. *Soil & Tillage Research*, **102**, 87–92.
- Lee C, Wu M, Asio VB, Chen Z (2006) Using a soil quality index to assess the effects of applying swine manure compost on soil quality under a crop rotation system in Taiwan. *Soil Science*, **171**, 210–222.
- Lee DK, Owens VN, Doolittle JJ (2007) Switchgrass and soil carbon sequestration response to ammonium nitrate, manure, and harvest frequency on conservation reserve program land. *Agronomy Journal*, **99**, 462–468.
- Leifeld J, Reiser R, Oberholzer HR (2009) Consequences of conventional versus organic farming on soil carbon: results from a 27-year field experiment. *Agronomy Journal*, **101**, 1204–1218.
- Liang B, Yang X, He X, Zhou J (2011) Effects of 17-year fertilization on soil microbial biomass C and N and soluble organic C and N in loessial soil during maize growth. *Biology and Fertility of Soils*, **47**, 121–128.
- Lou Y, Wang J, Liang W (2011) Impacts of 22-year organic and inorganic N managements on soil organic C fractions in a maize field, northeast China. *Catena*, **87**, 386–390.
- Ludwig B, Geisseler D, Michel K *et al.* (2011) Effects of fertilization and soil management on crop yields and carbon stabilization in soils A review. *Agronomy for Sustainable Development*, **31**, 361–372.
- Lynch DH, Voroney RP, Warman PR (2005) Soil physical properties and organic matter fractions under forages receiving composts, manure or fertilizer. *Compost Science & Utilization*, **13**, 252–261.
- Majumder B, Mandal B, Bandyopadhyay PK, Gangopadhyay A, Mani PK, Kundu AL, Mazumdar D (2008) Organic amendments influence soil organic carbon pools and rice-wheat productivity. *Soil Science Society of America Journal*, **72**, 775–785.
- Mandal B, Majumder B, Bandyopadhyay PK *et al.* (2007) The potential of cropping systems and soil amendments for carbon sequestration in soils under long-term experiments in subtropical India. *Global Change Biology*, **13**, 357–369.
- Mellek JE, Dieckow J, Da Silva VL, Favaretto N, Pauletti V, Vezzani FM, De Souza JLM (2010) Dairy liquid manure and no-tillage: Physical and hydraulic properties and carbon stocks in a Cambisol of Southern Brazil. *Soil and Tillage Research*, **110**, 69–76.
- Morari F, Lugato E, Berti A, Giardini L (2006) Long-term effects of recommended management practices on soil carbon changes and sequestration in north-eastern Italy. *Soil Use and Management*, **22**, 71–81.
- More SD, Hangarge DS (2003) Effect of integrated nutrient supply on crop productivity and soil characteristics with cotton-sorghum cropping sequence in Vertisol. *Journal of Maharashtra Agricultural Universities*, **28**, 8–12.
- Morlat R, Chaussod R (2008) Long-term Additions of Organic Amendments in a Loire Valley Vineyard. I. Effects on Properties of a Calcareous Sandy Soil. *American Journal of Enology and Viticulture*, **59**, 353–363.
- Nayak P, Patel D, Ramakrishnan B, Mishra AK, Samantaray RN (2009) Long-term application effects of chemical fertilizer and compost on soil carbon under intensive rice-rice cultivation. *Nutrient Cycling in Agroecosystems*, **83**, 259–269.
- Nayak AK, Gangwar B, Shukla AK *et al.* (2012) Long-term effect of different integrated nutrient management on soil organic carbon and its fractions and sustainability of rice-wheat system in Indo Gangetic Plains of India. *Field Crops Research*, **127**, 129–139.
- Ogle SM, Breidt FJ, Paustian K (2005) Agricultural management impacts on soil organic carbon storage under moist and dry climatic conditions of temperate and tropical regions. *Biogeochemistry*, **72**, 87–121.
- Parton WJ, Schimel DS, Cole CV, Ojima DS (1987) Analysis of Factors Controlling Soil Organic Matter Levels in Great Plains Grasslands. *Soil Science Society of America Journal*, **51**, 1173–1179.
- Rasmussen PE, Parton WJ (1994) Long-term effects of residue management in wheat-fallow: I. Inputs, yield, and soil organic matter. *Soil Science Society of America Journal*, **58**, 523–530.
- Rasool R, Kukul SS, Hira GS (2007) Soil physical fertility and crop performance as affected by long term application of FYM and inorganic fertilizers in rice-wheat system. *Soil & Tillage Research*, **96**, 64–72.
- Rasool R, Kukul SS, Hira GS (2008) Soil organic carbon and physical properties as affected by long-term application of FYM and inorganic fertilizers in maize-wheat system. *Soil & Tillage Research*, **101**, 31–36.
- Saha S, Mina BL, Gopinath KA, Kundu S, Gupta HS (2008) Organic amendments affect biochemical properties of a subtemperate soil of the Indian Himalayas. *Nutrient Cycling in Agroecosystems*, **80**, 233–242.
- Saha R, Mishra VK, Majumdar B, Laxminarayana K, Ghosh PK (2010a) Effect of integrated nutrient management on soil physical properties and crop productivity under a Maize (*Zea mays*)-mustard (*Brassica campestris*) cropping sequence in acidic soils of Northeast India. *Communications in Soil Science and Plant Analysis*, **41**, 2187–2200.
- Saha R, Nath V, Kumar D (2010b) Effects of farmyard manure on soil organic carbon stock, the pattern of fertility build-up, and plant growth in 'Mallika' mango (*Mangifera indica* L.). *Journal of Horticultural Science and Biotechnology*, **85**, 539–543.

- Sainju UM, Senwo ZN, Nyakatawa EZ, Tazisong IA, Reddy KC (2008a) Soil carbon and nitrogen sequestration as affected by long-term tillage, cropping systems, and nitrogen fertilizer sources. *Agriculture, Ecosystems & Environment*, **127**, 234–240.
- Sainju UM, Senwo ZN, Nyakatawa EZ, Tazisong IA, Reddy KC (2008b) Tillage, cropping systems, and nitrogen fertilizer source effects on soil carbon sequestration and fractions. *Journal of Environmental Quality*, **37**, 880–888.
- Sainju UM, Senwo ZN, Nyakatawa EZ, Tazisong IA, Reddy KC (2010) Poultry Litter Application Increases Nitrogen Cycling Compared with Inorganic Nitrogen Fertilization. *Agronomy Journal*, **102**, 917–925.
- Sarkar S, Singh SR, Singh RP (2003) The effect of organic and inorganic fertilizers on soil physical condition and the productivity of a rice-lentil cropping sequence in India. *Journal of Agricultural Science*, **140**, 419–425.
- Sharma KL, Grace JK, Mishra PK *et al.* (2011) Effect of soil and nutrient-management treatments on soil quality indices under cotton-based production system in rainfed semi-arid tropical vertisol. *Communications in Soil Science and Plant Analysis*, **42**, 1298–1315.
- Singh BR, Borresen T, Uhlen G, Ekeberg E (1998) Long term effects of crop rotation, cultivation practices, and fertilizers on carbon sequestration in soils of Norway. In: *Management of carbon sequestration in soil* (eds Lal R, Kimble JM, Follett RF, Stewart BA), pp. 195–208. CRC Press Inc, Boca Raton.
- Singh KN, Prasad B, Sinha SK (2001) Effect of integrated nutrient management on a Typic Haplaquent on yield and nutrient availability in a rice-wheat cropping system. *Australian Journal of Agricultural Research*, **52**, 855–858.
- Singh KP, Suman A, Singh PN, Srivastava TK (2007) Improving quality of sugarcane-growing soils by organic amendments under subtropical climatic conditions of India. *Biology and Fertility of Soils*, **44**, 367–376.
- Smith P, Powlson D, Glendining M, Smith JU (1997) Potential for carbon sequestration in European soils: preliminary estimates for five scenarios using results from long-term experiments. *Global Change Biology*, **3**, 67–79.
- Soltner D (2000) Les bases de la production végétale - Tome I: Le sol et son amélioration, Sciences et techniques agricoles.
- Srinivasarao C, Venkateswarlu B, Lal R *et al.* (2011) Long-term manuring and fertilizer effects on depletion of soil organic carbon stocks under pearl millet-cluster bean-castor rotation in Western India. *Land Degradation and Development*, doi: 10.1002/ldr.1158.
- Su Y-Z, Wang F, Suo D-R, Zhang Z-H, Du M-W (2006) Long-term effect of fertilizer and manure application on soil-carbon sequestration and soil fertility under the wheat-wheat-maize cropping system in northwest China. *Nutrient Cycling in Agroecosystems*, **75**, 285–295.
- Suman A, Singh KP, Singh P, Yadav RL (2009) Carbon input, loss and storage in subtropical Indian Inceptisol under multi-ratooning sugarcane. *Soil and Tillage Research*, **104**, 221–226.
- Thomsen IK, Christensen BT (2004) Yields of wheat and soil carbon and nitrogen contents following long-term incorporation of barley straw and ryegrass catch crops. *Soil Use and Management*, **20**, 432–438.
- Thomsen IK, Christensen BT (2010) Carbon sequestration in soils with annual inputs of maize biomass and maize-derived animal manure: evidence from ¹³C abundance. *Soil Biology & Biochemistry*, **42**, 1643–1646.
- Thomsen IK, Olesen JE, Møller HB, Sørensen P, Christensen BT (2013) Carbon dynamics and retention in soil after anaerobic digestion of dairy cattle feed and faeces. *Soil Biology and Biochemistry*, **58**, 82–87.
- Triberti L, Nastri A, Giordani G, Comellini F, Baldoni G, Toderi G (2008) Can mineral and organic fertilization help sequester carbon dioxide in cropland? *European Journal of Agronomy*, **29**, 13–20.
- Tummers B, Van Der Laan J (2006) Data Thief III - Version 1.6 - November 2010. Available at: <http://datathief.org/> (accessed 4 August 2011).
- Vandenbygaert AJ, Angers DA (2006) Towards accurate measurements of soil organic carbon stock change in agroecosystems. *Canadian Journal of Soil Science*, **86**, 465–471.
- Velthof GL, Bannink A, Oenema O, Van Der Meer HG, Spoelstra SF (2000) *Relationships between animal nutrition and manure quality - A literature review on C, N, P and S compounds*. Alterra-rapport 063, Alterra, Green World Research, Wageningen, 44 pp. Available at: <http://edepot.wur.nl/28901> (accessed 26 January 2010).
- Viaud V, Angers DA, Parnaudeau V, Morvan T, Aubry SM (2011) Response of organic matter to reduced tillage and animal manure in a temperate loamy soil. *Soil Use and Management*, **27**, 84–93.
- Walia MK, Walia SS, Dhaliwal SS (2010) Long-term effect of integrated nutrient management of properties of typic ustochrept after 23 cycles of an irrigated rice (*Oryza sativa* L.)-wheat (*Triticum aestivum* L.) system. *Journal of Sustainable Agriculture*, **34**, 724–743.
- Whalen JK, Chang C (2002) Macroaggregate Characteristics in Cultivated Soils after 25 Annual Manure Applications. *Soil Science Society of America Journal*, **66**, 1637–1647.
- Yang Z, Singh BR, Hansen S (2007) Aggregate associated carbon, nitrogen and sulfur and their ratios in long-term fertilized soils. *Soil & Tillage Research*, **95**, 161–171.
- Zhao B-Q, Li X-Y, Li X-P (2010) Long-term fertilizer experiment network in China: crop yields and soil nutrient trends. *Agronomy Journal*, **102**, 216–230.