

Three-Level Atom: Two Multi-Mode Filters Moment Equations

Jacob Ngaha

May 13, 2021

1 Hamiltonian and Master Equation

The Hamiltonian is

$$H = -\hbar \left(\frac{\alpha}{2} + \delta \right) |e\rangle \langle e| - 2\hbar\delta |f\rangle \langle f| + \hbar \frac{\Omega}{2} (\Sigma_+ + \Sigma_-) + \hbar \sum_{j=-N}^N \left(\omega_j^{(a)} a_j^\dagger a_j + \omega_j^{(b)} b_j^\dagger b_j \right) + \frac{i\hbar}{2} \sum_{j=-N}^N \left(\mathcal{E}_j^{(a)*} a_j \Sigma_+ - \mathcal{E}_j^{(a)} \Sigma_- a_j^\dagger \right) + \frac{i\hbar}{2} \sum_{j=-N}^N \left(\mathcal{E}_j^{(b)*} b_j \Sigma_+ - \mathcal{E}_j^{(b)} \Sigma_- b_j^\dagger \right) \quad (1)$$

where Ω is the Rabi frequency,

$$\alpha = \omega_{fe} - \omega_{eg}, \quad (2)$$

is the atomic anharmonicity, δ is given by

$$2\delta = 2\omega_d - \omega_{fg}, \quad (3)$$

is the *drive detuning from two-photon resonance*

$$\Sigma_- = \sigma_-^{eg} + \xi \sigma_-^{fe} = |g\rangle \langle e| + \xi |e\rangle \langle f|, \quad \Sigma_+ = \Sigma_-^\dagger, \quad (4)$$

is the atomic raising (lowering) operator, a^\dagger (b^\dagger) and a (b) are the cavity photon creation and annihilation operators for filter A (B), N is the number of modes either side of the central mode ($2N + 1$ total modes),

$$\omega_j^{(a)} = \omega_0^{(a)} + j\delta\omega^{(a)}, \quad \omega_j^{(b)} = \omega_0^{(b)} + j\delta\omega^{(b)} \quad (5)$$

is the resonance frequency of the j^{th} mode, with central frequency $\omega_0^{(a/b)}$ and mode frequency spacing $\delta\omega^{(a/b)}$, and

$$\mathcal{E}_j^{(a)} = \sqrt{\frac{\epsilon\gamma\kappa_a}{2(2N+1)}} e^{im\varphi_j}, \quad \mathcal{E}_j^{(b)} = \sqrt{\frac{\epsilon\gamma\kappa_b}{2(2N+1)}} e^{im\varphi_j}, \quad (6)$$

is the cascaded systems coupling of the j^{th} mode for filter A (B), where γ is the atomic decay rate, κ_a (κ_b) is the cavity decay rate for filter A (B), ϵ is the percentage of fluorescence sent to the filter, and

$$\varphi_j = \frac{j\pi}{N}, \quad (7)$$

sets the size of the frequency dependent time delay, with integer m .

The master equation is

$$\begin{aligned} \frac{d\rho}{dt} = \frac{1}{i\hbar} [H, \rho] + \frac{\gamma}{2} (1 - \epsilon) (2\sigma_- \rho \sigma_+ - \sigma_+ \sigma_- \rho - \rho \sigma_+ \sigma_-) \\ + \frac{\kappa_a}{2} \sum_{j=-N}^N \left(2a_j \rho a_j^\dagger - a_j^\dagger a_j \rho - \rho a_j^\dagger a_j \right) \end{aligned} \quad (8)$$

$$\begin{aligned}
& + \frac{1}{2} \sum_{j=-N}^N \left(2C_j^{(a)} \rho C_j^{(a)\dagger} - C_j^{(a)\dagger} C_j^{(a)} \rho - \rho C_j^{(a)\dagger} C_j^{(a)} \right) \\
& + \frac{\kappa_b}{2} \sum_{j=-N}^N \left(2b_j \rho b_j^\dagger - b_j^\dagger b_j \rho - \rho b_j^\dagger b_j \right) \\
& + \frac{1}{2} \sum_{j=-N}^N \left(2C_j^{(b)} \rho C_j^{(b)\dagger} - C_j^{(b)\dagger} C_j^{(b)} \rho - \rho C_j^{(b)\dagger} C_j^{(b)} \right),
\end{aligned} \tag{9}$$

where

$$C_j = \sqrt{\frac{\epsilon\gamma}{2N+1}} \sigma_- + \sqrt{\kappa} e^{im\varphi_j} a_j, \tag{10}$$

is the cascaded systems decay operator. Expanding the master equation out into neat terms, we arrive at:

$$\begin{aligned}
\frac{d\rho}{dt} = & i \left(\frac{\alpha}{2} + \delta \right) (\sigma^{ee} \rho - \rho \sigma^{ee}) + 2i\delta (\sigma^{ff} \rho - \rho \sigma^{ff}) - i \frac{\Omega}{2} (\Sigma_+ \rho - \rho \Sigma_+) \\
& - i \frac{\Omega}{2} (\Sigma_- \rho - \rho \Sigma_-) + \frac{\Gamma}{2} (2\Sigma_- \rho \Sigma_+ - \Sigma_+ \Sigma_- \rho - \rho \Sigma_+ \Sigma_-) \\
& - i \sum_{j=-N}^N \omega_j^{(a)} (a_j^\dagger a_j \rho - \rho a_j^\dagger a_j) + \kappa_a \sum_{j=-N}^N (2a_j \rho a_j^\dagger - a_j^\dagger a_j \rho - \rho a_j^\dagger a_j) \\
& - \sum_{j=-N}^N \mathcal{E}_j^{(a)} (a_j^\dagger \sigma_- \rho - \sigma_- \rho a_j^\dagger) - \sum_{j=-N}^N \mathcal{E}_j^{(a)*} (\rho \sigma_+ a_j - a_j \rho \sigma_+) \\
& - i \sum_{j=-N}^N \omega_j^{(b)} (b_j^\dagger b_j \rho - \rho b_j^\dagger b_j) + \kappa_b \sum_{j=-N}^N (2b_j \rho b_j^\dagger - b_j^\dagger b_j \rho - \rho b_j^\dagger b_j) \\
& - \sum_{j=-N}^N \mathcal{E}_j^{(b)} (b_j^\dagger \sigma_- \rho - \sigma_- \rho b_j^\dagger) - \sum_{j=-N}^N \mathcal{E}_j^{(b)*} (\rho \sigma_+ b_j - b_j \rho \sigma_+).
\end{aligned} \tag{11}$$

2 Operator Averages

2.1 First-Order: Atomic Equations

Rearranging the density operator equations, we can write the atomic moment equations in matrix form

$$\frac{d}{dt} \langle \sigma \rangle = \mathbf{M} \langle \sigma \rangle + \mathbf{B}, \tag{12}$$

where

$$\langle \sigma \rangle = \begin{pmatrix} \langle \sigma^{gg} \rangle \\ \langle \sigma_-^{eg} \rangle \\ \langle \sigma_+^{eg} \rangle \\ \langle \sigma^{ee} \rangle \\ \langle \sigma_-^{fe} \rangle \\ \langle \sigma_+^{fe} \rangle \\ \langle \sigma_-^{fg} \rangle \\ \langle \sigma_+^{fg} \rangle \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ \Gamma \xi^2 \\ i\xi \frac{\Omega}{2} \\ -i\xi \frac{\Omega}{2} \\ 0 \\ 0 \end{pmatrix}, \tag{13}$$

and

$$\mathbf{M} = \begin{pmatrix} 0 & -i\frac{\Omega}{2} & i\frac{\Omega}{2} & \Gamma & 0 & 0 & 0 & 0 \\ -i\frac{\Omega}{2} & -[\frac{\Gamma}{2} - i(\frac{\Omega}{2} + \delta)] & 0 & i\frac{\Omega}{2} & \Gamma\xi & 0 & -i\xi\frac{\Omega}{2} & 0 \\ i\frac{\Omega}{2} & 0 & -[\frac{\Gamma}{2} + i(\frac{\Omega}{2} + \delta)] & -i\frac{\Omega}{2} & 0 & \Gamma\xi & 0 & i\xi\frac{\Omega}{2} \\ -\Gamma\xi^2 & i\frac{\Omega}{2} & -i\frac{\Omega}{2} & -\Gamma(1+\xi^2) & -i\xi\frac{\Omega}{2} & i\xi\frac{\Omega}{2} & 0 & 0 \\ -i\xi\frac{\Omega}{2} & 0 & 0 & -i\xi\Omega & -[\frac{\Gamma}{2}(1+\xi^2) + i(\frac{\Omega}{2} - \delta)] & 0 & i\frac{\Omega}{2} & 0 \\ i\xi\frac{\Omega}{2} & 0 & 0 & i\xi\Omega & 0 & -[\frac{\Gamma}{2}(1+\xi^2) - i(\frac{\Omega}{2} - \delta)] & 0 & -i\frac{\Omega}{2} \\ 0 & -i\xi\frac{\Omega}{2} & 0 & 0 & i\frac{\Omega}{2} & 0 & -[\frac{\Gamma}{2}\xi^2 - 2i\delta] & 0 \\ 0 & 0 & i\xi\frac{\Omega}{2} & 0 & 0 & -i\frac{\Omega}{2} & 0 & -[\frac{\Gamma}{2}\xi^2 + 2i\delta] \end{pmatrix}. \quad (14)$$

This differential equation has solution

$$\langle \sigma(t) \rangle = e^{\mathbf{M}t} \langle \sigma(0) \rangle + (1 - e^{\mathbf{M}t}) \langle \sigma \rangle_{ss}, \quad (15)$$

where

$$\langle \sigma \rangle_{ss} = -\mathbf{M}^{-1} \mathbf{B}. \quad (16)$$

2.2 First-Order: Cavity Equations

$$\frac{d}{dt} \langle a_j \rangle = -(\kappa + i\omega_j) \langle a_j \rangle - \mathcal{E}_j \left(\langle \sigma_-^{eg} \rangle + \xi \langle \sigma_-^{fe} \rangle \right), \quad (17a)$$

$$\frac{d}{dt} \langle a_j^\dagger \rangle = -(\kappa - i\omega_j) \langle a_j^\dagger \rangle - \mathcal{E}_j \left(\langle \sigma_+^{eg} \rangle + \xi \langle \sigma_+^{fe} \rangle \right), \quad (17b)$$

$$\frac{d}{dt} \langle b_j \rangle = -(\kappa_b + i\omega_j^{(b)}) \langle b_j \rangle - \mathcal{E}_j^{(b)} \left(\langle \sigma_-^{eg} \rangle + \xi \langle \sigma_-^{fe} \rangle \right), \quad (17c)$$

$$\frac{d}{dt} \langle b_j^\dagger \rangle = -(\kappa_b - i\omega_j^{(b)}) \langle b_j^\dagger \rangle - \mathcal{E}_j^{(b)*} \left(\langle \sigma_+^{eg} \rangle + \xi \langle \sigma_+^{fe} \rangle \right). \quad (17d)$$

2.3 Second-Order: Cavity-Atom Coupled Equations

Using the vector notation, we have moment equations

$$\frac{d}{dt} \langle a_j \sigma \rangle = \mathbf{M}_j^{(a)} \langle a_j \sigma \rangle + \begin{pmatrix} -\mathcal{E}_j^{(a)} \langle \sigma_-^{eg} \rangle \\ -\mathcal{E}_j^{(a)} \xi \langle \sigma_-^{fg} \rangle \\ -\mathcal{E}_j^{(a)} \langle \sigma^{ee} \rangle \\ \Gamma\xi^2 \langle a_j \rangle - \mathcal{E}_j^{(a)} \xi \langle \sigma_-^{fe} \rangle \\ i\xi\frac{\Omega}{2} \langle a_j \rangle \\ -i\xi\frac{\Omega}{2} \langle a_j \rangle - \mathcal{E}_j^{(a)} \xi (1 - \langle \sigma^{gg} \rangle - \langle \sigma^{ee} \rangle) \\ 0 \\ -\mathcal{E}_j^{(a)} \langle \sigma_+^{fe} \rangle \end{pmatrix}, \quad (18a)$$

$$\frac{d}{dt} \langle b_j \sigma \rangle = \mathbf{M}_j^{(b)} \langle b_j \sigma \rangle + \begin{pmatrix} -\mathcal{E}_j^{(b)} \langle \sigma_-^{eg} \rangle \\ -\mathcal{E}_j^{(b)} \xi \langle \sigma_-^{fg} \rangle \\ -\mathcal{E}_j^{(b)} \langle \sigma^{ee} \rangle \\ \Gamma\xi^2 \langle b_j \rangle - \mathcal{E}_j^{(b)} \xi \langle \sigma_-^{fe} \rangle \\ i\xi\frac{\Omega}{2} \langle b_j \rangle \\ -i\xi\frac{\Omega}{2} \langle b_j \rangle - \mathcal{E}_j^{(b)} \xi (1 - \langle \sigma^{gg} \rangle - \langle \sigma^{ee} \rangle) \\ 0 \\ -\mathcal{E}_j^{(b)} \langle \sigma_+^{fe} \rangle \end{pmatrix}, \quad (18b)$$

where

$$\mathbf{M}_j^{(a)} = \mathbf{M} - \left(\kappa_a + i\omega_j^{(a)} \right) \mathbb{1}, \quad (19a)$$

$$\mathbf{M}_j^{(b)} = \mathbf{M} - \left(\kappa_b + i\omega_j^{(b)} \right) \mathbb{1}; \quad (19b)$$

and

$$\frac{d}{dt} \langle a_j^\dagger \boldsymbol{\sigma} \rangle = \mathbf{M}_j^{(a^\dagger)} \langle a_j^\dagger \boldsymbol{\sigma} \rangle + \begin{pmatrix} -\mathcal{E}_j^{(a)*} \langle \sigma_+^{eg} \rangle \\ -\mathcal{E}_j^{(a)*} \langle \sigma^{ee} \rangle \\ -\mathcal{E}_j^{(a)*} \xi \langle \sigma_+^{fg} \rangle \\ \Gamma \xi^2 \langle a_j^\dagger \rangle - \mathcal{E}_j^{(a)*} \xi \langle \sigma_+^{fe} \rangle \\ i\xi \frac{\Omega}{2} \langle a_j^\dagger \rangle - \mathcal{E}_j^{(a)*} \xi (1 - \langle \sigma^{gg} \rangle - \langle \sigma^{ee} \rangle) \\ -i\xi \frac{\Omega}{2} \langle a_j^\dagger \rangle \\ -\mathcal{E}_j^{(a)*} \langle \sigma_-^{fe} \rangle \\ 0 \end{pmatrix}, \quad (20a)$$

$$\frac{d}{dt} \langle b_j^\dagger \boldsymbol{\sigma} \rangle = \mathbf{M}_j^{(b^\dagger)} \langle b_j^\dagger \boldsymbol{\sigma} \rangle + \begin{pmatrix} -\mathcal{E}_j^{(b)*} \langle \sigma_+^{eg} \rangle \\ -\mathcal{E}_j^{(b)*} \langle \sigma^{ee} \rangle \\ -\mathcal{E}_j^{(b)*} \xi \langle \sigma_+^{fg} \rangle \\ \Gamma \xi^2 \langle b_j^\dagger \rangle - \mathcal{E}_j^{(b)*} \xi \langle \sigma_+^{fe} \rangle \\ i\xi \frac{\Omega}{2} \langle b_j^\dagger \rangle - \mathcal{E}_j^{(b)*} \xi (1 - \langle \sigma^{gg} \rangle - \langle \sigma^{ee} \rangle) \\ -i\xi \frac{\Omega}{2} \langle b_j^\dagger \rangle \\ -\mathcal{E}_j^{(b)*} \langle \sigma_-^{fe} \rangle \\ 0 \end{pmatrix}, \quad (20b)$$

$$(20c)$$

where

$$\mathbf{M}_j^{(a^\dagger)} = \mathbf{M} - \left(\kappa_a - i\omega_j^{(a)} \right) \mathbb{1}, \quad (21a)$$

$$\mathbf{M}_j^{(b^\dagger)} = \mathbf{M} - \left(\kappa_b - i\omega_j^{(b)} \right) \mathbb{1}. \quad (21b)$$

2.4 Second-Order: Cavity Equations

$$\begin{aligned} \frac{d}{dt} \langle a_j b_k \rangle &= - \left(\kappa_a + \kappa_b + i \left(\omega_j^{(a)} + \omega_k^{(b)} \right) \right) \langle a_j b_k \rangle \\ &\quad - \mathcal{E}_j^{(a)} \left(\langle b_k \sigma_-^{eg} \rangle + \xi \langle b_k \sigma_-^{fe} \rangle \right) - \mathcal{E}_k^{(b)} \left(\langle a_j \sigma_-^{eg} \rangle + \xi \langle a_j \sigma_-^{fe} \rangle \right), \end{aligned} \quad (22a)$$

$$\begin{aligned} \frac{d}{dt} \langle a_j^\dagger b_k^\dagger \rangle &= - \left(\kappa_a + \kappa_b - i \left(\omega_j^{(a)} + \omega_k^{(b)} \right) \right) \langle a_j^\dagger b_k^\dagger \rangle \\ &\quad - \mathcal{E}_j^{(a)*} \left(\langle b_k^\dagger \sigma_+^{eg} \rangle + \langle b_k^\dagger \sigma_+^{fe} \rangle \right) - \mathcal{E}_k^{(b)*} \left(\langle a_j^\dagger \sigma_+^{eg} \rangle + \langle a_j^\dagger \sigma_+^{fe} \rangle \right), \end{aligned} \quad (22b)$$

$$\begin{aligned} \frac{d}{dt} \langle a_j^\dagger a_k \rangle &= - \left(2\kappa_a - i \left(\omega_j^{(a)} - \omega_k^{(a)} \right) \right) \langle a_j^\dagger a_k \rangle \\ &\quad - \mathcal{E}_j^{(a)*} \left(\langle a_k \sigma_+^{eg} \rangle + \xi \langle a_k \sigma_+^{fe} \rangle \right) - \mathcal{E}_k^{(a)} \left(\langle a_j^\dagger \sigma_-^{eg} \rangle + \xi \langle a_j^\dagger \sigma_-^{fe} \rangle \right), \end{aligned} \quad (22c)$$

$$\frac{d}{dt} \langle b_j^\dagger b_k \rangle = - \left(2\kappa_b - i \left(\omega_j^{(b)} - \omega_k^{(b)} \right) \right) \langle b_j^\dagger b_k \rangle$$

$$-\mathcal{E}_j^{(b)*} \left(\langle b_k \sigma_+^{eg} \rangle + \xi \langle b_k \sigma_+^{fe} \rangle \right) - \mathcal{E}_k^{(b)} \left(\langle b_j^\dagger \sigma_-^{eg} \rangle + \xi \langle b_j^\dagger \sigma_-^{fe} \rangle \right), \quad (22d)$$

$$\begin{aligned} \frac{d}{dt} \langle a_j^\dagger b_k \rangle = & - \left(\kappa_a + \kappa_b - i \left(\omega_j^{(a)} - \omega_k^{(b)} \right) \right) \langle a_j^\dagger b_k \rangle \\ & - \mathcal{E}_j^{(a)*} \left(\langle b_k \sigma_+^{eg} \rangle + \xi \langle b_k \sigma_+^{fe} \rangle \right) - \mathcal{E}_k^{(b)} \left(\langle a_j^\dagger \sigma_-^{eg} \rangle + \xi \langle a_j^\dagger \sigma_-^{fe} \rangle \right), \end{aligned} \quad (22e)$$

$$\begin{aligned} \frac{d}{dt} \langle b_j^\dagger a_k \rangle = & - \left(\kappa_a + \kappa_b - i \left(\omega_j^{(b)} - \omega_k^{(a)} \right) \right) \langle b_j^\dagger a_k \rangle \\ & - \mathcal{E}_j^{(b)*} \left(\langle a_k \sigma_+^{eg} \rangle + \xi \langle a_k \sigma_+^{fe} \rangle \right) - \mathcal{E}_k^{(a)} \left(\langle b_j^\dagger \sigma_-^{eg} \rangle + \xi \langle b_j^\dagger \sigma_-^{fe} \rangle \right) \end{aligned} \quad (22f)$$

2.5 Third-Order: Cavity-Atom Coupled Equations

Using the vector notation, we have moment equations

$$\frac{d}{dt} \langle a_j b_k \boldsymbol{\sigma} \rangle = \mathbf{M}_{j,k}^{(ab)} \langle a_j b_k \boldsymbol{\sigma} \rangle + \begin{pmatrix} -\mathcal{E}_j^{(a)} \langle b_k \sigma_-^{eg} \rangle - \mathcal{E}_k^{(b)} \langle a_j \sigma_-^{eg} \rangle \\ -\mathcal{E}_j^{(a)} \xi \langle b_k \sigma_-^{fg} \rangle - \mathcal{E}_k^{(b)} \xi \langle a_j \sigma_-^{fg} \rangle \\ -\mathcal{E}_j^{(a)} \langle b_k \sigma_-^{ee} \rangle - \mathcal{E}_k^{(b)} \langle a_j \sigma_-^{ee} \rangle \\ \Gamma \xi \langle a_j b_k \rangle - \mathcal{E}_j^{(a)} \xi \langle b_k \sigma_-^{fe} \rangle - \mathcal{E}_k^{(b)} \xi \langle a_j \sigma_-^{fe} \rangle \\ i\xi \frac{\Omega}{2} \langle a_j b_k \rangle \\ -i\xi \frac{\Omega}{2} \langle a_j b_k \rangle - \mathcal{E}_j^{(a)} \xi (\langle b_k \rangle - \langle b_k \sigma^{gg} \rangle - \langle b_k \sigma^{ee} \rangle) - \mathcal{E}_k^{(b)} \xi (\langle a_j \rangle - \langle a_j \sigma^{gg} \rangle - \langle a_j \sigma^{ee} \rangle) \\ 0 \\ -\mathcal{E}_j^{(a)} \langle b_k \sigma_+^{fe} \rangle - \mathcal{E}_k^{(b)} \langle a_j \sigma_+^{fe} \rangle \end{pmatrix}, \quad (23a)$$

$$\frac{d}{dt} \langle a_j^\dagger b_k^\dagger \boldsymbol{\sigma} \rangle = \mathbf{M}_{j,k}^{(a^\dagger b^\dagger)} \langle a_j^\dagger b_k^\dagger \boldsymbol{\sigma} \rangle + \begin{pmatrix} -\mathcal{E}_j^{(a)*} \langle b_k^\dagger \sigma_+^{eg} \rangle - \mathcal{E}_k^{(b)*} \langle a_j^\dagger \sigma_+^{eg} \rangle \\ -\mathcal{E}_j^{(a)*} \langle b_k^\dagger \sigma_+^{ee} \rangle - \mathcal{E}_k^{(b)*} \langle a_j^\dagger \sigma_+^{ee} \rangle \\ -\mathcal{E}_j^{(a)*} \xi \langle b_k^\dagger \sigma_+^{fg} \rangle - \mathcal{E}_k^{(b)*} \xi \langle a_j^\dagger \sigma_+^{fg} \rangle \\ \Gamma \xi^2 \langle a_j^\dagger b_k^\dagger \rangle - \mathcal{E}_j^{(a)*} \xi \langle b_k^\dagger \sigma_+^{fe} \rangle - \mathcal{E}_k^{(b)*} \xi \langle a_j^\dagger \sigma_+^{fe} \rangle \\ i\xi \frac{\Omega}{2} \langle a_j^\dagger b_k^\dagger \rangle - \mathcal{E}_j^{(a)*} \xi (\langle b_k^\dagger \rangle - \langle b_k^\dagger \sigma^{gg} \rangle - \langle b_k^\dagger \sigma^{ee} \rangle) - \mathcal{E}_k^{(b)*} \xi (\langle a_j^\dagger \rangle - \langle a_j^\dagger \sigma^{gg} \rangle - \langle a_j^\dagger \sigma^{ee} \rangle) \\ -i\xi \frac{\Omega}{2} \langle a_j^\dagger b_k^\dagger \rangle \\ -\mathcal{E}_j^{(a)*} \langle b_k^\dagger \sigma_-^{fe} \rangle - \mathcal{E}_k^{(b)*} \langle a_j^\dagger \sigma_-^{fe} \rangle \\ 0 \end{pmatrix}, \quad (23b)$$

where

$$\mathbf{M}_{j,k}^{(ab)} = \mathbf{M} - \left(\kappa_a + \kappa_b + i \left(\omega_j^{(a)} + \omega_k^{(b)} \right) \right) \mathbb{1}, \quad (24a)$$

$$\mathbf{M}_{j,k}^{(a^\dagger b^\dagger)} = \mathbf{M} - \left(\kappa_a + \kappa_b - i \left(\omega_j^{(a)} + \omega_k^{(b)} \right) \right) \mathbb{1}; \quad (24b)$$

and

$$\frac{d}{dt} \langle a_j^\dagger a_k \boldsymbol{\sigma} \rangle = \mathbf{M}_{j,k}^{(a^\dagger a)} \langle a_j^\dagger a_k \boldsymbol{\sigma} \rangle + \begin{pmatrix} -\mathcal{E}_j^{(a)*} \langle a_k \sigma_+^{eg} \rangle - \mathcal{E}_k^{(a)} \langle a_j^\dagger \sigma_-^{eg} \rangle \\ -\mathcal{E}_j^{(a)*} \langle a_k \sigma_+^{ee} \rangle - \mathcal{E}_k^{(a)} \xi \langle a_j^\dagger \sigma_-^{fg} \rangle \\ -\mathcal{E}_j^{(a)*} \xi \langle a_k \sigma_+^{fg} \rangle - \mathcal{E}_k^{(a)} \langle a_j^\dagger \sigma_-^{ee} \rangle \\ \Gamma \xi^2 \langle a_j^\dagger a_k \rangle - \mathcal{E}_j^{(a)*} \xi \langle a_k \sigma_+^{fe} \rangle - \mathcal{E}_k^{(a)} \xi \langle a_j^\dagger \sigma_-^{fe} \rangle \\ i\xi \frac{\Omega}{2} \langle a_j^\dagger a_k \rangle - \mathcal{E}_j^{(a)*} \xi (\langle a_k \rangle - \langle a_k \sigma^{gg} \rangle - \langle a_k \sigma^{ee} \rangle) \\ -i\xi \frac{\Omega}{2} \langle a_j^\dagger a_k \rangle - \mathcal{E}_k^{(a)} \xi (\langle a_j^\dagger \rangle - \langle a_j^\dagger \sigma^{gg} \rangle - \langle a_j^\dagger \sigma^{ee} \rangle) \\ -\mathcal{E}_j^{(a)*} \langle a_k \sigma_-^{fe} \rangle \\ -\mathcal{E}_k^{(a)} \langle a_j^\dagger \sigma_+^{fe} \rangle \end{pmatrix}, \quad (25a)$$

$$\frac{d}{dt}\langle b_j^\dagger b_k \sigma \rangle = \mathbf{M}_{j,k}^{(b^\dagger b)} \langle b_j^\dagger b_k \sigma \rangle + \begin{pmatrix} -\mathcal{E}_j^{(b)*} \langle b_k \sigma_+^{eg} \rangle - \mathcal{E}_k^{(b)} \langle b_j^\dagger \sigma_-^{eg} \rangle \\ -\mathcal{E}_j^{(b)*} \langle b_k \sigma_+^{ee} \rangle - \mathcal{E}_k^{(b)} \xi \langle b_j^\dagger \sigma_-^{fg} \rangle \\ -\mathcal{E}_j^{(b)*} \xi \langle b_k \sigma_+^{fg} \rangle - \mathcal{E}_k^{(b)} \langle b_j^\dagger \sigma_-^{ee} \rangle \\ \Gamma \xi^2 \langle b_j^\dagger b_k \rangle - \mathcal{E}_j^{(b)*} \xi \langle b_k \sigma_+^{fe} \rangle - \mathcal{E}_k^{(b)} \xi \langle b_j^\dagger \sigma_-^{fe} \rangle \\ i\xi \frac{\Omega}{2} \langle b_j^\dagger b_k \rangle - \mathcal{E}_j^{(b)*} \xi (\langle b_k \rangle - \langle b_k \sigma^{gg} \rangle - \langle b_k \sigma^{ee} \rangle) \\ -i\xi \frac{\Omega}{2} \langle b_j^\dagger b_k \rangle - \mathcal{E}_k^{(b)} \xi (\langle b_j^\dagger \rangle - \langle b_j^\dagger \sigma^{gg} \rangle - \langle b_j^\dagger \sigma^{ee} \rangle) \\ -\mathcal{E}_j^{(b)*} \langle b_k \sigma_-^{fe} \rangle \\ -\mathcal{E}_k^{(b)} \langle b_j^\dagger \sigma_+^{fe} \rangle \end{pmatrix}, \quad (25b)$$

where

$$\mathbf{M}_{j,k}^{(a^\dagger a)} = \mathbf{M} - \left(2\kappa_a - i \left(\omega_j^{(a)} - i\omega_k^{(a)} \right) \right) \mathbb{1}, \quad (26a)$$

$$\mathbf{M}_{j,k}^{(b^\dagger b)} = \mathbf{M} - \left(2\kappa_b - i \left(\omega_j^{(b)} - i\omega_k^{(b)} \right) \right) \mathbb{1}; \quad (26b)$$

and

$$\frac{d}{dt}\langle a_j^\dagger b_k \sigma \rangle = \mathbf{M}_{j,k}^{(a^\dagger b)} \langle a_j^\dagger b_k \sigma \rangle + \begin{pmatrix} -\mathcal{E}_j^{(a)*} \langle b_k \sigma_+^{eg} \rangle - \mathcal{E}_k^{(b)} \langle a_j^\dagger \sigma_-^{eg} \rangle \\ -\mathcal{E}_j^{(a)*} \langle b_k \sigma_+^{ee} \rangle - \mathcal{E}_k^{(b)} \xi \langle a_j^\dagger \sigma_-^{fg} \rangle \\ -\mathcal{E}_j^{(a)*} \xi \langle b_k \sigma_+^{fg} \rangle - \mathcal{E}_k^{(b)} \langle a_j^\dagger \sigma_-^{ee} \rangle \\ \Gamma \xi^2 \langle a_j^\dagger b_k \rangle - \mathcal{E}_j^{(a)*} \xi \langle b_k \sigma_+^{fe} \rangle - \mathcal{E}_k^{(b)} \xi \langle a_j^\dagger \sigma_-^{fe} \rangle \\ i\xi \frac{\Omega}{2} \langle a_j^\dagger b_k \rangle - \mathcal{E}_j^{(a)*} \xi (\langle b_k \rangle - \langle b_k \sigma^{gg} \rangle - \langle b_k \sigma^{ee} \rangle) \\ -i\xi \frac{\Omega}{2} \langle a_j^\dagger b_k \rangle - \mathcal{E}_k^{(b)} \xi (\langle a_j^\dagger \rangle - \langle a_j^\dagger \sigma^{gg} \rangle - \langle a_j^\dagger \sigma^{ee} \rangle) \\ -\mathcal{E}_j^{(a)*} \langle b_k \sigma_-^{fe} \rangle \\ -\mathcal{E}_k^{(b)} \langle a_j^\dagger \sigma_+^{fe} \rangle \end{pmatrix}, \quad (27a)$$

$$\frac{d}{dt}\langle b_j^\dagger a_k \sigma \rangle = \mathbf{M}_{j,k}^{(b^\dagger a)} \langle b_j^\dagger a_k \sigma \rangle + \begin{pmatrix} -\mathcal{E}_j^{(b)*} \langle a_k \sigma_+^{eg} \rangle - \mathcal{E}_k^{(a)} \langle b_j^\dagger \sigma_-^{eg} \rangle \\ -\mathcal{E}_j^{(b)*} \langle a_k \sigma_+^{ee} \rangle - \mathcal{E}_k^{(a)} \xi \langle b_j^\dagger \sigma_-^{fg} \rangle \\ -\mathcal{E}_j^{(b)*} \xi \langle a_k \sigma_+^{fg} \rangle - \mathcal{E}_k^{(a)} \langle b_j^\dagger \sigma_-^{ee} \rangle \\ \Gamma \xi^2 \langle b_j^\dagger a_k \rangle - \mathcal{E}_j^{(b)*} \xi \langle a_k \sigma_+^{fe} \rangle - \mathcal{E}_k^{(a)} \xi \langle b_j^\dagger \sigma_-^{fe} \rangle \\ i\xi \frac{\Omega}{2} \langle b_j^\dagger a_k \rangle - \mathcal{E}_j^{(b)*} \xi (\langle a_k \rangle - \langle a_k \sigma^{gg} \rangle - \langle a_k \sigma^{ee} \rangle) \\ -i\xi \frac{\Omega}{2} \langle b_j^\dagger a_k \rangle - \mathcal{E}_k^{(a)} \xi (\langle b_j^\dagger \rangle - \langle b_j^\dagger \sigma^{gg} \rangle - \langle b_j^\dagger \sigma^{ee} \rangle) \\ -\mathcal{E}_j^{(b)*} \langle a_k \sigma_-^{fe} \rangle \\ -\mathcal{E}_k^{(a)} \langle b_j^\dagger \sigma_+^{fe} \rangle \end{pmatrix}, \quad (27b)$$

where

$$\mathbf{M}_{j,k}^{(a^\dagger b)} = \mathbf{M} - \left(\kappa_a + \kappa_b - i \left(\omega_j^{(a)} - i\omega_k^{(b)} \right) \right) \mathbb{1}, \quad (28a)$$

$$\mathbf{M}_{j,k}^{(b^\dagger a)} = \mathbf{M} - \left(\kappa_a + \kappa_b - i \left(\omega_j^{(b)} - i\omega_k^{(a)} \right) \right) \mathbb{1}; \quad (28b)$$

2.6 Third-Order: Cavity Equations

$$\begin{aligned} \frac{d}{dt}\langle a_j^\dagger a_k b_l \rangle &= - \left(2\kappa_a + \kappa_b - i \left(\omega_j^{(a)} - \omega_k^{(a)} - \omega_l^{(b)} \right) \right) \langle a_j^\dagger a_k b_l \rangle \\ &\quad - \mathcal{E}_j^{(a)*} \left(\langle a_k b_l \sigma_+^{eg} \rangle + \xi \langle a_k b_l \sigma_+^{fe} \rangle \right) \\ &\quad - \mathcal{E}_k^{(a)} \left(\langle a_j^\dagger b_l \sigma_-^{eg} \rangle + \xi \langle a_j^\dagger b_l \sigma_-^{fe} \rangle \right) \end{aligned}$$

$$-\mathcal{E}_l^{(b)} \left(\langle a_j^\dagger a_k \sigma_-^{eg} \rangle + \xi \langle a_j^\dagger a_k \sigma_-^{fe} \rangle \right), \quad (29a)$$

$$\begin{aligned} \frac{d}{dt} \langle b_j^\dagger b_k a_l \rangle = & - \left(\kappa_a + 2\kappa_b - i \left(\omega_j^{(b)} - \omega_k^{(b)} - \omega_l^{(a)} \right) \right) \langle b_j^\dagger b_k a_l \rangle \\ & - \mathcal{E}_j^{(b)*} \left(\langle b_k a_l \sigma_+^{eg} \rangle + \xi \langle b_k a_l \sigma_+^{fe} \rangle \right) \\ & - \mathcal{E}_k^{(b)} \left(\langle b_j^\dagger a_l \sigma_-^{eg} \rangle + \xi \langle b_j^\dagger a_l \sigma_-^{fe} \rangle \right) \\ & - \mathcal{E}_l^{(a)} \left(\langle b_j^\dagger b_k \sigma_-^{eg} \rangle + \xi \langle b_j^\dagger b_k \sigma_-^{fe} \rangle \right), \end{aligned} \quad (29b)$$

and

$$\begin{aligned} \frac{d}{dt} \langle b_j^\dagger a_k^\dagger a_l \rangle = & - \left(2\kappa_a + \kappa_b - i \left(\omega_j^{(b)} + \omega_k^{(a)} - \omega_l^{(a)} \right) \right) \langle b_j^\dagger a_k^\dagger a_l \rangle \\ & - \mathcal{E}_j^{(b)*} \left(\langle a_k^\dagger a_l \sigma_+^{eg} \rangle + \xi \langle a_k^\dagger a_l \sigma_+^{fe} \rangle \right) \\ & - \mathcal{E}_k^{(a)*} \left(\langle b_j^\dagger a_l \sigma_-^{eg} \rangle + \xi \langle b_j^\dagger a_l \sigma_-^{fe} \rangle \right) \\ & - \mathcal{E}_l^{(a)} \left(\langle b_j^\dagger a_k^\dagger \sigma_-^{eg} \rangle + \xi \langle b_j^\dagger a_k^\dagger \sigma_-^{fe} \rangle \right), \end{aligned} \quad (30a)$$

$$\begin{aligned} \frac{d}{dt} \langle a_j^\dagger b_k^\dagger b_l \rangle = & - \left(\kappa_a + 2\kappa_b - i \left(\omega_j^{(a)} + \omega_k^{(b)} - \omega_l^{(b)} \right) \right) \langle a_j^\dagger b_k^\dagger b_l \rangle \\ & - \mathcal{E}_j^{(a)*} \left(\langle b_k^\dagger b_l \sigma_+^{eg} \rangle + \xi \langle b_k^\dagger b_l \sigma_+^{fe} \rangle \right) \\ & - \mathcal{E}_k^{(b)*} \left(\langle a_j^\dagger b_l \sigma_-^{eg} \rangle + \xi \langle a_j^\dagger b_l \sigma_-^{fe} \rangle \right) \\ & - \mathcal{E}_l^{(b)} \left(\langle a_j^\dagger b_k^\dagger \sigma_-^{eg} \rangle + \xi \langle a_j^\dagger b_k^\dagger \sigma_-^{fe} \rangle \right), \end{aligned} \quad (30b)$$

2.7 Fourth-Order: Cavity-Atom Coupled Equations

Using the vector notation, we have moment equations

$$\begin{aligned} \frac{d}{dt} \langle a_j^\dagger a_k b_l \sigma \rangle = & \mathbf{M}_{j,k,l}^{(a^\dagger ab)} \langle a_j^\dagger a_k b_l \sigma \rangle + \\ & \left(\begin{aligned} & -\mathcal{E}_j^{(a)*} \langle a_k b_l \sigma_+^{eg} \rangle - \mathcal{E}_k^{(a)} \langle a_j^\dagger b_l \sigma_-^{eg} \rangle - \mathcal{E}_l^{(b)} \langle a_j^\dagger a_k \sigma_-^{eg} \rangle \\ & -\mathcal{E}_j^{(a)*} \langle a_k b_l \sigma_-^{ee} \rangle - \mathcal{E}_k^{(a)} \xi \langle a_j^\dagger b_l \sigma_-^{fg} \rangle - \mathcal{E}_l^{(b)} \xi \langle a_j^\dagger a_k \sigma_-^{fg} \rangle \\ & -\mathcal{E}_j^{(a)*} \xi \langle a_k b_l \sigma_+^{fg} \rangle - \mathcal{E}_k^{(a)} \langle a_j^\dagger b_l \sigma_-^{ee} \rangle - \mathcal{E}_l^{(b)} \langle a_j^\dagger a_k \sigma_-^{ee} \rangle \\ & \Gamma \xi^2 \langle a_j^\dagger a_k b_l \rangle - \mathcal{E}_j^{(a)*} \xi \langle a_k b_l \sigma_+^{fe} \rangle - \mathcal{E}_k^{(a)} \xi \langle a_j^\dagger b_l \sigma_-^{fe} \rangle - \mathcal{E}_l^{(b)} \xi \langle a_j^\dagger a_k \sigma_-^{fe} \rangle \\ & i \xi \frac{\Omega}{2} \langle a_j^\dagger a_k b_l \rangle - \mathcal{E}_j^{(a)*} \xi \left(\langle a_k b_l \rangle - \langle a_k b_l \sigma^{gg} \rangle - \langle a_k b_l \sigma^{ee} \rangle \right) \\ & -i \xi \frac{\Omega}{2} \langle a_j^\dagger a_k b_l \rangle - \mathcal{E}_k^{(a)} \xi \left(\langle a_j^\dagger a_k \rangle - \langle a_j^\dagger b_l \sigma^{gg} \rangle - \langle a_j^\dagger b_l \sigma^{ee} \rangle \right) - \mathcal{E}_l^{(b)} \xi \left(\langle a_j^\dagger b_l \rangle - \langle a_j^\dagger a_k \sigma^{gg} \rangle - \langle a_j^\dagger a_k \sigma^{ee} \rangle \right) \\ & -\mathcal{E}_j^{(a)*} \langle a_k b_l \sigma_-^{fe} \rangle \\ & -\mathcal{E}_k^{(a)} \langle a_j^\dagger b_l \sigma_+^{fe} \rangle - \mathcal{E}_l^{(b)} \langle a_j^\dagger a_k \sigma_-^{fe} \rangle \end{aligned} \right), \end{aligned} \quad (31a)$$

$$\frac{d}{dt} \langle b_j^\dagger b_k a_l \sigma \rangle = \mathbf{M}_{j,k,l}^{(b^\dagger ba)} \langle b_j^\dagger b_k a_l \sigma \rangle +$$

$$\left(\begin{array}{c} -\mathcal{E}_j^{(b)*} \langle b_k a_l \sigma_+^{eg} \rangle - \mathcal{E}_k^{(b)} \langle b_j^\dagger a_l \sigma_-^{eg} \rangle - \mathcal{E}_l^{(a)} \langle b_j^\dagger b_k \sigma_-^{eg} \rangle \\ -\mathcal{E}_j^{(b)*} \langle b_k a_l \sigma_-^{ee} \rangle - \mathcal{E}_k^{(b)} \xi \langle b_j^\dagger a_l \sigma_-^{fg} \rangle - \mathcal{E}_l^{(a)} \xi \langle b_j^\dagger b_k \sigma_-^{fg} \rangle \\ -\mathcal{E}_j^{(b)*} \xi \langle b_k a_l \sigma_+^{fg} \rangle - \mathcal{E}_k^{(b)} \langle b_j^\dagger a_l \sigma_-^{ee} \rangle - \mathcal{E}_l^{(a)} \langle b_j^\dagger b_k \sigma_-^{ee} \rangle \\ \Gamma \xi^2 \langle b_j^\dagger b_k a_l \rangle - \mathcal{E}_j^{(b)*} \xi \langle b_k a_l \sigma_+^{fe} \rangle - \mathcal{E}_k^{(b)} \xi \langle b_j^\dagger a_l \sigma_-^{fe} \rangle - \mathcal{E}_l^{(a)} \xi \langle b_j^\dagger b_k \sigma_-^{fe} \rangle \\ i \xi \frac{\Omega}{2} \langle b_j^\dagger b_k a_l \rangle - \mathcal{E}_j^{(b)*} \xi (\langle b_k a_l \rangle - \langle b_k a_l \sigma^{gg} \rangle - \langle b_k a_l \sigma^{ee} \rangle) \\ -i \xi \frac{\Omega}{2} \langle b_j^\dagger b_k a_l \rangle - \mathcal{E}_k^{(b)} \xi (\langle b_j^\dagger b_k \rangle - \langle b_j^\dagger a_l \sigma^{gg} \rangle - \langle b_j^\dagger a_l \sigma^{ee} \rangle) - \mathcal{E}_l^{(a)} \xi (\langle b_j^\dagger a_l \rangle - \langle b_j^\dagger b_k \sigma^{gg} \rangle - \langle b_j^\dagger b_k \sigma^{ee} \rangle) \\ -\mathcal{E}_j^{(b)*} \langle b_k a_l \sigma_-^{fe} \rangle \\ -\mathcal{E}_k^{(b)} \langle b_j^\dagger a_l \sigma_+^{fe} \rangle - \mathcal{E}_l^{(a)} \langle b_j^\dagger b_k \sigma_-^{fe} \rangle \end{array} \right), \quad (31b)$$

where

$$\mathbf{M}_{j,k,l}^{(a^\dagger ab)} = \mathbf{M} - \left(2\kappa_a + \kappa_b - i \left(\omega_j^{(a)} - \omega_k^{(a)} - \omega_l^{(b)} \right) \right) \mathbb{1}, \quad (32a)$$

$$\mathbf{M}_{j,k,l}^{(b^\dagger ba)} = \mathbf{M} - \left(\kappa_a + 2\kappa_b - i \left(\omega_j^{(b)} - \omega_k^{(b)} - \omega_l^{(a)} \right) \right) \mathbb{1}; \quad (32b)$$

and

$$\frac{d}{dt} \langle b_j^\dagger a_k^\dagger a_l \sigma \rangle = \mathbf{M}_{j,k,l}^{(b^\dagger a^\dagger a)} \langle b_j^\dagger a_k^\dagger a_l \sigma \rangle + \left(\begin{array}{c} -\mathcal{E}_j^{(b)*} \langle a_k^\dagger a_l \sigma_+^{eg} \rangle - \mathcal{E}_k^{(a)*} \langle b_j^\dagger a_l \sigma_+^{eg} \rangle - \mathcal{E}_l^{(a)} \langle b_j^\dagger a_k^\dagger \sigma_-^{eg} \rangle \\ -\mathcal{E}_j^{(b)*} \langle a_k^\dagger a_l \sigma_-^{ee} \rangle - \mathcal{E}_k^{(a)*} \langle b_j^\dagger a_l \sigma_-^{ee} \rangle - \mathcal{E}_l^{(a)} \xi \langle b_j^\dagger a_k^\dagger \sigma_-^{fg} \rangle \\ -\mathcal{E}_j^{(b)*} \xi \langle a_k^\dagger a_l \sigma_+^{fg} \rangle - \mathcal{E}_k^{(a)*} \xi \langle b_j^\dagger a_l \sigma_+^{fg} \rangle - \mathcal{E}_l^{(a)} \langle b_j^\dagger a_k^\dagger \sigma_-^{ee} \rangle \\ \Gamma \xi^2 \langle b_j^\dagger a_k^\dagger a_l \rangle - \mathcal{E}_j^{(b)*} \xi \langle a_k^\dagger a_l \sigma_+^{fe} \rangle - \mathcal{E}_k^{(a)*} \xi \langle b_j^\dagger a_l \sigma_-^{fe} \rangle - \mathcal{E}_l^{(a)} \xi \langle b_j^\dagger a_k^\dagger \sigma_-^{fe} \rangle \\ i \xi \frac{\Omega}{2} \langle b_j^\dagger a_k^\dagger a_l \rangle - \mathcal{E}_j^{(b)*} \xi (\langle a_k^\dagger a_l \rangle - \langle a_k^\dagger a_l \sigma^{gg} \rangle - \langle a_k^\dagger a_l \sigma^{ee} \rangle) - \mathcal{E}_k^{(a)*} \xi (\langle b_j^\dagger a_l \rangle - \langle b_j^\dagger a_l \sigma^{gg} \rangle - \langle b_j^\dagger a_l \sigma^{ee} \rangle) \\ -i \xi \frac{\Omega}{2} \langle b_j^\dagger a_k^\dagger a_l \rangle - \mathcal{E}_l^{(a)} \xi (\langle b_j^\dagger a_k^\dagger \rangle - \langle b_j^\dagger a_k^\dagger \sigma^{gg} \rangle - \langle b_j^\dagger a_k^\dagger \sigma^{ee} \rangle) \\ -\mathcal{E}_j^{(b)*} \langle a_k^\dagger a_l \sigma_-^{fe} \rangle - \mathcal{E}_k^{(a)*} \langle b_j^\dagger a_l \sigma_-^{fe} \rangle \\ -\mathcal{E}_l^{(a)} \langle b_j^\dagger a_k^\dagger \sigma_+^{fe} \rangle \end{array} \right), \quad (33a)$$

$$\frac{d}{dt} \langle a_j^\dagger b_k^\dagger b_l \sigma \rangle = \mathbf{M}_{j,k,l}^{(a^\dagger b^\dagger b)} \langle a_j^\dagger b_k^\dagger b_l \sigma \rangle + \left(\begin{array}{c} -\mathcal{E}_j^{(a)*} \langle b_k^\dagger b_l \sigma_+^{eg} \rangle - \mathcal{E}_k^{(b)*} \langle a_j^\dagger b_l \sigma_+^{eg} \rangle - \mathcal{E}_l^{(b)} \langle a_j^\dagger b_k^\dagger \sigma_-^{eg} \rangle \\ -\mathcal{E}_j^{(a)*} \langle b_k^\dagger b_l \sigma_-^{ee} \rangle - \mathcal{E}_k^{(b)*} \langle a_j^\dagger b_l \sigma_-^{ee} \rangle - \mathcal{E}_l^{(b)} \xi \langle a_j^\dagger b_k^\dagger \sigma_-^{fg} \rangle \\ -\mathcal{E}_j^{(a)*} \xi \langle b_k^\dagger b_l \sigma_+^{fg} \rangle - \mathcal{E}_k^{(b)*} \xi \langle a_j^\dagger b_l \sigma_+^{fg} \rangle - \mathcal{E}_l^{(b)} \langle a_j^\dagger b_k^\dagger \sigma_-^{ee} \rangle \\ \Gamma \xi^2 \langle a_j^\dagger b_k^\dagger b_l \rangle - \mathcal{E}_j^{(a)*} \xi \langle b_k^\dagger b_l \sigma_+^{fe} \rangle - \mathcal{E}_k^{(b)*} \xi \langle a_j^\dagger b_l \sigma_-^{fe} \rangle - \mathcal{E}_l^{(b)} \xi \langle a_j^\dagger b_k^\dagger \sigma_-^{fe} \rangle \\ i \xi \frac{\Omega}{2} \langle a_j^\dagger b_k^\dagger b_l \rangle - \mathcal{E}_j^{(a)*} \xi (\langle b_k^\dagger b_l \rangle - \langle b_k^\dagger b_l \sigma^{gg} \rangle - \langle b_k^\dagger b_l \sigma^{ee} \rangle) - \mathcal{E}_k^{(b)*} \xi (\langle a_j^\dagger b_l \rangle - \langle a_j^\dagger b_l \sigma^{gg} \rangle - \langle a_j^\dagger b_l \sigma^{ee} \rangle) \\ -i \xi \frac{\Omega}{2} \langle a_j^\dagger b_k^\dagger b_l \rangle - \mathcal{E}_l^{(b)} \xi (\langle a_j^\dagger b_k^\dagger \rangle - \langle a_j^\dagger b_k^\dagger \sigma^{gg} \rangle - \langle a_j^\dagger b_k^\dagger \sigma^{ee} \rangle) \\ -\mathcal{E}_j^{(a)*} \langle b_k^\dagger b_l \sigma_-^{fe} \rangle - \mathcal{E}_k^{(b)*} \langle a_j^\dagger b_l \sigma_-^{fe} \rangle \\ -\mathcal{E}_l^{(b)} \langle a_j^\dagger b_k^\dagger \sigma_+^{fe} \rangle \end{array} \right), \quad (33b)$$

where

$$\mathbf{M}_{j,k,l}^{(b^\dagger a^\dagger a)} = \mathbf{M} - \left(2\kappa_a + \kappa_b - i \left(\omega_j^{(b)} + \omega_k^{(a)} - \omega_l^{(a)} \right) \right) \mathbb{1}, \quad (34a)$$

$$\mathbf{M}_{j,k,l}^{(a^\dagger b^\dagger b)} = \mathbf{M} - \left(\kappa_a + 2\kappa_b - i \left(\omega_j^{(a)} + \omega_k^{(b)} - \omega_l^{(b)} \right) \right) \mathbb{1}. \quad (34b)$$

2.8 Fourth-Order: Cavity Equation

Finally, we have

$$\begin{aligned}
\frac{d}{dt} \langle a_j^\dagger b_k^\dagger b_l a_m \rangle = & - \left(2\kappa_a + 2\kappa_b - i \left(\omega_j^{(a)} + \omega_k^{(b)} \right) + i \left(\omega_l^{(b)} + \omega_m^{(a)} \right) \right) \langle a_j^\dagger b_k^\dagger b_l a_m \rangle \\
& - \mathcal{E}_j^{(a)*} \left(\langle b_k^\dagger b_l a_m \sigma_+^{eg} \rangle + \xi \langle b_k^\dagger b_l a_m \sigma_+^{fe} \rangle \right) \\
& - \mathcal{E}_k^{(b)*} \left(\langle a_j^\dagger a_l a_m \sigma_+^{eg} \rangle + \xi \langle a_j^\dagger a_l a_m \sigma_+^{fe} \rangle \right) \\
& - \mathcal{E}_l^{(b)} \left(\langle a_j^\dagger b_k^\dagger a_m \sigma_-^{eg} \rangle + \xi \langle a_j^\dagger b_k^\dagger a_m \sigma_-^{fe} \rangle \right) \\
& - \mathcal{E}_m^{(a)} \left(\langle a_j^\dagger b_k^\dagger b_l \sigma_-^{eg} \rangle + \xi \langle a_j^\dagger b_k^\dagger b_l \sigma_-^{fe} \rangle \right).
\end{aligned} \tag{35}$$

3 Second-Order Cross-Correlation Function

The second-order correlation function for the filtered output field in the steady state is given by

$$G_{\text{cross}}^{(2)}(\tau) = \langle A^\dagger(0) B^\dagger B(\tau) A(0) \rangle = \sum_{j,k=-N}^N \langle A^\dagger(0) b_j^\dagger b_k(\tau) A(0) \rangle, \tag{36}$$

with the normalised second-order correlation function given by

$$g_{\text{cross}}^{(2)}(\tau) = \frac{G_{\text{cross}}^{(2)}(\tau)}{\langle A^\dagger A \rangle_{ss} \langle B^\dagger B \rangle_{ss}}. \tag{37}$$

Using the Quantum Regression Theorem, we can solve for the second-order correlation function by solving the following moment equations

$$\frac{d}{d\tau} \langle A^\dagger(0) \boldsymbol{\sigma} A(0) \rangle = \mathbf{M} \langle A^\dagger(0) \boldsymbol{\sigma} A(0) \rangle + \mathbf{B}, \tag{38a}$$

$$\frac{d}{d\tau} \langle A^\dagger(0) b_j^\dagger(\tau) A(0) \rangle = - \left(\kappa_b - i\omega_j^{(b)} \right) \langle A^\dagger(0) b_j^\dagger(\tau) A(0) \rangle - \mathcal{E}_j^{(b)*} \left(\langle A^\dagger(0) \sigma_+^{eg}(\tau) A(0) \rangle + \xi \langle A^\dagger(0) \sigma_+^{fe} A(0) \rangle \right), \tag{38b}$$

$$\frac{d}{d\tau} \langle A^\dagger(0) b_j(\tau) A(0) \rangle = - \left(\kappa_b + i\omega_j^{(b)} \right) \langle A^\dagger(0) b_j(\tau) A(0) \rangle - \mathcal{E}_j^{(b)} \left(\langle A^\dagger(0) \sigma_-^{eg}(\tau) A(0) \rangle + \xi \langle A^\dagger(0) \sigma_-^{fe} A(0) \rangle \right), \tag{38c}$$

with

$$\langle \boldsymbol{\sigma} \rangle = \begin{pmatrix} \langle A^\dagger(0) \sigma^{gg}(\tau) A(0) \rangle \\ \langle A^\dagger(0) \sigma_-^{eg}(\tau) A(0) \rangle \\ \langle A^\dagger(0) \sigma_+^{eg}(\tau) A(0) \rangle \\ \langle A^\dagger(0) \sigma^{ee}(\tau) A(0) \rangle \\ \langle A^\dagger(0) \sigma_-^{fe}(\tau) A(0) \rangle \\ \langle A^\dagger(0) \sigma_+^{fe}(\tau) A(0) \rangle \\ \langle A^\dagger(0) \sigma_-^{fg}(\tau) A(0) \rangle \\ \langle A^\dagger(0) \sigma_+^{fg}(\tau) A(0) \rangle \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ \Gamma \xi^2 \langle A^\dagger A \rangle_{ss} \\ i\xi \frac{\Omega}{2} \langle A^\dagger A \rangle_{ss} \\ -i\xi \frac{\Omega}{2} \langle A^\dagger A \rangle_{ss} \\ 0 \\ 0 \end{pmatrix}. \tag{39}$$

We also need to solve

$$\frac{d}{d\tau} \langle A^\dagger(0) b_j \boldsymbol{\sigma}(\tau) A(0) \rangle = \mathbf{M}_j^{(b)} \langle A^\dagger(0) b_j \boldsymbol{\sigma}(\tau) A(0) \rangle$$

$$+ \begin{pmatrix} -\mathcal{E}_j^{(b)} \langle A^\dagger(0) \sigma_-^{eg}(\tau) A(0) \rangle \\ -\mathcal{E}_j^{(b)} \xi \langle A^\dagger(0) \sigma_-^{fg}(\tau) A(0) \rangle \\ -\mathcal{E}_j^{(b)} \langle A^\dagger(0) \sigma^{ee}(\tau) A(0) \rangle \\ \Gamma \xi^2 \langle A^\dagger(0) b_j(\tau) A(0) \rangle - \mathcal{E}_j^{(b)} \xi \langle A^\dagger(0) \sigma_-^{fe}(\tau) A(0) \rangle \\ i\xi \frac{\Omega}{2} \langle A^\dagger(0) b_j(\tau) A(0) \rangle \\ -i\xi \frac{\Omega}{2} \langle A^\dagger(0) b_j(\tau) A(0) \rangle - \mathcal{E}_j^{(b)} \xi (\langle A^\dagger A \rangle_{ss} - \langle A^\dagger(0) \sigma^{gg}(\tau) A(0) \rangle - \langle A^\dagger(0) \sigma^{ee}(\tau) A(0) \rangle) \\ 0 \\ -\mathcal{E}_j^{(b)} \langle A^\dagger(0) \sigma_+^{fe}(\tau) A(0) \rangle \end{pmatrix}, \quad (40a)$$

and

$$\frac{d}{d\tau} \langle A^\dagger(0) b_j^\dagger \sigma(\tau) A(0) \rangle = \mathbf{M}_j^{(b^\dagger)} \langle A^\dagger(0) b_j^\dagger \sigma(\tau) A(0) \rangle + \begin{pmatrix} -\mathcal{E}_j^{(b)*} \langle A^\dagger(0) \sigma_+^{eg}(\tau) A(0) \rangle \\ -\mathcal{E}_j^{(b)*} \langle A^\dagger(0) \sigma^{ee}(\tau) A(0) \rangle \\ -\mathcal{E}_j^{(b)*} \xi \langle A^\dagger(0) \sigma_+^{fg}(\tau) A(0) \rangle \\ \Gamma \xi^2 \langle A^\dagger(0) b_j^\dagger(\tau) A(0) \rangle - \mathcal{E}_j^{(b)*} \xi \langle A^\dagger(0) \sigma_+^{fe}(\tau) A(0) \rangle \\ i\xi \frac{\Omega}{2} \langle A^\dagger(0) b_j^\dagger(\tau) A(0) \rangle - \mathcal{E}_j^{(b)*} \xi (\langle A^\dagger A \rangle_{ss} - \langle A^\dagger(0) \sigma^{gg}(\tau) A(0) \rangle - \langle A^\dagger(0) \sigma^{ee}(\tau) A(0) \rangle) \\ -i\xi \frac{\Omega}{2} \langle A^\dagger(0) b_j^\dagger(\tau) A(0) \rangle \\ -\mathcal{E}_j^{(b)*} \langle A^\dagger(0) \sigma_-^{fe}(\tau) A(0) \rangle \\ 0 \end{pmatrix}, \quad (40b)$$

(40c)

where

$$\mathbf{M}_j^{(b)} = \mathbf{M} - (\kappa_b + i\omega_j^{(b)}) \mathbb{1}, \quad (41a)$$

$$\mathbf{M}_j^{(b^\dagger)} = \mathbf{M} - (\kappa_b - i\omega_j^{(b)}) \mathbb{1}. \quad (41b)$$

Finally, we will also need to solve

$$\begin{aligned} \frac{d}{d\tau} \langle A^\dagger(0) b_k^\dagger b_l(\tau) A(0) \rangle &= - \left(2\kappa_b - i\omega_k^{(b)} + i\omega_l^{(b)} \right) \langle A^\dagger(0) b_k^\dagger b_l(\tau) A(0) \rangle \\ &\quad - \mathcal{E}_k^{(b)*} \left(\langle A^\dagger(0) b_l \sigma_+^{eg}(\tau) A(0) \rangle + \xi \langle A^\dagger(0) b_l \sigma_+^{fe}(\tau) A(0) \rangle \right) \\ &\quad - \mathcal{E}_l^{(b)} \left(\langle A^\dagger(0) b_k^\dagger \sigma_-^{eg}(\tau) A(0) \rangle + \xi \langle A^\dagger(0) b_k^\dagger \sigma_-^{fe}(\tau) A(0) \rangle \right). \end{aligned} \quad (42)$$