

# Modelling the Milky Way Galaxy & the Large Magellanic Cloud

August, 2015

## 1 Useful Quantities and definitions

In this section some common quantities useful for describe the density profiles are defined and explained.

### 1.1 Critical density of the Universe:

The Critical density of the universe is defined as:

$$\rho_c = \frac{3H^2}{8\pi G} \quad (1)$$

Where  $H$  is the Hubble parameter and this parameter depends on the cosmological parameters. This density ...

### 1.2 Virialization

A dark matter halo is virialized when it is in equilibrium, such an equilibrium occurs after the dark matter have collapsed and the force of gravity equals the **relaxation** processes [1]. This happens when the dark matter reach an overdensity value  $\Delta_{vir}$ . This overdensity corresponds to a radius and a mass  $r_{vir}$  &  $M_{vir}$  respectively.

$\Delta_{vir} = \frac{\rho_{vir}}{\rho_c}$ . For a cosmology with  $(\Omega_m + \Omega_\Lambda = 1)$  the solution for the **Top Hat** model can be approximated by:

$$\Delta_{vir} = (18\pi^2 + 82x - 39x^2)/\Omega(z) \quad (2)$$

[3, 2] Where  $x = \Omega(z) - 1$ . For the present time ( $z = 0$ )  $\Delta_{vir} = 360$ .

The behavior of this function is shown in Fig.??




Figure 1: The solid blue line shows the behaviour of  $\Delta_{vir}$  as function of the redshift. The black line show the value of  $\Delta_{vir} = 173$  at  $z > 4$

The virial density now can be expressed in terms of  $\Delta_{vir}$ :

$$\rho_{vir} = \frac{3M_{vir}}{4\pi r_{vir}^3} = \Delta_{vir}\Omega_m\rho_{crit} \quad (3)$$

Where  $\Omega_m$  is the density parameter that give as the abundance of matter in the Universe, it is define as  $\Omega_m = \rho/\rho_c$  and the actual value is  $\Omega_m \simeq 0.3$  Once the virial density is defined with 3 for a given  $z$  then the radius and the virial mass can be related:

$$r_{vir} = \left( \frac{3M_{vir}}{4\pi\Delta_{vir}\Omega_m\rho_{crit}} \right)^{1/3} \quad (4)$$

For example for a halo of mass  $M = 1 \times 10^{12}M_\odot$  the corresponding radius is  $r_{vir} = 262.4$  Kpc

### 1.3 $r_{200}$ & $M_{200}$

There is another radius and mass of particular interest. This is the radius that enclosed a density of 200 times the density of the Universe.  $M_{200}$  is defined as:

$$M_{200} = 200\rho_c \frac{4}{3}\pi r_{200}^3 \quad (5)$$

In the same way  $M_{vir}$  is defined as:

$$M_{vir} = \Delta_{vir}\Omega_m\rho_c \frac{4}{3}\pi r_{vir}^3 \quad (6)$$

The critical density  $\rho_c$  is the same for both cases, then it is possible to relate both masses from Eq5 and Eq6 as follows:

$$\frac{M_{200}}{M_{vir}} = \left( \frac{200}{\Delta_{vir}\Omega_m} \right) \left( \frac{r_{200}}{r_{vir}} \right)^3 \quad (7)$$

Here is common to call  $q = \left( \frac{200}{\Delta_{vir}\Omega_m} \right)$ , at  $z = 0$   $q = 2.053$ .

$$\frac{M_{200}}{M_{vir}} = q \left( \frac{r_{200}}{r_{vir}} \right)^3 \quad (8)$$

This Eq.8 relates  $M_{vir}$  and  $M_{200}$  for a given  $r_{vir}$  and  $r_{200}$ .


## 2 Densities profiles

### 2.1 Plummer

The plumer density profile is one of the simplest models which describes a constant density near the center and falls at large radius.


$$\Phi_P(r) = -\frac{GM}{\sqrt{r^2 + a^2}} \quad (9)$$

Where  $a$  is call the scale length. The scale length set the length  $a$  in which the majority of the density is enclosed. Note that if  $a$  is zero the plummer potential would be exactly as the potential of a point mass. In the other hand if  $a$  goes to infty the potential is rewpresenting a very extended mass source. In other words the scale length set up the size of the volume in which the mass  $M$  is enclosed.



$$\nabla^2 \Phi_P(r) = 4\pi G \rho_P(r) = \frac{1}{r^2} \frac{d}{dr} \left( r^2 \frac{d\Phi_P(r)}{dr} \right) \quad (10)$$


$$\rho_P(r) = \frac{3M}{4\pi a^3} \left( 1 + \frac{r^2}{a^2} \right)^{-5/2} \quad (11)$$



The enclosed mass can be derived from the density by integrating over the volume.

$$M_P(< r) = 4\pi \int_0^r r'^2 \frac{3M}{4\pi a^3} \left(1 + \frac{r'^2}{a^2}\right)^{-5/2} dr' = \frac{3M}{a^3} \left( \frac{a^4 r^3 \sqrt{r^2/a^2 + 1}}{3(r^2 + a^2)^2} \right) \quad (12)$$


$$M_P(< r) = M \frac{r^3}{(a^2 + r^2)^{3/2}} \quad (13)$$



$$F_g = \frac{GmM}{r^2} = ma_c = m \frac{v_c^2}{r} \quad (14)$$

$$v_c = \sqrt{\frac{GM(< r)}{r}} \quad (15)$$

$$v_c = \sqrt{GM \left( \frac{r^2}{(r^2 + a^2)^{3/2}} \right)} \quad (16)$$



## 2.2 Hernquist profile

The Hernquist profile is derived in such a way that it follows the


$$\rho_{Hernquist}(r) = \frac{M}{2\pi} \frac{a}{r(r+a)^3} \quad (17)$$




$$M_{Hernquist}(< r) = 2aM \int \frac{r}{(r+a)^3} dr \quad (18)$$

$$M_{Hernquist}(< r) = M \frac{r^2}{(r+a)^2} \quad (19)$$

$$\Phi = -\frac{GM}{r+a} \quad (20)$$

$$v_c(r) = GM \frac{r}{(r+a)^2} \quad (21)$$





## 2.3 Singular Isothermal Sphere

The Singular Isothermal Sphere (SIS) describes a system in which the particles follow a Maxwellian density distribution. With this distribution and the Poisson equation the follow density profiles could be derived.

$$\rho_{iso}(r) = \frac{\sigma^2}{2\pi G r^2} \quad (22)$$


Following the same procedure as with the previous profiles we find  $M$ ,  $\Phi$  and  $v_c$ .



$$M_{iso}(< r) = \frac{2\sigma r}{G} \quad (23)$$

$$\Phi_{iso}(r) = 2\sigma^2 \ln(r) + const. \quad (24)$$

$$v_c(r) = \sqrt{2}\sigma \quad (25)$$

This profile is quite different to the previous ones due to the fact that here the input is the velocity instead of the total Mass.





## 2.4 NFW

$$\rho_{NFW}(r) = \frac{M}{2\pi a^3 (r/a)(1+r/a)^2} \quad (26)$$

$$M_{NFW}(r) = M \left( \ln(1+x) - \frac{x}{1+x} \right) \quad (27)$$

Where  $x = r/a$ , is useful to define the function  $f(x)$  as:

$$f(x) = \ln(1+x) - \frac{x}{1+x} \quad (28)$$

Then 27 can be expressed as:



$$M_{NFW} = 4\pi\rho_a a^3 f(x) \quad (29)$$




$$\Phi_{NFW} = -4\pi GM \frac{\ln(1+r/a)}{r} \quad (30)$$

$$c(M_{vir}) = 9.60 \left( \frac{M_{vir}}{10^{12} h^{-1} M_{\odot}} \right)^{-0.075} \quad (31)$$

$$v_c(r) = \sqrt{\left( \frac{M(r)G}{r} \right)} = \sqrt{\left( \frac{2M (\ln(1+c) - \frac{c}{1+c})}{r} \right)} \quad (32)$$





### 3 Conversion from NFW to the Hernquist profile

The average density of the NFW distribution can be expressed as:

$$\bar{\rho}_{NFW}(r) = \frac{3M_{NFW}(r)}{4\pi r^3} \quad (33)$$

And with eq.29 the  $\bar{\rho}_{NFW}(r)$  takes de form:


$$\bar{\rho}_{NFW}(r) = 3\rho_a \left(\frac{a}{r}\right)^3 f(x) \quad (34)$$

Now if we want to find the relationship between  $r_{200}$  and  $r_{vir}$  for the NFW profile we have to apply eq8.

$$q = \frac{3\rho_a \frac{a}{r_{200}} f(c_{200})}{3\rho_a \frac{a}{r_{vir}} f(c)} = \frac{c_{200}^3 f(c_{200})}{c_{vir}^3 f(c_{vir})} \quad (35)$$

$$?? \frac{c_{200}}{c_{vir}} = \left( \frac{f(c_{200})}{q f(c_{vir})} \right)^{1/3} \quad (36)$$

For  $c_{vir} = 10$  this function is shown in Fig.??, where  $y = \frac{c_{200}}{c_{vir}} - \left( \frac{f(c_{200})}{q f(c_{vir})} \right)^{1/3}$



Note that the solution of Eq.?? is when  $y = 0$ , one solution is  $c_{200} = 0$  but this is not of particular interest for us.

The other solution is computed analytically using the bisection algorithm.  $c_{200} = 7.4$

In order to seek the equivalence between the NFW and the Hernquist profile, We have to match the same enclosed mass of both profiles at a given radius. To his end we have to find  $M_H$  in terms of  $r_s$ .

$$M_H(r) = M_{NFW}(r) \quad (37)$$

$$\frac{M_H r^2}{a^2 + r^2} = 4\pi\rho_s r_s^3 \left[ \ln(1+x) - \frac{x}{1+x} \right] \quad (38)$$

In the limit  $r \rightarrow 0$

$$M_H = \frac{4\pi\rho_s r_s^3 a^2}{r^2} \left[ \left( x - \frac{x^2}{2} \right) - x \right] \quad (39)$$

$$M_H = 4\pi\rho_s r_s^3 \frac{a^2}{r^2} \left( -\frac{r^2}{2r_s^2} \right) \quad (40)$$

$$M_H = 2\pi r_s a^2 \quad (41)$$

With this relation is possible now to match both profiles at a given radius  $\tilde{r}$

$$M_H(\tilde{r}) = M_{NFW}(\tilde{r}) \quad (42)$$

$$2\pi\rho_s a^2 r_s \frac{\tilde{r}^2}{a^2} \frac{1}{\left(1 + \frac{\tilde{r}}{a}\right)^2} = 4\pi\rho_s r_s^3 \left( \text{Ln} \left(1 + \frac{\tilde{r}}{r_s}\right) - \frac{\tilde{r}}{\tilde{r} + r_s} \right) \quad (43)$$

$$\frac{\tilde{r}^2 a^2}{(a + \tilde{r})^2} = 2r_s^2 \left( \text{Ln} \left(1 + \frac{\tilde{r}}{r_s}\right) - \frac{\tilde{r}}{\tilde{r} + r_s} \right) \quad (44)$$

$$\frac{\tilde{r}^2 a^2}{(a + \tilde{r})^2} = 2r_s^2 f(\tilde{x}) \quad (45)$$

$$\left(\frac{a}{r_s}\right)^2 = \frac{2}{\tilde{r}^2} (a + \tilde{r})^2 f(\tilde{x}) \quad (46)$$

$$\frac{a}{r_s} = \frac{[2f(\tilde{x})]^{1/2}}{\tilde{r}} (a + \tilde{r}) \quad (47)$$

$$\left(\frac{a}{r_s}\right) \left(1 - \frac{[2f(\tilde{x})]^{1/2}}{\tilde{x}}\right) = [2f(\tilde{x})]^{1/2} \quad (48)$$

$$\frac{a}{r_s} = \frac{[2f(\tilde{x})]^{1/2}}{\left(1 - \frac{[2f(\tilde{x})]^{1/2}}{\tilde{x}}\right)} \quad (49)$$

$$\frac{a}{r_s} = \frac{[2f(\tilde{x})]^{1/2} \tilde{x}}{\tilde{x} - (2f(\tilde{x})^{1/2})} = \frac{1}{\left([2f(\tilde{x})]^{-1/2} - \frac{1}{\tilde{x}}\right)} \quad (50)$$


Finally the ratio of the enclosed mass of the Hernquist and the NFW profiles is:

$$\frac{M_H}{M_{vir}} = \frac{2\pi\rho_s a^2 r_s}{4\pi\rho_s r_s^3 f(c_{vir})} = \frac{1}{2f(c_{vir})} \left(\frac{a}{r_s}\right)^2 \quad (51)$$

## 4 Miyamoto-Nagai Disk

$$\Phi_M(R, z) = -\frac{GM}{\sqrt{R^2 + (a + \sqrt{(z^2 + b^2)})^2}} \quad (52)$$

$$\rho_M(R, Z) = \left(\frac{b^2 M}{4\pi}\right) \frac{aR^2 + (a + 3\sqrt{z^2 + b^2})(a + \sqrt{z^2 + b^2})^2}{[R^2 + (a^2 + \sqrt{z^2 + b^2})^2]^{5/2} (z^2 + b^2)^{3/2}} \quad (53)$$




## 5 Logarithmic Profile

Disc profile

$$\Phi_L(R, z) = \frac{1}{2} v_0^2 \ln \left( R_c^2 + R^2 + \frac{z^2}{q_\phi^2} \right) + \text{constant} \quad (54)$$

The circular velocity at  $z = 0$  is  $v_c^2(R, z = 0) = r \frac{d\Phi}{dR}$ :

$$v_c(R, z = 0) = r \frac{d\Phi_L}{dr} = \frac{v_0 R}{\sqrt{r_c^2 + R^2}} \quad (55)$$




To derive the density we make use of Poisson's equation in cylindrical coordinates:

$$\rho_L(R, z) = \frac{\nabla^2 \Phi_L}{4\pi G} = \frac{1}{4\pi G} \left( \frac{1}{r} \frac{d}{dR} R \frac{d}{dR} + \frac{d^2}{dz^2} \right) \Phi_L \quad (56)$$

$$\rho_L(R, z) = \frac{v_0^2}{8\pi G} \left( \frac{1}{R} \frac{4R(R_c^2 + R^2 + \frac{z^2}{q_\phi^2}) - 4R^3}{(R_c^2 + R^2 + \frac{z^2}{q_\phi^2})^2} + \frac{\frac{2}{q_\phi^2}(R_c^2 + R^2 + \frac{z^2}{q_\phi^2}) - \frac{4z^2}{q_\phi^4}}{(R_c^2 + R^2 + \frac{z^2}{q_\phi^2})^2} \right) \quad (57)$$

$$\rho_L(R, z) = \frac{v_0^2}{4\pi G q_\phi^2} \frac{(2q_\phi^2 + 1)R_c^2 + r^2 + (2 - q_\phi^2)z^2}{(R_c^2 + r^2 + z^2 q_\phi^{-2})^2} \quad (58)$$



## 6 Logarithmic profile (Law, Johnston & Majewski)

This profile is almost the same that the one explained in the previous section 5

$$\Phi = v_{halo}^2 \ln[r^2 + (z^2/q^2) + d^2] \quad (59)$$

$$v_c = \frac{\sqrt{2}vr}{\sqrt{r^2 + d^2}} \quad (60)$$

with  $q = 0.8 - 1.45$ ,  $v_{halo}$  should match  $v_{circ,\odot}$ ,  $d = 1 - 20kpc$

## 7 Logarithmic (Law & Majewski 2010)

[http://iopscience.iop.org/0004-637X/714/1/229/pdf/apj\\_714\\_1\\_229.pdf](http://iopscience.iop.org/0004-637X/714/1/229/pdf/apj_714_1_229.pdf)

$$\Phi = v_{halo}^2 \ln(C_1 x^2 + C_2 y^2 + C_3 xy + (z/q_z)^2 r_{halo}^2) \quad (61)$$

$R, r$  (Cylindrical, spherical)


$$C_1 = \left( \frac{\cos^2 \phi}{q_1^2} + \frac{\sin^2 \phi}{q_2^2} \right) \quad (62)$$

$$C_2 = \left( \frac{\cos^2 \phi}{q_2^2} + \frac{\sin^2 \phi}{q_1^2} \right) \quad (63)$$

$$C_3 = 2 \sin \phi \cos \phi \left( \frac{1}{q_1^2} - \frac{1}{q_2^2} \right) \quad (64)$$

## 8 Modelling the MW

Component	Besla07	LM2010	Roeland12	Gomez15
Disk Model	Miyamoto-Nagai	Miyamoto-Nagai		
Disk Mass( $M_\odot$ )	$5.5^{10}$	$1.0 \times 10^{11}$		
Disk Param	$R_d = 3.5, z = r_{disk}/5.0$	$\alpha = 1$		$, a = 6.5kpc, b = 0.26Kpc$
Bulge Model	Hernquist	Hernquist		Hernquist
Bulge Mass( $M_\odot$ )	$10^{10}$	$3.4 \times 10^{10}$		
Bulge Param	$0.6kpc$	$3.4 \times 10^{10} M_\odot, c = 0.7kpc$	$0.6Kpc$	
DM halo Model	NFW	7	Hernquist(NFW)	
DM halo mass( $M_\odot$ )	$10^{12}$	$\times 10^{10}$		
Halo Param	$c = 11, r_{vir} = 258Kpc$	$r_{halo} = 12Kpc$		
Solar distance $R_\odot$ (kpc)	8.0			
reference	Besla07	LM2010		



## 9 TO-DO:

- Study the properties and different parameters ( $M$ ,  $\rho$ ,  $v_c$ ,  $a$ )
- Plummer
- Henrquist
- Isothermal
- NFW
- Miyamoto-Nagai (Lunes)
- Exponential (Martes)
- Jhonston and Bullock 2005 (Logarithmic) (Martes)
- Besla 2007 (Miercoles)
- Law & Majewusky 2010 triaxial (Miercoles)
- Try to reproduce MW rotation curve van Der Marel 2012 ( $M_{vir}$  &  $c$ ) (Jueves)
- note: Klypin relation between  $c$  and  $M_{vir}$  Doesn't take into account adiabatic contraction.
- work in the code that integrates the orbits using the accelerations. (Viernes)

## References

1. J. Binney and S. Tremaine. *Galactic Dynamics: Second Edition*. Princeton University Press, 2008.



2. G. L. Bryan and M. L. Norman. Statistical Properties of X-Ray Clusters: Analytic and Numerical Comparisons. , 495:80–99, March 1998.
3. V. R. Eke, S. Cole, and C. S. Frenk. Cluster evolution as a diagnostic for Omega. , 282:263–280, September 1996.