

## THE IMPACT OF GAS BULK ROTATION ON THE LYMAN- $\alpha$ LINE

JUAN N. GARAVITO-CAMARGO<sup>1</sup>, JAIME E. FORERO-ROMERO<sup>1</sup>, MARK DIJKSTRA<sup>2</sup>

*Submitted for publication in ApJ*

### ABSTRACT

We present results of radiative transfer calculations to measure the impact of gas bulk rotation on the morphology of the Lyman  $\alpha$  line in galaxies. We model a galaxy as a sphere with a homogeneous mixture of dust and hydrogen at a constant temperature. These spheres have a solid-body rotation with maximum velocities in the range  $0 - 300 \text{ km s}^{-1}$  and neutral hydrogen optical depths in the range  $\tau_{\text{H}} = 10^5 - 10^7$ . We also consider two kinds of spatial distribution for the radiation sources with respect to the sphere: central and homogeneous. We find that the line width and the intensity at the line's centre increases with rotational velocity. In the case of homogeneously distributed sources, for large rotational velocities the line transforms from a double peak to a single peak at the line center. Under the same conditions the escape fraction increases  $\sim 30\%$ . For radiation sources located off-centre, the line morphology presents a range of single, double and triple peaked lines. We show how these results are useful to interpret recent spectroscopic results of distant  $z \sim 2 - 3$  star forming galaxies.

*Subject headings:* galaxies: high-redshift, galaxies: star formation, line: formation

### 1. INTRODUCTION

The detection of strong Ly $\alpha$  emission lines has become an essential method in extragalactic astronomy to find distant star-forming galaxies (Partridge & Peebles 1967; Rhoads et al. 2000; Gawiser et al. 2007; Koehler et al. 2007; Ouchi et al. 2008; Yamada et al. 2012; Schenker et al. 2012). The galaxies detected using this method receive the name of Ly $\alpha$  emitters (LAEs). A detailed examination of this galaxy population has diverse implications for galaxy formation, reionization and the large scale structure of the Universe. Attempts to fully exploit the physical information included in the Ly $\alpha$  line require an understanding of all the physical factors involved in shaping the line. Due to the resonant nature of this line, these physical factors notably include temperature, density and bulk velocity field of the neutral Hydrogen in the emitting galaxy and its surroundings.

A basic understanding of the quantitative behaviour of the Ly $\alpha$  line has been reached through analytical studies in the case of a static configurations, such as uniform slabs (Harrington 1973; Neufeld 1990) and uniform spheres (Dijkstra et al. 2006). Analytical studies of configurations including some kind of bulk flow only include the case of a sphere with a Hubble like expansion flow (Loeb & Rybicki 1999).

A quantitative description of the Ly $\alpha$  line has been reached through Monte Carlo simulations (Auer 1968; Avery & House 1968; Adams 1972). In the last two decades these studies have become popular due to the availability of computing power. Early into the 21st century the first studies focused on homogeneous and static media (Ahn et al. 2000, 2001; Zheng & Miralda-Escudé 2002). Later on the effects of clumpy media (Hansen & Oh 2006) and expanding/contracting shell/spherical geometries started to be studied (Verhamme et al. 2006;

Dijkstra et al. 2006). Similar codes have applied these results to semi-analytical models of galaxy formation (Orsi et al. 2012) and results of large hydrodynamical simulations (Forero-Romero et al. 2011, 2012; Behrens & Niemeyer 2013). Recently Monte Carlo codes have also been applied to the results of high resolution hydrodynamical simulations of individual galaxies (Laursen et al. 2009; Barnes et al. 2011; Verhamme et al. 2012; Yajima et al. 2012). While, recent developments have been focused on the systematic study of clumpy outflows (Dijkstra & Kramer 2012) and anisotropic velocity configurations (Zheng & Wallace 2013).

The recent studies of galaxies in hydrodynamical simulations (Laursen et al. 2009; Barnes et al. 2011; Verhamme et al. 2012; Yajima et al. 2012) have all shown systematic variations in the Ly $\alpha$  line with the viewing angle. These variations are a complex superposition of anisotropic density configurations (i.e. edge-on vs. face-on view of a galaxy), the inflows observed by gas cooling and the outflows included in the supernova feedback process of the simulation. These bulk flows physically correspond to the circumgalactic and intergalactic medium (CGM and IGM). These effects are starting to be studied in simplified configurations that vary the density and wind characteristics (Zheng & Wallace 2013).

However, in all these efforts the effect of rotation, which is an ubiquitous feature in galaxies, has not been systematically studied. The processing of the Ly $\alpha$  photons in a rotating interstellar medium (ISM) must have some kind of impact in the Ly $\alpha$  line morphology.

Performing that study is the main goal of this paper. We investigate for the first time the impact of rotation on the morphology of the Lyman  $\alpha$  line. We focus on a simplified system, a spherical gas cloud with homogeneous density and solid body rotation, to study the line morphology and the escape fraction in the presence of dust. We base our work on two independent Monte Carlo based radiative transfer codes CLARA (Forero-Romero et al. 2011) and XX (Dijkstra & Kramer 2012).

<sup>1</sup> Departamento de Física, Universidad de los Andes, Cra. 1 No. 18A-10, Edificio Ip, Bogotá, Colombia

<sup>2</sup> Max Planck Institute for Astrophysics, Karl-Schwarzschild-Str. 1, 85741, Garching, Germany

This paper is structured as follows. In §2 we present the implementation of bulk rotation into the Monte Carlo codes, paying special attention to coordinate definitions. We also list the different physical parameters in the simulated grid of models. In the next §3 we present the results of the simulations, with special detail to quantities in the line that show a clear evolution as a function of the sphere rotational velocity. In §4 we discuss the implications of our results in the interpretation of LAEs observations and further refinements that can be implemented in the theoretical modelling of the Ly $\alpha$  line. In the last section we present our conclusions.

In this paper we express a photon's frequency in terms of the dimensionless variable  $x \equiv (\nu - \nu_a)/\Delta\nu_\alpha$ , where  $\nu_\alpha = 2.46 \times 10^{15}$  Hz is the Ly $\alpha$  resonance frequency,  $\Delta\nu_\alpha \equiv \nu_\alpha \sqrt{2kT/m_p c^2} \equiv \nu_a v_{\text{th}}/c$  is the doppler broadening of the line which depends on the neutral gas temperature  $T$  scattering the radiation or equivalently the thermal velocity  $v_{\text{th}}$  of the atoms.

## 2. MODELS OF BULK GAS ROTATION

Describing the kinematics of gas rotation in all generality is a complex task, specially at high redshifts where there is still missing a thorough observational account of rotation in galaxies beyond  $z > 1.0$ . Even at low redshifts there is a great variation in the shape of the rotation curve as observed in HI emission as a function of the distance to the galaxy center. However there are two recurrent features. First, in the central galactic region the velocity increases proportional to the radius, following a solid rotation behaviour. Second, beyond a certain radius the rotation curve tends to flatten.

An ab-initio description of realistic rotation curves in simulations depends on having access to the dynamic evolution of all mass components in the galaxy: stars, gas and dark matter. Such level of realism is extremely complex to achieve, specially if one wants to get a systematic description based on statistics of simulated objects.

Following the tradition of studies of Ly $\alpha$  emitting systems, we implement a model with simplified geometry. We assume that the gas is homogeneously distributed in a sphere that rotates as a solid body with constant angular velocity. This simple model will contain only one free parameter: the linear velocity at the sphere's surface,  $V_{\text{max}}$ .

### 2.1. Detailed Implementation of Rotation

In the MonteCarlo code we define a cartesian coordinate system to describe the position of each photon. The origin of this system coincides with the center of the sphere and the rotation axis is defined to be  $z$ -axis. With this choice, the components of the gas bulk velocity field,  $\vec{v} = v_x \hat{i} + v_y \hat{j} + v_z \hat{k}$ , can be written as

$$v_x = -\frac{y}{R} V_{\text{max}}, \quad (1)$$

$$v_y = \frac{x}{R} V_{\text{max}}, \quad (2)$$

$$v_z = 0, \quad (3)$$

where  $R$  is the radius of the sphere and  $V_{\text{max}}$  is the linear velocity at the sphere's surface. The minus/plus sign in the  $x/y$ -component of the velocity indicates the direction

Physical Parameter (units)	Symbol	Values
Velocity (km s <sup>-1</sup> )	$V_{\text{max}}$	0, 50, 100, 200, 300
Hydrogen Optical Depth	$\tau_H$	10 <sup>5</sup> , 10 <sup>6</sup> , 10 <sup>7</sup>
Dust Optical Depth	$\tau_a$	0,1
Photons Distributions		Central, Homogeneous

TABLE 1

LIST OF THE PHYSICAL PARAMETERS THAT DEFINE THE SPHERICAL MODELS SIMULATED IN OUR MONTE CARLO CALCULATIONS. FOR EACH PARAMETER WE VARY THE VALUES IN THE RANGE LISTED IN THE LAST COLUMN. TAKING INTO ACCOUNT ALL THE POSSIBLE COMBINATIONS WE END UP WITH 60 DIFFERENT MODELS.

of rotation. In this case we take the angular velocity in the same direction as the  $\hat{k}$  unit vector. With these definitions we can write the angular velocity as  $\omega = V_{\text{max}}/R$ .

For each photon in the simulation we have its initial position inside the sphere, direction of propagation  $\hat{k}_{in}$  and reduced frequency  $x_{in}$ . The photon's propagation stops once they cross the surface of the sphere. At this point we store the position, the outgoing direction of propagation  $\hat{k}_{out}$  and the reduced frequency  $x_{out}$ . We define the angle  $\Theta$  by  $\cos \Theta = \hat{k} \cdot \hat{k}_{out}$ , that is the polar angle of the outgoing photon with respect to the  $z$  axis. Following Zheng & Wallace (2013) we make the study of the anisotropic emission in terms of this angle..

### 2.2. Grid of Simulated Galaxies

In the Monte Carlo calculations we follow the propagation of  $N_\gamma = 10^5$  numerical photons through different spherical galaxies. For each galaxy we vary at least one of the following parameters: the maximum rotational velocity  $V_{\text{max}}$ , the hydrogen optical depth  $\tau_H$ , the dust optical depth  $\tau_a$  and the initial distribution of photons with respect to the gas. There are 60 models in total combining all variations of the input parameters. Table 1 summarizes the models.

Additionally, we have used two independently developed Monte Carlo codes (Forero-Romero et al. 2011; Dijkstra & Kramer 2012) to perform the calculations of the non-dusty models. The results we report are robust in the sense that they are obtained by both codes.

## 3. RESULTS

The central result of this paper is summarized in Figure 1. It shows the evident impact of rotation on the morphology of the emergent Ly $\alpha$  line. Both panels focus on the results for  $\tau_H = 10^7$ , showing that the influence of rotation is present both when the photons are either homogeneously or centrally initialized over the gas volume.

In the next subsections we characterize the line morphology by the half-width at half intensity and the peak maxima. In order to interpret the morphological changes in the line we also report the median number of scatter for each Ly $\alpha$  photon in the simulation. For the models where dust is included we measure the escape fraction as a function of rotational velocity. Finally, we make an estimate of the anisotropic emission of the models in comparison with static spheres.

### 3.1. Line width and peak maxima

The first quantitative conclusion of the effect of rotation in the Ly $\alpha$  line is that double peaks broaden and re-




FIG. 1.— Shape of the Ly $\alpha$  line for different velocities rotational velocities for spherical distributions with  $\tau_H = 10^7$ . The left (right) panel shows the central (homogeneous) photon distribution. All photons were taken into account regardless of their final direction of propagation.




FIG. 2.— Half-width for the non-dusty models as a function of rotational velocity  $V_{\max}$ . Continuous (dashed) lines correspond to homogeneous (central) source distributions.

duce their intensity while the line center rises. This gives the impression that, as the rotational velocity increases, the double peaks are merged into a single broad emission peak. This is most evident at the highest rotational velocities for the homogeneously distributed sources (right panel in Figure 1).

To quantify the line broadening we use the width at half maximum (FWHM),  $W$ . Figure 2 shows how  $W$  increases with rotational velocity. Continuous (dashed) lines connect the results for homogeneous (central) source distribution. For the temperature  $T = 10^4\text{K}$  used in our radiative transfer calculations the thermal velocity is  $v_{th} = 12.8\text{ km s}^{-1}$ . For a model with  $\tau_H$  it means that the FWHM increase up to  $350\text{ km s}^{-1}$  (at  $V_{\max} = 300\text{ km s}^{-1}$ ) compared to  $150\text{ km s}^{-1}$  in the static case.

Figure 3 shows the position for the peak maxima as




FIG. 3.— Position of the peak maxima as a function of rotational velocity  $V_{\max}$ . Continuous (dashed) lines correspond to homogeneous (central) source distributions. A value of  $x_{\max} = 0$  indicates that line becomes single peaked.

a function of the rotational velocity  $V_{\max}$ . This Figure clearly shows that in the case of central distributed sources there is barely any change with rotational velocity in the range of explored parameters. However, in the case of homogeneously emitted sources the maxima position remain close to constant until beyond some velocity threshold the line becomes single peaked with  $x_m = 0\text{ km s}^{-1}$ .

The transition to a single peaked line seems to occur in systems where it is easier for the bulk of the photons to escape with the lowest number of scatterings possible, allowing them not to move very far from the center of the line. This might explain how the single peak stage is easily achieved in the homogeneous source distribution. In this case there is a fraction of the photons that are inside a photosphere region with  $\tau_{H,r} \ll \tau_H$  where  $\tau_{H,r}$  is the optical depth from the radius of emission to the




FIG. 4.— Logarithm of the average number of scatterings as function of rotational velocity. Continuous (dashed) lines represent an homogeneous (central) distribution of sources.

sphere’s surface. These conditions allows the photons to escape with much less scatterings compared to the photons emitted at the very center of the sphere. In turn, it gives the photons less scatterings to be placed far from the line center.

Increasing the rotational velocity  $V_{\max}$  reduces the optical depth making the photosphere region effectively larger, increasing the number of photons escaping close to the lines’s center. In our models we find the following correspondence between the optical depth  $\tau_H = \{10^5, 10^6, 10^7\}$  and the transitional velocities  $V_{\text{trans}} = \{50 - 100 \text{ km s}^{-1}, 100 - 200 \text{ km s}^{-1}, 200 - 300 \text{ km s}^{-1}\}$  which can only be constrained to be in the range of velocities in the models that gave a  $x_{\max} \neq 0$  and  $x_{\max} = 0$ .

For the central emission the transition to a single peak is never completed in the range of explored parameters. The non appearance of a single peak phase can be in part explained to the absence of a photosphere, as is the case in the homogeneous distribution. Nevertheless, there is a rise in the intensity at the line center as the rotational velocity increases. This hints that the encounters with a non static medium are inefficient in pushing photons outside the line’s center.

In the next sub-section we quantify the number of scatterings and explore to what extent this is related to the emergence of single peaked emission.

### 3.2. Average Number of Scatterings

The number of Ly $\alpha$  photon scatterings affects its final frequency after escaping the galaxy. In static gas geometries, a large value of the optical depth correlates with a high number of scatterings, increasing the probability of finding a Ly $\alpha$  photon far from the center of the line. In these cases the peak maxima shifts away from the line center as the amount of neutral hydrogen increases.

In Figure 4 we show the average number of scatterings  $\langle N_{\text{scatt}} \rangle$  as a function of the rotational velocity  $V_{\max}$ . For the central distributions the average number of scatterings has a modest change for increasing rotational ve-

locities,  $\langle N_{\text{scatt}} \rangle$  changes less than 0.5% for different velocities. In the same setup the average number of scatterings is proportional to the optical depth, as expected in analogy from the analytic result for the homogeneous infinite-slab  $\langle N_{\text{scatt}} \rangle = 1.612\tau_H$  (Adams 1972; Harrington 1973). In the spheres with centrally distributed sources we find  $\langle N_{\text{scatt}} \rangle = (1.50, 1.00, 0.92)\tau_H$  for optical depths  $\tau_H = (10^5, 10^6, 10^7)$ .

Figure 4 shows that for the homogeneous distribution there is a clear decrease of  $\langle N_{\text{scatt}} \rangle$  as the  $V_{\max}$  increases. This effect is more pronounced for lower values of the optical depth. For  $\tau_H = 10^5$  the average number of scatterings decreases by 61% at  $V_{\max} = 300 \text{ km s}^{-1}$  in comparison to the static case.

The analytic expectation for the slab with homogeneously distributed sources is  $\langle N_{\text{scatt}} \rangle = 1.16\tau_H$  (Harrington 1973), a factor of 0.72 lower than the case of the centrally emitted photons. In our case we find that for the static, while for homogeneously distributed source  $\langle N_{\text{scatt}} \rangle = (0.99, 0.59, 0.51)\tau_H$ , this represents factors of (0.66, 0.59, 0.55) lower than the centrally emitted photons.

In order to gain a deeper understanding of these results we make 2D histograms for the number of scatterings as a function of the outgoing dimensionless frequency  $x$ . In Figure 5 we show such histogram in the case  $\tau_H = 10^5$  for the static case and  $V_{\max} = 300 \text{ km s}^{-1}$ . The upper (lower) panels show the results for the homogeneous (central) source distribution. The color scale is logarithmic in the number of photons at a certain value  $x - N_{\text{scatt}}$ .

Figure 5 supports our hypothesis about the photosphere in the homogeneous distribution. In that case most of the photons that left with  $x \sim 0$  have escaped with less than 10 scatterings. This explains the origin of a single central peak. However, for a central distribution the situation is different. In this case the number of scatterings remains high, in the order of the optical depth, but the two peaks do get closer to each other. In this case the physical picture is that each scattering, due to the bulk velocity of the gas, is inefficient in driving the photon outside the line center.

### 3.3. Dusty Clouds: Escape Fraction

The average number of scatterings is also reflected in the amount of photons absorbed by dust. Under these considerations we do not expect any change in the escape fraction for a dusty cloud with central source of radiation given that the number of scatterings remains close to constant. On the other hand, in the case of a homogeneous radiation source the number of scatterings drops as  $V_{\max}$  increases, which has to be reflected as an increasing escape fraction.

Figure 6 shows the dependence of the escape fraction as a function of the maximum rotational velocity, confirming our expectations. For the central source distribution the escape fraction barely shows any change, while for the homogeneous case there is a clear increase in the escape fraction for high rotational velocities. Rotation has a higher relative impact in the models with low optical depth  $\tau_H = 10^5, 10^6$ , where it can rise from (0.26, 0.28) ( $V = 0 \text{ km s}^{-1}$ ) up to (0.37, 0.39) ( $V_{\text{rot}} = 300 \text{ km s}^{-1}$ ).

In Figure 7 we put these results in the context of the analytical solution for the infinite slab (Neufeld 1990). In




FIG. 5.— 2D histogram of  $N_{\text{scatt}}$  vs  $x$ . The upper (lower) panels show the homogeneous (central) source distribution. Left corresponds to the static case and the right  $V_{\text{max}} = 300 \text{ km/s}$ . The colour scale is logarithmic on the number of photons with given values of  $N_{\text{scatt}}$  and  $x$ .

Neufeld's set-up the analytic solution depends uniquely on the product  $(a\tau_{\text{H}})^{1/3}\tau_a$ , an approximation that is valid only in the limits where  $a\tau_{\text{H}} \gg 1$ . The dashed lines in Figure 7 correspond to the cases of different velocities. First of all we note that the escape fraction does not increase from  $\tau_{\text{H}} = 10^5$  to  $\tau_{\text{H}} = 10^6$ , this is explained since we are in a regime where the condition for the analytic expectations ( $a\tau_h \gg 1$ ) does not hold. This figure also shows that the escape fraction is higher by factors of 2 – 10 than the expectation for a slab with centrally distributed sources. Furthermore, the decrease of the escape fraction with increasing optical depth is also slower.

#### 3.4. Off-Centred emission

The model with homogeneous radiation sources allows to study the effect of radiation sources distributed with a different symmetry than the gas. We select photons emitted in spheres that are off-centred with respect to

the gas distribution.

Figure 8 shows the two spherical regions we define. Each has a radius of  $0.5R$  and can have two possible centres positions  $C$  along the  $\hat{i}$  direction:  $\pm C\hat{i}$ . We also place an observer along the  $\hat{j}$  direction, such that the emitting sphere centred on  $+C\hat{i}$  moves away from the observer and the sphere centred on  $-C\hat{i}$  moves towards the observer. We consider two values for  $C = (0.25, 0.50)R$ .

We study first the deviations of the received flux on the surface of the sphere. We follow Zheng & Wallace (2013) to estimate the flux seen by an observer located at an angle  $\Theta$  (distant observer angle) normalized to the isotropic flux:

$$F(\mu) = \frac{2\Delta N}{N\Delta\mu}, \quad (4)$$

Where  $\mu = \cos \Theta$ ,  $N$  is the total number of outgoing pho-




FIG. 6.— Escape fraction as a function of rotational velocity. The continuous (dashed) lines correspond to homogeneous (central) models.




FIG. 7.— Escape fraction as a function of the product  $(a\tau_H)^{1/3}\tau_a$ . The analytic solution for the infinite slab is shown as a continuous line. Different dashed lines correspond to different rotational velocities.

tons,  $\Delta N$  is the number of photons in an angular bin  $\Delta\Theta$ . This definition satisfies the condition  $\int_{-1}^1 F(\mu) d\mu/2 = 1$ .

For the central and homogeneous source distributions we find that the level of isotropy angular dependency represented by  $\mu$  doesn't show a considerable variation in particular the observer would infer luminosities different from the isotropical value only at levels  $< 3\%$ . This values were obtained for a run with  $V_{\max} = (50 \text{ km s}^{-1}, 100 \text{ km s}^{-1}, 200 \text{ km s}^{-1}, 300 \text{ km s}^{-1})$  and  $\tau_H = 10^5, 10^6, 10^7$  and  $N_\gamma = 10^5$  photons.

For the Off-center emission Fig.9 the variations of  $\mu$  increase as the rotational velocity increase, for  $V_{\max} = (50 \text{ km s}^{-1} \Delta\mu \leq 7\%, \text{ for } V_{\max} = (100 \text{ km s}^{-1} \Delta\mu \leq 12\%$




FIG. 8.— The two solid circles show the two different spherical off-centered emitting regions. The angular velocity vector is defined to be along the  $\hat{i}$  direction and the observer is along the  $\hat{j}$  direction (where the reader is).




FIG. 9.— Flux angular dependency for Off-center distribution, different line widths illustrate different off-center positions.

and finally for  $V_{\max} = (200 \text{ km s}^{-1} \Delta\mu \leq 15\%$ . This would imply that the amount of flux we measure would depend in the angle between our line of sight and the rotation axis of the galaxy.

The outgoing spectra for two different spheres located at  $\pm 0.5R$  are shown in Fig.10, we have shown this positions because the resulting profiles show new interesting morphologies than are not found in the  $\pm 0.25R$  case.

The main effect we found in the off-center emission is that for some models ( $\tau = 10^6, V = 100 \text{ km s}^{-1}$  and  $\tau = 10^7, V = 200 \text{ km s}^{-1}$ ) the line presents three peaks (blue-shifted, centred, red-shifted), to obtain clearest profiles concerning this results we run three models with  $\tau_\gamma = 10^6$ , This profiles are shown in Fig.11, where we clearly observe the presence of three peaks, and also we do not find any asymmetry in the line produced by the off-center emission.

For models with  $([\tau = 10^5, V = 100 \text{ km s}^{-1}], [\tau =$

$10^6, V = 200 \text{ km s}^{-1}$ ] and  $[\tau = 10^7, V = 300 \text{ km s}^{-1}]$ ) the centred line is dominant but the presence of the redder and bluer peak is also present which is also a new result. Also as in the homogeneous / central model the line is broader at high rotational velocities.

#### 4. DISCUSSION

Gas bulk rotation has a noticeable effect on the morphology of the Ly $\alpha$  line. In this section we discuss the implications of these findings for the interpretation of available observational data. Our discussion will focus on the qualitative aspects of diverse morphological features observed by Kulas et al. (2012) and Yamada et al. (2012).

We follow a classification of Ly $\alpha$  profiles made by Kulas et al. (2012). They presented observational results for star forming galaxies at  $z = 2-3$  that showed multiple peaks in their emission. They classified the observed lines into five groups. In Group I the red peak is stronger than the blue, while in Group II the blue peak is stronger, in both cases the peaks are symmetrically located around the line center. In Group III the blue peak also dominates but there is an overall redward shift of the zero point. Group IV presents two similar peaks symmetrically located around the zero point. Group V presents three peaks.

The asymmetries in Groups I, II and III cannot be fully explained by rotation. However, some galaxies in those groups have a high flux between the two peaks; an effect that can be induced by rotation. The same argument applies to galaxies in the group IV in which the flux in between of the two peaks is high ( $\geq 70\%$  of the intensity in the highest peak). Finally, the triple peaked lines in Galaxies in Group V can be easily reproduced by rotation with off-centred emission.

Another relevant sample of lines was presented by Yamada et al. (2012) for 91 LAEs at  $z = 3.1$ . They find that  $\sim 33\%$  of the galaxies present a single peak profile. Some of the peaks have a strong symmetry, while others do not. The symmetric single peaked profile can be explained by rotation effects, while the asymmetric cases naturally arise in outflow/inflow models (Verhamme et al. 2008; Dijkstra et al. 2006). Among the 91 LAEs, 8 reveal triple peaked profiles with a central peak and two smaller peaks at the redder and bluer side, a feature that is present in our off-centre emission models with rotation.

The presence of single peaked profiles has been associated to inflow/outflow dynamics (?). In this paper we show that gas bulk rotation can also be considered as a probable origin for that behaviour, provided that the observed single peak is highly symmetric. Similarly, in the case of double peaked lines with a high level of flux at the line center, rotation also deserves to be considered in the pool of possible bulk flows responsible for that feature, specially if the two peaks have similar intensities. The case of triple peaked lines is another clear feature of gas rotation, this time with the additional condition of an off-centred emission.

#### 5. CONCLUSIONS

In this paper, we have estimated the effects of gas bulk rotation on the morphology of the Lyman  $\alpha$  line in star forming galaxies. The results are based on the study of

a homogeneous sphere rotating as a solid body. We explored a range of models by varying the rotation speed, hydrogen optical depth, dust optical depth and initial distribution of Ly $\alpha$  photons with respect to the gas density. As a cross-validation, we obtain our results from two independently developed Monte-Carlo radiative transfer codes.

Our main result is that rotation clearly impacts the Ly $\alpha$  line morphology. Double peaked lines can make transitions into single peaked lines as the rotational velocity increases, off-centred can result into triple peaked lines.

We have quantified the results in different aspects. In Section 3.1 we show that the line width increases up to a factor of 2 – 3 in comparison to the static case. Simultaneously, the flux in the line center also increases with rotational velocity. This increase induces the transition from a double peak to a single peak line in the models where the radiation sources are homogeneously distributed.

In the case of homogeneously distributed sources, the changes in line morphology are linked to the reduction in the average number of scatterings. In the second part of Section 3.1 we show that, as rotational velocity increases, a large fraction of photons escape with less than 10 scatterings. On average there are 40% less scatterings. This makes highly improbable that a photon's frequency can move away from the line's center.

In contrast, for centrally located sources, the number of scatterings does not significantly decrease. However, due to the increasing bulk velocity of the gas, each scattering is inefficient in shifting photons away from the line's center.

These changes in the number of scatterings have a direct bearing on the escape fraction in the presence of dust. In Section 3.3 we find that the escape fraction increases in 20% – 30% for the homogeneous source distribution. In the centrally located emission, the rotational effects are negligible on the escape fraction.

As many observational spectra Ly $\alpha$  emission line (Kulas et al, Yamada et al) is double peaked, these peaks provide important information concerning gas kinematics and geometry, which can be partially explained with inflows/outflows of gas content. We study the effect of rotation in the position of these peaks, and we find that the position of the maxima does change with rotation for the homogeneous models when the double peak merged into a single peak as velocity increases. This effect is not seen for the central distribution when the double peak remains constant as the velocity increases. We also find that there is no dependency in the observer viewing angle with the maxima position.

Comparing our results with recent observed LAEs we find that many morphological features such as high central line flux, single peak profiles and multi-peaked profiles could be explained by gas bulk rotation present in these LAEs.

This paper illustrates for the first time the main effects of rotation in the morphology of the Ly $\alpha$  emission line, we estimate the range of these effects for simplified models.

#### ACKNOWLEDGEMENTS




FIG. 10.— Ly $\alpha$  profiles for the off-center positions. dashed lines represents spheres locate at  $-0.5R$  while solid lines represents spheres at  $0.5R$ )

## REFERENCES

- Adams, T. F. 1972, *ApJ*, 174, 439  
Ahn, S.-H., Lee, H.-W., & Lee, H. M. 2000, *Journal of Korean Astronomical Society*, 33, 29  
—. 2001, *ApJ*, 554, 604  
Auer, L. H. 1968, *ApJ*, 153, 783  
Avery, L. W., & House, L. L. 1968, *ApJ*, 152, 493  
Barnes, L. A., Haehnelt, M. G., Tescari, E., & Viel, M. 2011, *MNRAS*, 416, 1723  
Behrens, C., & Niemeyer, J. 2013, *A&A*, 556, A5  
Dijkstra, M., Haimes, Z., & Spaans, M. 2006, *ApJ*, 649, 14  
Dijkstra, M., & Kramer, R. 2012, *MNRAS*, 424, 1672  
Forero-Romero, J. E., Yepes, G., Gottlöber, S., Knollmann, S. R., Cuesta, A. J., & Prada, F. 2011, *MNRAS*, 415, 3666  
Forero-Romero, J. E., Yepes, G., Gottlöber, S., & Prada, F. 2012, *MNRAS*, 419, 952  
Gawiser, E., Francke, H., Lai, K., Schawinski, K., Gronwall, C., Ciardullo, R., Quadri, R., Orsi, A., Barrientos, L. F., Blanc, G. A., Fazio, G., & Feldmeier, J. J. 2007, *ApJ*, 671, 278  
Hansen, M., & Oh, S. P. 2006, *MNRAS*, 367, 979  
Harrington, J. P. 1973, *MNRAS*, 162, 43  
Koehler, R. S., Schuecker, P., & Gebhardt, K. 2007, *A&A*, 462, 7  
Kulas, K. R., Shapley, A. E., Kollmeier, J. A., Zheng, Z., Steidel, C. C., & Hainline, K. N. 2012, *ApJ*, 745, 33  
Laursen, P., Sommer-Larsen, J., & Andersen, A. C. 2009, *ApJ*, 704, 1640  
Loeb, A., & Rybicki, G. B. 1999, *ApJ*, 524, 527  
Neufeld, D. A. 1990, *ApJ*, 350, 216  
Orsi, A., Lacey, C. G., & Baugh, C. M. 2012, *MNRAS*, 425, 87  
Ouchi, M., Shimasaku, K., Akiyama, M., Simpson, C., Saito, T., Ueda, Y., Furusawa, H., Sekiguchi, K., Yamada, T., Kodama, T., Kashikawa, N., Okamura, S., Iye, M., Takata, T., Yoshida, M., & Yoshida, M. 2008, *ApJS*, 176, 301  
Partridge, R. B., & Peebles, P. J. E. 1967, *ApJ*, 147, 868  
Rhoads, J. E., Malhotra, S., Dey, A., Stern, D., Spinrad, H., & Jannuzi, B. T. 2000, *ApJ*, 545, L85






FIG. 11.— Ly $\alpha$  profiles for the off-center positions. green lines represents spheres locate at  $-0.5R$  while red lines represents spheres at  $0.5R$ )

Schenker, M. A., Stark, D. P., Ellis, R. S., Robertson, B. E.,  
Dunlop, J. S., McLure, R. J., Kneib, J.-P., & Richard, J. 2012,  
ApJ, 744, 179  
Verhamme, A., Dubois, Y., Blaizot, J., Garel, T., Bacon, R.,  
Devriendt, J., Guiderdoni, B., & Slyz, A. 2012, A&A, 546, A111  
Verhamme, A., Schaerer, D., Atek, H., & Tapken, C. 2008, 111, 89  
Verhamme, A., Schaerer, D., & Maselli, A. 2006, A&A, 460, 397

Yajima, H., Li, Y., Zhu, Q., Abel, T., Gronwall, C., & Ciardullo,  
R. 2012, ApJ, 754, 118  
Yamada, T., Nakamura, Y., Matsuda, Y., Hayashino, T.,  
Yamauchi, R., Morimoto, N., Kousai, K., & Umemura, M.  
2012, AJ, 143, 79  
Zheng, Z., & Miralda-Escudé, J. 2002, ApJ, 578, 33  
Zheng, Z., & Wallace, J. 2013, ArXiv e-prints