# BDA Project: Hurricane forecasting in Stan

José Miguel Ramírez & Jonas Lindblad Aalto University

▶ Destructive storms occuring in the late summer and fall in the northern hemisphere's tropical region.



- Destructive storms occurring in the late summer and fall in the northern hemisphere's tropical region.
- Classified by their wind intensity at the eye wall.



- Destructive storms occurring in the late summer and fall in the northern hemisphere's tropical region.
- Classified by their wind intensity at the eye wall.
- They can cause extreme levels of flooding and destroy many buildings.



- Destructive storms occurring in the late summer and fall in the northern hemisphere's tropical region.
- Classified by their wind intensity at the eye wall.
- ▶ They can cause extreme levels of flooding and destroy many buildings.
- Monetary damages and loss of lives increase with an almost exponential character as a function of storm intensity.



► Forecasters predict two quantities: track and intensity

- Forecasters predict two quantities: track and intensity
- ▶ Two kinds of models: dynamical and statistical

- Forecasters predict two quantities: track and intensity
- ► Two kinds of models: dynamical and statistical
  - Dynamical: simulate the laws of physics

- ► Forecasters predict two quantities: track and intensity
- ► Two kinds of models: dynamical and statistical
  - Dynamical: simulate the laws of physics
  - Statistical: estimate based on historical data

- Forecasters predict two quantities: track and intensity
- ► Two kinds of models: dynamical and statistical
  - Dynamical: simulate the laws of physics
    - Statistical: estimate based on historical data
- Dynamical vs. statistical: good at long- and short-range forecasts respectively

- ► Forecasters predict two quantities: track and intensity
- Two kinds of models: dynamical and statistical
  - Dynamical: simulate the laws of physics
    - Statistical: estimate based on historical data
- Dynamical vs. statistical: good at long- and short-range forecasts respectively
- ... but for hurricane forecasts short-range is usually more interesting

- Forecasters predict two quantities: track and intensity
- Two kinds of models: dynamical and statistical
  - Dynamical: simulate the laws of physics
    - Statistical: estimate based on historical data
- Dynamical vs. statistical: good at long- and short-range forecasts respectively
- ...but for hurricane forecasts short-range is usually more interesting
- Rapid intensification: forecasted better by dynamical models

- Forecasters predict two quantities: track and intensity
- ► Two kinds of models: dynamical and statistical
  - Dynamical: simulate the laws of physics
  - Statistical: estimate based on historical data
- Dynamical vs. statistical: good at long- and short-range forecasts respectively
- ...but for hurricane forecasts short-range is usually more interesting
- Rapid intensification: forecasted better by dynamical models
- ► This project: a *statistical* model for *intensity*

The US government forecasting agency, the National Hurricane Center (NHC), uses a large number of models operationally. The models (together: the *model ensemble*) are used together with experienced meteorologists' judgment to provide the official forecast.

Surprisingly, the best single (short-range) model is a multiple linear regression!

- Surprisingly, the best single (short-range) model is a multiple linear regression!
- ► The NHC regression model: Statistical Hurricane Intensity Prediction Scheme (SHIPS)


- Surprisingly, the best single (short-range) model is a multiple linear regression!
- ► The NHC regression model: Statistical Hurricane Intensity Prediction Scheme (SHIPS)
- ➤ SHIPS: ~140 covariates, many calculated from data sources more easily available to the NHC

- Surprisingly, the best single (short-range) model is a multiple linear regression!
- ► The NHC regression model: Statistical Hurricane Intensity Prediction Scheme (SHIPS)
- ➤ SHIPS: ~140 covariates, many calculated from data sources more easily available to the NHC
- ► SHIPS dataset: publically available with no restrictions (link: SHIPS Development)


- Surprisingly, the best single (short-range) model is a multiple linear regression!
- ► The NHC regression model: Statistical Hurricane Intensity Prediction Scheme (SHIPS)
- ➤ SHIPS: ~140 covariates, many calculated from data sources more easily available to the NHC
- SHIPS dataset: publically available with no restrictions (link: SHIPS Development)
- ... but the documentation is terrible

- Surprisingly, the best single (short-range) model is a multiple linear regression!
- ► The NHC regression model: Statistical Hurricane Intensity Prediction Scheme (SHIPS)
- ➤ SHIPS: ~140 covariates, many calculated from data sources more easily available to the NHC
- SHIPS dataset: publically available with no restrictions (link: SHIPS Development)
- ... but the documentation is terrible
- ► SHIPS: only a point estimate; our project: a predictive distribution

### Hurricane forecasting basics: our selection




The SHIPS Blunder: a simple linear regression




Model 2: regression with skewness

errors not symmetric around the mean prediction!




Model 2: regression with skewness

- errors not symmetric around the mean prediction!
- rapid intensification!




Model 3: regression with a linear model for standard deviation

fewer storms reach higher values of VMAX



Model 3: regression with a linear model for standard deviation

- fewer storms reach higher values of VMAX
- allow for higher variance to account for larger historical uncertainty



# Model: limitations

Model: posterior predictive checking








# Forecasting

# Forecasting: Model Comparison (1)

Dataset comparison for the linear regression model (LOOCV)


Dataset	elpd_diff	se_diff
С	0.0	0.0
В	-25.0	6.5
Α	-27.4	6.3



# Forecasting: Model Comparison (2)

Dataset comparison for the skewed regression model (LOOCV)


Dataset	elpd_diff	se_diff
С	0.0	0.0
В	-23.2	6.2
Α	-28.7	6.2



# Forecasting: Model Comparison (3)

Dataset comparison for the Changing variance model (LOOCV)


Dataset	elpd_diff	se_diff
С	0.0	0.0
В	-32.6	8.2
Α	-37.1	8.2



# Forecasting: Model Comparison (4)

Model comparison using the Dataset C (LOOCV)

Model	elpd_diff	se_diff
Variance	0.0	0.0
Skew	-176.3	27.8
Linear	-205.5	34.9



## Problems to solve & development ideas

variable selection in full SHIPS dataset

## Problems to solve & development ideas

- variable selection in full SHIPS dataset
- more time series autoregressive components

## Problems to solve & development ideas

- variable selection in full SHIPS dataset
- more time series autoregressive components
- use LGEM model (will explain)



### Additional information

#### R Markdown

This is an R Markdown presentation. Markdown is a simple formatting syntax for authoring HTML, PDF, and MS Word documents. For more details on using R Markdown see http://rmarkdown.rstudio.com.

When you click the **Knit** button a document will be generated that includes both content as well as the output of any embedded R code chunks within the document.

### Slide with Bullets

▶ Bullet 1

### Slide with Bullets

- ▶ Bullet 1
- ▶ Bullet 2

#### Slide with Bullets

- ▶ Bullet 1
- ▶ Bullet 2
- ▶ Bullet 3

### Slide with R Output

#### summary(cars)

```
##
       speed
                     dist
##
   Min. : 4.0
                Min. : 2.00
##
   1st Qu.:12.0
                1st Qu.: 26.00
##
   Median: 15.0 Median: 36.00
##
   Mean :15.4
                Mean : 42.98
##
   3rd Qu.:19.0
                3rd Qu.: 56.00
##
   Max. :25.0
                Max. :120.00
```

#### Slide with Plot

