

BDA Project: Hurricane forecasting in Stan

José Miguel Ramírez & Jonas Lindblad
Aalto University

Hurricane introduction

Hurricane introduction

- Destructive storms occurring in the late summer and fall in the northern hemisphere's tropical region.



Hurricane introduction

- Destructive storms occurring in the late summer and fall in the northern hemisphere's tropical region.
- Classified by their wind intensity at the eye wall.



Hurricane introduction

- Destructive storms occurring in the late summer and fall in the northern hemisphere's tropical region.
- Classified by their wind intensity at the eye wall.
- They can cause extreme levels of flooding and destroy many buildings.



Hurricane introduction

- Destructive storms occurring in the late summer and fall in the northern hemisphere's tropical region.
- Classified by their wind intensity at the eye wall.
- They can cause extreme levels of flooding and destroy many buildings.
- Monetary damages and loss of lives increase with an almost exponential character as a function of storm intensity.



Hurricane forecasting basics

- Forecasters predict two quantities: track and intensity

Hurricane forecasting basics

- Forecasters predict two quantities: track and intensity
- Two kinds of models: *dynamical* and *statistical*

Hurricane forecasting basics

- Forecasters predict two quantities: track and intensity
- Two kinds of models: *dynamical* and *statistical*
 - Dynamical: simulate the laws of physics

Hurricane forecasting basics

- Forecasters predict two quantities: track and intensity
- Two kinds of models: *dynamical* and *statistical*
 - Dynamical: simulate the laws of physics
 - Statistical: estimate based on historical data

Hurricane forecasting basics

- Forecasters predict two quantities: track and intensity
- Two kinds of models: *dynamical* and *statistical*
 - Dynamical: simulate the laws of physics
 - Statistical: estimate based on historical data
- Dynamical vs. statistical: good at long- and short-range forecasts respectively

Hurricane forecasting basics

- Forecasters predict two quantities: track and intensity
- Two kinds of models: *dynamical* and *statistical*
 - Dynamical: simulate the laws of physics
 - Statistical: estimate based on historical data
- Dynamical vs. statistical: good at long- and short-range forecasts respectively
- ... but for hurricane forecasts short-range is usually more interesting

Hurricane forecasting basics

- Forecasters predict two quantities: track and intensity
- Two kinds of models: *dynamical* and *statistical*
 - Dynamical: simulate the laws of physics
 - Statistical: estimate based on historical data
- Dynamical vs. statistical: good at long- and short-range forecasts respectively
- ... but for hurricane forecasts short-range is usually more interesting
- *Rapid intensification*: forecasted better by dynamical models

Hurricane forecasting basics

- Forecasters predict two quantities: track and intensity
- Two kinds of models: *dynamical* and *statistical*
 - Dynamical: simulate the laws of physics
 - Statistical: estimate based on historical data
- Dynamical vs. statistical: good at long- and short-range forecasts respectively
- ... but for hurricane forecasts short-range is usually more interesting
- *Rapid intensification*: forecasted better by dynamical models
- This project: a *statistical* model for *intensity*

Hurricane forecasting basics: the SHIPS data

The US government forecasting agency, the National Hurricane Center (NHC), uses a large number of models operationally. The models (together: the *model ensemble*) are used together with experienced meteorologists' judgment to provide the official forecast.

- Surprisingly, the best *single* (short-range) model is a multiple linear regression!

Hurricane forecasting basics: the SHIPS data

The US government forecasting agency, the National Hurricane Center (NHC), uses a large number of models operationally. The models (together: the *model ensemble*) are used together with experienced meteorologists' judgment to provide the official forecast.

- Surprisingly, the best *single* (short-range) model is a multiple linear regression!
- The NHC regression model: Statistical Hurricane Intensity Prediction Scheme (SHIPS)

Hurricane forecasting basics: the SHIPS data

The US government forecasting agency, the National Hurricane Center (NHC), uses a large number of models operationally. The models (together: the *model ensemble*) are used together with experienced meteorologists' judgment to provide the official forecast.

- Surprisingly, the best *single* (short-range) model is a multiple linear regression!
- The NHC regression model: Statistical Hurricane Intensity Prediction Scheme (SHIPS)
- SHIPS: ~140 covariates, many calculated from data sources more easily available to the NHC

Hurricane forecasting basics: the SHIPS data

The US government forecasting agency, the National Hurricane Center (NHC), uses a large number of models operationally. The models (together: the *model ensemble*) are used together with experienced meteorologists' judgment to provide the official forecast.

- Surprisingly, the best *single* (short-range) model is a multiple linear regression!
- The NHC regression model: Statistical Hurricane Intensity Prediction Scheme (SHIPS)
- SHIPS: ~140 covariates, many calculated from data sources more easily available to the NHC
- SHIPS dataset: publically available with no restrictions (link: SHIPS Development)

Hurricane forecasting basics: the SHIPS data

The US government forecasting agency, the National Hurricane Center (NHC), uses a large number of models operationally. The models (together: the *model ensemble*) are used together with experienced meteorologists' judgment to provide the official forecast.

- Surprisingly, the best *single* (short-range) model is a multiple linear regression!
- The NHC regression model: Statistical Hurricane Intensity Prediction Scheme (SHIPS)
- SHIPS: ~140 covariates, many calculated from data sources more easily available to the NHC
- SHIPS dataset: publically available with no restrictions (link: SHIPS Development)
- ... but the documentation is terrible

Hurricane forecasting basics: the SHIPS data

The US government forecasting agency, the National Hurricane Center (NHC), uses a large number of models operationally. The models (together: the *model ensemble*) are used together with experienced meteorologists' judgment to provide the official forecast.

- Surprisingly, the best *single* (short-range) model is a multiple linear regression!
- The NHC regression model: Statistical Hurricane Intensity Prediction Scheme (SHIPS)
- SHIPS: ~140 covariates, many calculated from data sources more easily available to the NHC
- SHIPS dataset: publically available with no restrictions (link: SHIPS Development)
- ... but the documentation is terrible
- SHIPS: only a point estimate; our project: a predictive distribution

The SHIPS developmental data is confusing!

[illegible]

Hurricane forecasting basics: the SHIPS data

We are making *synoptic* models.


$$T = 0$$
[illegible]

[illegible]

Hurricane forecasting basics: our selection

We have not done statistical variable selection. Choice of variable subset is based on theory.

- **CSST**: (climatological) sea surface temperature
- **RHLO**: low-altitude relative humidity
- **T200**: air temperature at 200 mb (very high altitude)
- **SHRD**: wind shear between 850 and 200 mb
- **VMPI**: maximum potential intensity




- for testing, we have variable sets *A*, *B*, *C*
- *A*: LAT/LON, VMAX, CSST, SHRD
- *B*: LAT/LON, VMAX, CSST, SHRD, VMPI
- *C*: LAT/LON, VMAX, CSST, SHRD, VMPI, RHLO, T200

Intensity change predictive model

Intensity change predictive model


The SHIPS Blunder: a simple linear regression



Intensity change predictive model

Model 2: regression with skewness


- errors not symmetric around the mean prediction!



Intensity change predictive model

Model 2: regression with skewness


- errors not symmetric around the mean prediction!
- rapid intensification!



Intensity change predictive model

Model 3: regression with a linear model for standard deviation


- fewer storms reach higher values of VMAX



Intensity change predictive model

Model 3: regression with a linear model for standard deviation

- fewer storms reach higher values of VMAX
- allow for higher variance to account for larger historical uncertainty



Models: some remarks

- SHIPS: predict V_{max} ; our models: predict ΔV_{max}

Models: some remarks

- SHIPS: predict V_{max} ; our models: predict ΔV_{max}
- we also decided to include latitude and longitude as variables; geographic region is known to be important for storm development

Models: some remarks

- SHIPS: predict V_{max} ; our models: predict ΔV_{max}
- we also decided to include latitude and longitude as variables; geographic region is known to be important for storm development
- reminder: we do 12-hour forecasts

Models: some remarks

- SHIPS: predict V_{max} ; our models: predict ΔV_{max}
- we also decided to include latitude and longitude as variables; geographic region is known to be important for storm development
- reminder: we do 12-hour forecasts
- we standardized all of our data; priors chosen to be weak in standardized scale

Models: some remarks

- SHIPS: predict V_{max} ; our models: predict ΔV_{max}
- we also decided to include latitude and longitude as variables; geographic region is known to be important for storm development
- reminder: we do 12-hour forecasts
- we standardized all of our data; priors chosen to be weak in standardized scale
- SHIPS data: 1982-2019; our restriction: 2017-2019

Models: some remarks

- SHIPS: predict V_{max} ; our models: predict ΔV_{max}
- we also decided to include latitude and longitude as variables; geographic region is known to be important for storm development
- reminder: we do 12-hour forecasts
- we standardized all of our data; priors chosen to be weak in standardized scale
- SHIPS data: 1982-2019; our restriction: 2017-2019
- poor problem setup! True model is the laws of physics, but we are fitting a regression

Models: some remarks




- SHIPS: predict V_{max} ; our models: predict ΔV_{max}
- we also decided to include latitude and longitude as variables; geographic region is known to be important for storm development
- reminder: we do 12-hour forecasts
- we standardized all of our data; priors chosen to be weak in standardized scale
- SHIPS data: 1982-2019; our restriction: 2017-2019
- poor problem setup! True model is the laws of physics, but we are fitting a regression
- models were programmed in Stan; sampling with `rstan` using default parameters; resulted in no divergences or issues except for the skew model and it was solved by increasing max tree depth to 15

Forecasting: Model Comparison

Forecasting: Model Comparison

Dataset comparison for the **linear regression model** (LOOCV)




Dataset	elpd_diff	se_diff
C	0.0	0.0
<i>B</i>	-25.0	6.5
<i>A</i>	-27.4	6.3



Forecasting: Model Comparison (2)

Dataset comparison for the **skewed regression model** (LOOCV)




Dataset	elpd_diff	se_diff
C	0.0	0.0
<i>B</i>	-23.2	6.2
<i>A</i>	-28.7	6.2



Forecasting: Model Comparison (3)

Dataset comparison for the **Changing variance model** (LOOCV)

Dataset	elpd_diff	se_diff
<i>C</i>	0.0	0.0
<i>B</i>	-32.6	8.2
<i>A</i>	-37.1	8.2




Forecasting: Model Comparison (4)

Model comparison using the Dataset C (LOOCV)


Model	elpd_diff	se_diff
Variance	0.0	0.0
Skew	-176.3	27.8
Linear	-205.5	34.9

Marginal posteriors




Forecasting: checking predictions

Hurricane Dorian 2019. The image shows a 90% credible interval.
Black dotted line: true VMAX




Forecasting: checking predictions

Hurricane Florence 2018. The image shows a 90% credible interval.
Black dotted line: true VMAX



Forecasting: checking predictions

Hurricane Helene 2018. The image shows a 90% credible interval.
Black dotted line: true VMAX



Concluding section

Concluding section

Further development ideas:

- variable selection in full SHIPS dataset

Concluding section

Further development ideas:

- variable selection in full SHIPS dataset
- more time series autoregressive components

Concluding section

Further development ideas:

- variable selection in full SHIPS dataset
- more time series autoregressive components
- use LGEM model (will explain)

A final takeaway

Talking about the 2017 category 5 hurricane Irma:

“The hurricane will be moving through an environment of low vertical wind shear, a moist mid-level atmosphere, and increasing upper-ocean heat content, and this is ideal for some additional intensification. However, given that the SHIPS models do not show any significant change in the intensity, the NHC forecast keeps Irma a powerful hurricane through five days.” — NHC 5AM advisory Sep. 5th

While the NHC published the advisory, Irma was undergoing rapid intensification from 130 to 175 knots over a span of only a few hours.

Conclusions & contact info

- Takeaway: the SHIPS model is terrible!
- Simple changes to the predictive distribution can improve the model
- Further development using Bayesian methods seems promising and there are several possible directions

More at our Github repo:

<https://github.com/jnlb/bda-hurricane-modeling>

The SHIPS website: http://rammb.cira.colostate.edu/research/tropical_cyclones/ships/index.asp

Contact info:

José Miguel Ramírez

- rocket-chat: @jose

Jonas Lindblad

- rocket-chat: @jnlb

Additional information

The SHIPS model:

$$y_i \sim \mathcal{N}(\alpha + X_i \cdot \beta_{N-1}, \sigma), \quad i = 1, \dots, r,$$

where we let X_i denote the i :th row of the data, β_{N-1} is an $N - 1$ -dimensional parameter vector, and r is the number of observations (rows) in the data. Its priors were

$$\begin{bmatrix} \alpha_0 \\ \beta_{N-1,0} \end{bmatrix} \sim \mathcal{N}(\mathbf{0}_N, 10 \cdot \mathbf{I}_N), \sigma_0 \sim \text{Inv-}\chi^2\left(\frac{1}{10}\right).$$

The skew-normal regression model:

$$y_i \sim \text{SkewNormal}(\alpha + X_i \cdot \beta_{N-1}, \sigma, \psi), \quad i = 1, \dots, r,$$

with priors

$$\begin{bmatrix} \alpha_0 \\ \beta_{N-1,0} \end{bmatrix} \sim \mathcal{N}(\mathbf{0}_N, 10 \cdot \mathbf{I}_N), \sigma_0 \sim \text{Inv-}\chi^2\left(\frac{1}{10}\right), \psi_0 \sim \mathcal{N}(0, 1).$$

Additional information

The variance model:

$$y_i \sim \mathcal{N}(\alpha + X_i \cdot \beta_{N-1}, \sigma + \gamma |V_{\max,i}|), \quad i = 1, \dots, r,$$

where, again, we use the same notation as before and let $V_{\max,i}$ denote the V_{\max} -value of the i :th row. The priors that were fed into Stan were

$$\begin{bmatrix} \alpha_0 \\ \beta_{N-1,0} \end{bmatrix} \sim \mathcal{N}(\mathbf{0}_N, 10 \cdot \mathbf{I}_N), \quad \sigma_0 \sim \text{Inv-}\chi^2\left(\frac{1}{10}\right), \quad \gamma_0 \sim \Gamma(1, 1).$$

Additional information

The variance regression was run in rstan with the following options

```
variance_m <- rstan::stan_model(file = file.path(mod_path,
  "minimal3.stan"))
variance_model <- rstan::sampling(variance_m,
  data = stan_data,
  control = list(max_treedepth = 10),
  iter=4000, seed = SEED)
```


Additional information




Image from *Introduction to Tropical Meteorology, 2nd Ed.*, 2011, by A. Laing & J-L Evans.

Additional information




Image from *Introduction to Tropical Meteorology, 2nd Ed.*, 2011, by A. Laing & J-L Evans.