## BDA Project: Hurricane forecasting in Stan

José Miguel Ramírez & Jonas Lindblad Aalto University

▶ Destructive storms occuring in the late summer and fall in the northern hemisphere's tropical region.



- Destructive storms occurring in the late summer and fall in the northern hemisphere's tropical region.
- Classified by their wind intensity at the eye wall.



- Destructive storms occurring in the late summer and fall in the northern hemisphere's tropical region.
- Classified by their wind intensity at the eye wall.
- They can cause extreme levels of flooding and destroy many buildings.



- Destructive storms occurring in the late summer and fall in the northern hemisphere's tropical region.
- Classified by their wind intensity at the eye wall.
- ▶ They can cause extreme levels of flooding and destroy many buildings.
- Monetary damages and loss of lives increase with an almost exponential character as a function of storm intensity.



► Forecasters predict two quantities: track and intensity

- ► Forecasters predict two quantities: track and intensity
- ► Two kinds of models: dynamical and statistical

- ► Forecasters predict two quantities: track and intensity
- ► Two kinds of models: dynamical and statistical
  - Dynamical: simulate the laws of physics

- ► Forecasters predict two quantities: track and intensity
- ► Two kinds of models: dynamical and statistical
  - Dynamical: simulate the laws of physics
  - Statistical: estimate based on historical data


- Forecasters predict two quantities: track and intensity
- ► Two kinds of models: dynamical and statistical
  - Dynamical: simulate the laws of physics
  - Statistical: estimate based on historical data
- Dynamical vs. statistical: good at long- and short-range forecasts respectively


- Forecasters predict two quantities: track and intensity
- Two kinds of models: dynamical and statistical
  - Dynamical: simulate the laws of physics
  - Statistical: estimate based on historical data
- Dynamical vs. statistical: good at long- and short-range forecasts respectively
- Rapid intensification: forecasted better by dynamical models

- Forecasters predict two quantities: track and intensity
- ► Two kinds of models: dynamical and statistical
  - Dynamical: simulate the laws of physics
  - Statistical: estimate based on historical data
- Dynamical vs. statistical: good at long- and short-range forecasts respectively
- Rapid intensification: forecasted better by dynamical models
- This project: a statistical model for intensity

# Hurricane forecasting basics: the SHIPS data

### Hurricane forecasting basics: our selection





# Model: limitations

Model: posterior predictive checking








# Forecasting

# Forecasting: model comparison



# Problems to solve & development ideas



### Additional information

#### R Markdown

This is an R Markdown presentation. Markdown is a simple formatting syntax for authoring HTML, PDF, and MS Word documents. For more details on using R Markdown see http://rmarkdown.rstudio.com.

When you click the **Knit** button a document will be generated that includes both content as well as the output of any embedded R code chunks within the document.

### Slide with Bullets

▶ Bullet 1

### Slide with Bullets

- ▶ Bullet 1
- ▶ Bullet 2

### Slide with Bullets

- ▶ Bullet 1
- ▶ Bullet 2
- ▶ Bullet 3

### Slide with R Output

### summary(cars)

```
##
       speed
                     dist
##
   Min. : 4.0
                Min. : 2.00
##
   1st Qu.:12.0
                1st Qu.: 26.00
##
   Median: 15.0 Median: 36.00
##
   Mean :15.4
                Mean : 42.98
##
   3rd Qu.:19.0
                3rd Qu.: 56.00
##
   Max. :25.0
                Max. :120.00
```

### Slide with Plot

