MolProbity Ramachandran analysis

7sd4.H.pdb, all models



94.4% (1265/1340) of all residues were in favored (98%) regions.
99.3% (1331/1340) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

- [1] 9 Asn (-170.5, -144.4)
- [1] 58 Gly (-159.9, -49.4)
- [2] 10 Asn (76.3, -141.9)
- [5] 9 Asn (-165.9, -86.5)
- [5] 26 Leu (76.1, 123.6)
- [6] 26 Leu (72.3, 119.1)
- [7] 9 Asn (-167.0, -86.9)
- [7] 26 Leu (77.3, 120.8)
- [9] 9 Asn (-173.0, -71.4)

http://kinemage.biochem.duke.edu

Lovell, Davis, et al. Proteins 50:437 (2003)

MolProbity Ramachandran analysis

7sd4.H.pdb, model 1



93.3% (125/134) of all residues were in favored (98%) regions.
98.5% (132/134) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[1] 9 Asn (-170.5, -144.4)
[1] 58 Gly (-159.9, -49.4)

http://kinemage.biochem.duke.edu

Lovell, Davis, et al. Proteins 50:437 (2003)

MolProbity Ramachandran analysis

7sd4.H.pdb, model 2



95.5% (128/134) of all residues were in favored (98%) regions.
99.3% (133/134) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[2] 10 Asn (76.3, -141.9)

MolProbity Ramachandran analysis

7sd4.H.pdb, model 3



94.8% (127/134) of all residues were in favored (98%) regions.
100.0% (134/134) of all residues were in allowed (>99.8%) regions.

There were no outliers.

http://kinemage.biochem.duke.edu

Lovell, Davis, et al. Proteins 50:437 (2003)

MolProbity Ramachandran analysis

7sd4.H.pdb, model 4



94.0% (126/134) of all residues were in favored (98%) regions.
100.0% (134/134) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

7sd4.H.pdb, model 5



94.0% (126/134) of all residues were in favored (98%) regions.
98.5% (132/134) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[5] 9 Asn (-165.9, -86.5)
[5] 26 Leu (76.1, 123.6)

MolProbity Ramachandran analysis

7sd4.H.pdb, model 6



94.0% (126/134) of all residues were in favored (98%) regions.
99.3% (133/134) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[6] 26 Leu (72.3, 119.1)

Lovell, Davis, et al. Proteins 50:437 (2003)

MolProbity Ramachandran analysis

7sd4.H.pdb, model 7



94.0% (126/134) of all residues were in favored (98%) regions.
98.5% (132/134) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[7] 9 Asn (-167.0, -86.9)
[7] 26 Leu (77.3, 120.8)

MolProbity Ramachandran analysis

7sd4.H.pdb, model 8



94.8% (127/134) of all residues were in favored (98%) regions.
100.0% (134/134) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

7sd4.H.pdb, model 9



94.0% (126/134) of all residues were in favored (98%) regions.
99.3% (133/134) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[9] 9 Asn (-173.0, -71.4)

MolProbity Ramachandran analysis

7sd4.H.pdb, model 10



95.5% (128/134) of all residues were in favored (98%) regions.
100.0% (134/134) of all residues were in allowed (>99.8%) regions.

There were no outliers.