




7p2o.H.pdb, all models









[14] 17 Lys (-15.8, -69.1)
[15] 17 Lys (-14.3, -70.4)
[15] 44 Ser (177.9, -158.4)

[1] 17 Lys (-22.4, -58.0)	[7] 23 Gly (134.5, 66.9)
[2] 17 Lys (-18.5, -66.3)	[8] 17 Lys (-16.7, -63.5)
[2] 45 Asn (-121.7, -52.5)	[10] 45 Asn (-116.0, -52.5)
[3] 17 Lys (-19.6, -61.4)	[11] 17 Lys (-17.7, -69.4)
[4] 17 Lys (-13.1, -76.6)	[12] 17 Lys (-14.2, -72.1)
[5] 17 Lys (-16.2, -74.6)	[12] 23 Gly (133.3, 51.0)
[6] 17 Lys (-27.0, -69.4)	[13] 17 Lys (-15.4, -69.0)
[7] 17 Lys (-16.2, -66.0)	[13] 23 Gly (133.4, 62.7)

[15] 4 Ser (-17.9, -155.4)
[16] 17 Lys (-18.1, -60.4)
[16] 23 Gly (132.3, 59.9)
[17] 17 Lys (-16.8, -61.5)
[17] 23 Gly (131.1, 59.1)
[18] 17 Lys (-11.1, -75.9)
[19] 17 Lys (-18.8, -64.7)
[20] 17 Lys (-14.8, -75.7)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 1









89.1% (57/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[1] 17 Lys (-22.4, -58.0)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 2









81.2% (52/64) of all residues were in favored (98%) regions.
96.9% (62/64) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[2] 17 Lys (-18.5, -66.3)
[2] 45 Asn (-121.7, -52.5)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 3









87.5% (56/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[3] 17 Lys (-19.6, -61.4)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 4









87.5% (56/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[4] 17 Lys (-13.1, -76.6)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 5









90.6% (58/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[5] 17 Lys (-16.2, -74.6)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 6









85.9% (55/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[6] 17 Lys (-27.0, -69.4)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 7








85.9% (55/64) of all residues were in favored (98%) regions.
96.9% (62/64) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[7] 17 Lys (-16.2, -66.0)
[7] 23 Gly (134.5, 66.9)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 8









85.9% (55/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[8] 17 Lys (-16.7, -63.5)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 9









82.8% (53/64) of all residues were in favored (98%) regions.
100.0% (64/64) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

7p2o.H.pdb, model 10









81.2% (52/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[10] 45 Asn (-116.0, -52.5)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 11









87.5% (56/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[11] 17 Lys (-17.7, -69.4)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 12









87.5% (56/64) of all residues were in favored (98%) regions.
96.9% (62/64) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[12] 17 Lys (-14.2, -72.1)
[12] 23 Gly (133.3, 51.0)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 13









89.1% (57/64) of all residues were in favored (98%) regions.
96.9% (62/64) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[13] 17 Lys (-15.4, -69.0)
[13] 23 Gly (133.4, 62.7)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 14




87.5% (56/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[14] 17 Lys (-15.8, -69.1)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 15








87.5% (56/64) of all residues were in favored (98%) regions.
96.9% (62/64) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[15] 17 Lys (-14.3, -70.4)
[15] 44 Ser (177.9, -158.4)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 16









85.9% (55/64) of all residues were in favored (98%) regions.
96.9% (62/64) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[16] 17 Lys (-18.1, -60.4)
[16] 23 Gly (132.3, 59.9)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 17









89.1% (57/64) of all residues were in favored (98%) regions.
96.9% (62/64) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[17] 17 Lys (-16.8, -61.5)
[17] 23 Gly (131.1, 59.1)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 18









90.6% (58/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[18] 17 Lys (-11.1, -75.9)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 19









89.1% (57/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[19] 17 Lys (-18.8, -64.7)

MolProbity Ramachandran analysis

7p2o.H.pdb, model 20



87.5% (56/64) of all residues were in favored (98%) regions.
98.4% (63/64) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[20] 17 Lys (-14.8, -75.7)