


MolProbity Ramachandran analysis

8dwa.H.pdb, model 1



84.8% (502/592) of all residues were in favored (98%) regions.
96.1% (569/592) of all residues were in allowed (>99.8%) regions.

There were 23 outliers (phi, psi):

A 337 Pro (-69.6, -80.7)	H 14 Pro (-44.8, 163.1)	L 68 Gly (34.0, -89.5)
A 338 Phe (58.4, -59.6)	H 17 Ser (173.0, 136.7)	L 92 Tyr (87.4, -160.0)
A 361 Cys (161.3, 171.1)	H 29 Phe (37.5, -84.2)	L 94 Asn (54.5, 134.9)
A 365 Tyr (78.3, -112.8)	H 30 Gly (-28.0, -50.2)	L 95 Leu (84.7, 132.2)
A 366 Ser (42.0, -81.1)	H 43 Gln (-164.9, -102.2)	L 142 Arg (-61.7, -93.2)
A 372 Ala (-85.4, -132.2)	H 99 Thr (-61.1, -144.1)	L 143 Glu (10.1, 76.9)
A 448 Asn (-20.0, 107.2)	H 158 Ala (54.5, -89.5)	L 168 Ser (-27.4, -42.9)
A 487 Asn (45.2, -153.9)	H 191 Thr (94.5, -100.2)	