







## 7mhl.H.pdb, all models



# MolProbity Ramachandran analysis

7mhl.H.pdb, model 1









92.4% (281/304) of all residues were in favored (98%) regions.  
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [1] 3 Phe (40.8, 114.0)
- [1] 154 Tyr (-21.1, 71.9)
- [1] 155 Asp (160.1, -83.9)
- [1] 277 Asn (54.3, -1.2)
- [1] 300 Cys (-63.9, -156.7)
- [1] 303 Val (-154.9, 56.6)
- [1] 305 Phe (158.9, -3.8)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 2




90.5% (275/304) of all residues were in favored (98%) regions.  
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

|                            |                             |
|----------------------------|-----------------------------|
| [2] 3 Phe (51.1, 115.2)    | [2] 303 Val (-171.1, 85.7)  |
| [2] 48 Asp (-171.0, -28.9) | [2] 305 Phe (153.3, -109.1) |
| [2] 50 Leu (-58.4, -104.8) |                             |
| [2] 155 Asp (145.4, -21.1) |                             |
| [2] 277 Asn (108.0, -46.9) |                             |
| [2] 300 Cys (76.0, -131.7) |                             |
| [2] 301 Ser (-33.7, 81.1)  |                             |
| [2] 302 Gly (62.6, -27.8)  |                             |

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 3




91.8% (279/304) of all residues were in favored (98%) regions.  
96.4% (293/304) of all residues were in allowed (>99.8%) regions.

There were 11 outliers (phi, psi):

|                             |                             |
|-----------------------------|-----------------------------|
| [3] 2 Gly (163.2, 7.5)      | [3] 302 Gly (32.2, -110.6)  |
| [3] 3 Phe (36.4, 104.4)     | [3] 304 Thr (169.6, -173.6) |
| [3] 49 Met (-34.7, -31.5)   | [3] 305 Phe (142.3, 40.3)   |
| [3] 72 Asn (88.6, -63.0)    |                             |
| [3] 198 Thr (-25.7, 141.8)  |                             |
| [3] 299 Gln (-67.6, -148.7) |                             |
| [3] 300 Cys (10.5, -129.6)  |                             |
| [3] 301 Ser (16.2, 83.6)    |                             |

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 4









91.4% (278/304) of all residues were in favored (98%) regions.  
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [4] 3 Phe (37.6, 99.6)
- [4] 154 Tyr (-2.7, -57.2)
- [4] 300 Cys (40.5, -92.1)
- [4] 304 Thr (172.5, -131.4)
- [4] 305 Phe (170.8, -162.5)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 5




92.8% (282/304) of all residues were in favored (98%) regions.  
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[5] 3 Phe (86.0, 121.3)  
[5] 71 Gly (30.7, 81.9)  
[5] 143 Gly (-58.9, 18.5)  
[5] 300 Cys (24.9, -117.2)  
[5] 301 Ser (42.0, -4.1)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 6









91.4% (278/304) of all residues were in favored (98%) regions.  
96.1% (292/304) of all residues were in allowed (>99.8%) regions.

There were 12 outliers (phi, psi):

|                             |                             |
|-----------------------------|-----------------------------|
| [6] 2 Gly (36.1, -96.9)     | [6] 300 Cys (-24.0, -94.4)  |
| [6] 3 Phe (134.4, 129.3)    | [6] 302 Gly (-47.5, 68.1)   |
| [6] 24 Thr (-31.9, -78.7)   | [6] 303 Val (-58.0, 65.2)   |
| [6] 48 Asp (-30.8, -33.9)   | [6] 304 Thr (128.6, -121.6) |
| [6] 72 Asn (61.6, -15.9)    |                             |
| [6] 155 Asp (170.5, -5.2)   |                             |
| [6] 192 Gln (63.3, 110.7)   |                             |
| [6] 299 Gln (-56.5, -100.6) |                             |

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 7



93.1% (283/304) of all residues were in favored (98%) regions.  
96.7% (294/304) of all residues were in allowed (>99.8%) regions.






There were 10 outliers (phi, psi):

|                            |                           |
|----------------------------|---------------------------|
| [7] 2 Gly (-2.1, -105.9)   | [7] 303 Val (-6.7, 106.4) |
| [7] 3 Phe (170.4, 131.4)   | [7] 304 Thr (53.6, -94.9) |
| [7] 71 Gly (35.4, -105.8)  |                           |
| [7] 189 Gln (-65.8, 63.2)  |                           |
| [7] 191 Ala (-56.6, -88.8) |                           |
| [7] 192 Gln (63.2, 132.5)  |                           |
| [7] 277 Asn (44.3, 19.4)   |                           |
| [7] 301 Ser (-68.7, 32.8)  |                           |



# MolProbity Ramachandran analysis

7mhl.H.pdb, model 8




92.8% (282/304) of all residues were in favored (98%) regions.  
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [8] 154 Tyr (28.5, -41.8)
- [8] 278 Gly (39.8, 16.7)
- [8] 301 Ser (-178.4, 82.9)
- [8] 304 Thr (55.8, -75.6)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 9









91.1% (277/304) of all residues were in favored (98%) regions.  
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[9] 49 Met (154.7, 4.2)  
[9] 119 Asn (16.0, 73.9)  
[9] 142 Asn (-32.1, 145.8)  
[9] 154 Tyr (116.2, -142.9)  
[9] 301 Ser (169.7, -4.5)  
[9] 303 Val (-161.1, -158.3)  
[9] 304 Thr (6.5, -92.5)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 10




92.8% (282/304) of all residues were in favored (98%) regions.  
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [10] 23 Gly (45.7, -97.0)
- [10] 46 Ser (-34.8, -92.4)
- [10] 47 Glu (30.5, -68.5)
- [10] 49 Met (15.7, 100.9)
- [10] 50 Leu (-172.1, 47.5)
- [10] 303 Val (-42.6, -82.4)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 11









90.8% (276/304) of all residues were in favored (98%) regions.  
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

|                              |                              |
|------------------------------|------------------------------|
| [11] 23 Gly (33.1, -164.1)   | [11] 303 Val (-80.3, -90.7)  |
| [11] 24 Thr (23.1, -59.0)    | [11] 304 Thr (-52.2, -123.6) |
| [11] 46 Ser (17.6, -69.0)    |                              |
| [11] 48 Asp (-72.5, -112.4)  |                              |
| [11] 73 Val (-37.6, 106.1)   |                              |
| [11] 154 Tyr (-74.6, -135.7) |                              |
| [11] 300 Cys (176.8, 57.8)   |                              |
| [11] 301 Ser (152.7, 95.8)   |                              |

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 12









95.1% (289/304) of all residues were in favored (98%) regions.  
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [12] 46 Ser (-28.3, -75.9)
- [12] 47 Glu (-7.4, -61.5)
- [12] 48 Asp (-56.8, -146.0)
- [12] 49 Met (39.0, -33.3)
- [12] 154 Tyr (88.4, -91.3)
- [12] 191 Ala (-36.6, 100.9)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 13









93.4% (284/304) of all residues were in favored (98%) regions.  
99.0% (301/304) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):  
[13] 154 Tyr (-3.2, -54.7)  
[13] 278 Gly (163.3, -4.2)  
[13] 303 Val (-129.9, -129.7)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 14









94.4% (287/304) of all residues were in favored (98%) regions.  
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[14] 154 Tyr (-20.3, -45.4)  
[14] 302 Gly (32.6, -102.6)  
[14] 303 Val (154.5, -51.9)  
[14] 305 Phe (1.6, -101.7)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 15



91.1% (277/304) of all residues were in favored (98%) regions.  
97.7% (297/304) of all residues were in allowed (>99.8%) regions.






There were 7 outliers (phi, psi):

[15] 46 Ser (-46.5, -16.0)  
[15] 51 Asn (160.4, 79.2)  
[15] 71 Gly (53.8, -100.3)  
[15] 154 Tyr (119.8, -27.9)  
[15] 277 Asn (35.9, 77.5)  
[15] 301 Ser (-18.7, 127.7)  
[15] 303 Val (110.9, 2.0)



# MolProbity Ramachandran analysis

7mhl.H.pdb, model 16




93.4% (284/304) of all residues were in favored (98%) regions.  
99.3% (302/304) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):  
[16] 301 Ser (43.9, 131.6)  
[16] 303 Val (-167.3, -57.7)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 17




92.8% (282/304) of all residues were in favored (98%) regions.  
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[17] 47 Glu (-66.5, 58.2)  
[17] 48 Asp (172.4, 51.4)  
[17] 301 Ser (59.6, 130.6)  
[17] 303 Val (-170.7, -49.0)  
[17] 305 Phe (-76.6, -125.4)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 18









94.1% (286/304) of all residues were in favored (98%) regions.  
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [18] 46 Ser (57.0, -76.5)
- [18] 48 Asp (153.8, 40.5)
- [18] 51 Asn (-155.4, 35.8)
- [18] 73 Val (-33.4, 84.1)
- [18] 154 Tyr (-15.9, -45.3)
- [18] 156 Cys (176.3, 116.9)
- [18] 301 Ser (151.4, -24.8)
- [18] 302 Gly (-6.1, -129.5)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 19




93.8% (285/304) of all residues were in favored (98%) regions.  
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[19] 46 Ser (-66.3, -102.3)  
[19] 47 Glu (0.8, -49.5)  
[19] 71 Gly (44.6, -104.6)  
[19] 154 Tyr (21.4, -90.2)  
[19] 301 Ser (18.7, 92.3)  
[19] 302 Gly (-131.9, -66.7)  
[19] 303 Val (51.0, 12.6)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 20









93.1% (283/304) of all residues were in favored (98%) regions.  
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

[20] 46 Ser (-18.6, -37.1)  
[20] 154 Tyr (17.0, -85.7)  
[20] 213 Ile (-69.1, 21.2)  
[20] 277 Asn (22.9, 55.5)  
[20] 301 Ser (52.0, 165.7)  
[20] 304 Thr (176.1, -87.7)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 21









91.8% (279/304) of all residues were in favored (98%) regions.  
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[21] 71 Gly (50.5, -83.5)  
[21] 154 Tyr (49.1, -94.7)  
[21] 303 Val (81.9, 136.3)  
[21] 305 Phe (94.6, -56.9)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 22









93.8% (285/304) of all residues were in favored (98%) regions.  
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [22] 154 Tyr (14.2, -78.1)
- [22] 276 Met (-57.7, 45.4)
- [22] 277 Asn (-10.2, 113.3)
- [22] 278 Gly (60.7, -22.7)
- [22] 301 Ser (22.5, -160.7)
- [22] 303 Val (53.2, -176.4)
- [22] 304 Thr (-13.6, 90.4)
- [22] 305 Phe (0.9, -61.9)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 23



92.1% (280/304) of all residues were in favored (98%) regions.  
98.7% (300/304) of all residues were in allowed (>99.8%) regions.







There were 4 outliers (phi, psi):

[23] 47 Glu (-58.9, 2.9)  
[23] 169 Thr (-60.7, 8.6)  
[23] 193 Ala (11.4, 91.2)  
[23] 303 Val (75.2, -97.1)



# MolProbity Ramachandran analysis

7mhl.H.pdb, model 24









92.1% (280/304) of all residues were in favored (98%) regions.  
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

|                              |                            |
|------------------------------|----------------------------|
| [24] 23 Gly (37.9, -97.8)    | [24] 305 Phe (74.8, -79.9) |
| [24] 46 Ser (-27.3, -92.4)   |                            |
| [24] 47 Glu (-18.2, -43.3)   |                            |
| [24] 51 Asn (-134.9, 23.3)   |                            |
| [24] 142 Asn (14.3, 92.4)    |                            |
| [24] 277 Asn (-32.8, -127.8) |                            |
| [24] 301 Ser (14.3, -130.6)  |                            |
| [24] 302 Gly (52.7, 91.0)    |                            |

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 25









90.5% (275/304) of all residues were in favored (98%) regions.  
95.1% (289/304) of all residues were in allowed (>99.8%) regions.

There were 15 outliers (phi, psi):

|                              |                              |
|------------------------------|------------------------------|
| [25] 24 Thr (-18.5, -84.7)   | [25] 192 Gln (-50.1, 169.7)  |
| [25] 47 Glu (-30.8, -114.5)  | [25] 193 Ala (166.9, 150.2)  |
| [25] 48 Asp (-31.6, -30.3)   | [25] 277 Asn (50.0, -70.6)   |
| [25] 49 Met (-39.9, -17.1)   | [25] 300 Cys (-67.1, -86.2)  |
| [25] 52 Pro (-111.5, 164.3)  | [25] 302 Gly (178.9, 91.3)   |
| [25] 154 Tyr (47.2, -95.3)   | [25] 303 Val (-157.2, -76.2) |
| [25] 189 Gln (-37.9, 173.9)  | [25] 304 Thr (-79.6, -93.6)  |
| [25] 190 Thr (139.6, -179.4) |                              |

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 26









93.1% (283/304) of all residues were in favored (98%) regions.  
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[26] 23 Gly (-8.9, -56.0)  
[26] 46 Ser (-3.1, -57.2)  
[26] 71 Gly (14.8, -77.4)  
[26] 154 Tyr (-4.7, -39.1)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 27









92.4% (281/304) of all residues were in favored (98%) regions.  
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [27] 23 Gly (51.6, -78.5)
- [27] 24 Thr (-78.4, -96.6)
- [27] 155 Asp (-174.9, -10.2)
- [27] 301 Ser (-48.6, -137.2)
- [27] 303 Val (-154.8, 1.9)
- [27] 304 Thr (153.4, 12.5)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 28









91.8% (279/304) of all residues were in favored (98%) regions.  
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

|                              |                             |
|------------------------------|-----------------------------|
| [28] 73 Val (-21.9, 123.1)   | [28] 303 Val (-152.3, 84.4) |
| [28] 154 Tyr (92.4, -36.0)   | [28] 305 Phe (157.6, 87.4)  |
| [28] 167 Leu (-81.6, -161.3) |                             |
| [28] 168 Pro (-87.6, -35.8)  |                             |
| [28] 278 Gly (160.0, 6.9)    |                             |
| [28] 298 Arg (-54.6, -79.4)  |                             |
| [28] 300 Cys (-52.9, -98.0)  |                             |
| [28] 301 Ser (-59.3, -120.3) |                             |

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 29




93.1% (283/304) of all residues were in favored (98%) regions.  
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

[29] 24 Thr (-33.0, -12.8)  
[29] 52 Pro (-121.5, 148.5)  
[29] 72 Asn (-51.4, -73.3)  
[29] 73 Val (21.8, 58.3)  
[29] 154 Tyr (34.3, -47.1)  
[29] 301 Ser (-167.7, -104.9)  
[29] 303 Val (174.0, 67.5)  
[29] 305 Phe (161.6, -14.5)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 30









94.1% (286/304) of all residues were in favored (98%) regions.  
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[30] 154 Tyr (172.1, -79.1)  
[30] 169 Thr (-41.5, -4.7)  
[30] 277 Asn (21.1, -70.6)  
[30] 301 Ser (-162.5, -89.5)  
[30] 305 Phe (-175.5, -11.7)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 31



94.7% (288/304) of all residues were in favored (98%) regions.  
98.4% (299/304) of all residues were in allowed (>99.8%) regions.


There were 5 outliers (phi, psi):

[31] 153 Asp (-53.0, -128.4)  
[31] 301 Ser (-49.6, -142.4)  
[31] 302 Gly (73.0, 97.1)  
[31] 304 Thr (92.6, -111.0)  
[31] 305 Phe (-18.3, 137.2)



# MolProbity Ramachandran analysis

7mhl.H.pdb, model 32









92.8% (282/304) of all residues were in favored (98%) regions.  
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[32] 46 Ser (26.8, -78.9)  
[32] 47 Glu (-53.8, 11.3)  
[32] 48 Asp (-172.1, 72.3)  
[32] 193 Ala (-37.3, 146.6)  
[32] 302 Gly (-179.9, 69.6)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 33









92.1% (280/304) of all residues were in favored (98%) regions.  
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [33] 47 Glu (-40.1, -26.4)
- [33] 48 Asp (175.2, 60.5)
- [33] 49 Met (-44.8, -4.4)
- [33] 142 Asn (-16.1, 121.7)
- [33] 154 Tyr (16.4, -69.9)
- [33] 191 Ala (-20.8, 133.8)
- [33] 301 Ser (-21.3, 140.5)
- [33] 303 Val (77.8, -8.4)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 34









92.4% (281/304) of all residues were in favored (98%) regions.  
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[34] 51 Asn (-147.8, 25.5)  
[34] 191 Ala (-37.0, -86.5)  
[34] 192 Gln (28.7, 110.1)  
[34] 299 Gln (-44.3, -19.4)  
[34] 304 Thr (-163.1, 6.6)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 35









94.1% (286/304) of all residues were in favored (98%) regions.  
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[35] 49 Met (-11.2, -29.0)  
[35] 154 Tyr (33.5, -99.4)  
[35] 299 Gln (-25.0, 53.8)  
[35] 303 Val (86.1, 37.0)  
[35] 304 Thr (-55.8, -0.9)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 36









94.4% (287/304) of all residues were in favored (98%) regions.  
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[36] 71 Gly (52.9, -68.4)  
[36] 154 Tyr (-3.4, -64.4)  
[36] 191 Ala (-42.1, 93.8)  
[36] 305 Phe (-168.3, -142.4)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 37




94.1% (286/304) of all residues were in favored (98%) regions.  
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [37] 49 Met (-47.4, -3.9)
- [37] 142 Asn (-9.9, 121.2)
- [37] 167 Leu (-84.6, -161.3)
- [37] 168 Pro (-86.0, -40.3)
- [37] 195 Gly (-57.6, -162.0)
- [37] 301 Ser (-4.0, 123.1)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 38









92.8% (282/304) of all residues were in favored (98%) regions.  
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[38] 141 Leu (-64.9, -169.0)  
[38] 142 Asn (-32.7, -174.0)  
[38] 143 Gly (20.4, 16.2)  
[38] 154 Tyr (-21.7, -34.4)  
[38] 193 Ala (178.7, 107.9)  
[38] 195 Gly (-51.7, -132.7)  
[38] 301 Ser (38.5, -156.7)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 39



94.4% (287/304) of all residues were in favored (98%) regions.  
97.7% (297/304) of all residues were in allowed (>99.8%) regions.







There were 7 outliers (phi, psi):

[39] 46 Ser (13.2, -70.3)  
[39] 154 Tyr (-2.1, -89.9)  
[39] 193 Ala (162.0, 86.0)  
[39] 194 Ala (-18.1, -104.3)  
[39] 299 Gln (-55.0, 25.6)  
[39] 301 Ser (115.0, -145.6)  
[39] 303 Val (163.2, 65.4)



# MolProbity Ramachandran analysis

7mhl.H.pdb, model 40









92.8% (282/304) of all residues were in favored (98%) regions.  
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

|                             |                              |
|-----------------------------|------------------------------|
| [40] 23 Gly (52.8, -101.9)  | [40] 305 Phe (-174.5, -33.1) |
| [40] 73 Val (-29.2, 113.4)  |                              |
| [40] 154 Tyr (6.1, -86.8)   |                              |
| [40] 189 Gln (-53.4, 76.3)  |                              |
| [40] 194 Ala (-56.9, -80.4) |                              |
| [40] 277 Asn (33.5, 79.5)   |                              |
| [40] 303 Val (-153.3, 23.8) |                              |
| [40] 304 Thr (-167.8, 43.8) |                              |

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 41









92.8% (282/304) of all residues were in favored (98%) regions.  
99.0% (301/304) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[41] 49 Met (-70.1, 23.5)  
[41] 154 Tyr (58.6, -82.4)  
[41] 303 Val (-50.5, -87.4)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 42









95.7% (291/304) of all residues were in favored (98%) regions.  
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[42] 23 Gly (51.0, -74.0)  
[42] 24 Thr (-79.5, -96.5)  
[42] 49 Met (-62.3, 8.6)  
[42] 277 Asn (55.0, 91.7)  
[42] 278 Gly (38.8, 26.9)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 43









94.1% (286/304) of all residues were in favored (98%) regions.  
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[43] 142 Asn (-58.8, 99.1)  
[43] 154 Tyr (41.9, -90.7)  
[43] 300 Cys (-62.2, 57.4)  
[43] 301 Ser (-176.2, -84.2)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 44




93.4% (284/304) of all residues were in favored (98%) regions.  
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[44] 48 Asp (144.2, -23.0)  
[44] 52 Pro (-120.4, 135.9)  
[44] 303 Val (-93.9, -101.7)  
[44] 305 Phe (161.8, -24.1)

# MolProbity Ramachandran analysis

7mhl.H.pdb, model 45



91.8% (279/304) of all residues were in favored (98%) regions.  
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

|                             |                              |
|-----------------------------|------------------------------|
| [45] 47 Glu (-22.5, 72.5)   | [45] 300 Cys (-70.3, 19.5)   |
| [45] 48 Asp (169.7, -26.6)  | [45] 302 Gly (-156.9, -89.2) |
| [45] 50 Leu (-24.0, -65.9)  |                              |
| [45] 168 Pro (-108.3, 64.9) |                              |
| [45] 169 Thr (-170.5, -3.9) |                              |
| [45] 191 Ala (-59.7, 92.4)  |                              |
| [45] 277 Asn (56.9, -67.3)  |                              |
| [45] 278 Gly (179.7, 13.0)  |                              |