


MolProbity Ramachandran analysis

7mhm.H.pdb, all models



93.9% (12281/13072) of all residues were in favored (98%) regions.
98.0% (12805/13072) of all residues were in allowed (>99.8%) regions.
This list is truncated; use the MolProbity multi-chart.html for complete list.







There were 267 outliers (phi, psi):

[1] 51 Asn (-166.1, 42.9)	[2] 301 Ser (3.6, -92.3)	[4] 24 Thr (41.9, 11.1)
[1] 154 Tyr (51.0, -69.1)	[2] 303 Val (125.6, -159.9)	[4] 154 Tyr (60.2, -91.2)
[1] 303 Val (67.8, -179.9)	[2] 305 Phe (121.7, -32.9)	[4] 303 Val (140.9, -56.1)
[1] 305 Phe (-173.4, 49.0)	[3] 302 Gly (-58.4, -145.0)	[4] 304 Thr (-173.6, -148.7)
[2] 23 Gly (-25.1, 120.6)	[3] 303 Val (114.6, -97.0)	[5] 52 Pro (-106.8, 129.3)
[2] 24 Thr (102.8, -25.5)	[3] 304 Thr (169.2, -71.3)	[5] 71 Gly (45.7, -57.4)
[2] 71 Gly (57.7, -84.0)	[3] 305 Phe (171.2, -76.6)	[5] 72 Asn (-165.0, 11.3)
[2] 154 Tyr (40.3, -99.3)	[4] 23 Gly (24.8, 112.3)	[5] 154 Tyr (52.4, -89.7)
		[5] 301 Ser (-47.8, 73.8)
		[5] 302 Gly (-160.9, -56.6)
		[5] 303 Val (97.2, -16.1)

[5] 305 Phe (-39.6, -86.6)	[8] 154 Tyr (39.2, -90.2)	[10] 277 Asn (30.1, 38.9)
[6] 154 Tyr (31.0, -68.7)	[8] 277 Asn (34.1, 29.0)	[10] 303 Val (90.9, 72.7)
[6] 301 Ser (-10.7, 63.5)	[8] 301 Ser (131.8, -40.5)	[10] 304 Thr (27.7, -173.5)
[6] 302 Gly (150.7, 26.6)	[8] 303 Val (56.6, -46.2)	[11] 154 Tyr (67.7, -98.9)
[6] 303 Val (6.8, -62.3)	[9] 154 Tyr (41.2, -90.4)	[11] 191 Ala (-45.5, 95.5)
[7] 71 Gly (50.3, -70.1)	[9] 155 Asp (-71.3, -129.9)	[11] 303 Val (90.7, 78.9)
[7] 154 Tyr (74.4, -120.7)	[9] 156 Cys (42.2, 97.0)	[11] 304 Thr (30.9, -179.4)
[7] 278 Gly (156.4, 6.0)	[9] 300 Cys (132.3, -25.7)	[12] 71 Gly (-18.5, 100.0)
[7] 301 Ser (110.8, -57.4)	[9] 304 Thr (-175.3, -73.7)	[12] 72 Asn (116.7, -58.6)
[7] 303 Val (134.0, -78.1)	[9] 305 Phe (169.2, -89.3)	[12] 305 Phe (-60.7, 72.7)
[7] 304 Thr (-71.9, -133.2)	[10] 72 Asn (-51.0, -78.6)	[13] 71 Gly (3.7, 109.5)
[8] 46 Ser (-45.5, -16.1)	[10] 195 Gly (-53.0, -165.7)	[13] 303 Val (80.3, -19.8)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 1









95.4% (290/304) of all residues were in favored (98%) regions.
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [1] 51 Asn (-166.1, 42.9)
- [1] 154 Tyr (51.0, -69.1)
- [1] 303 Val (67.8, -179.9)
- [1] 305 Phe (-173.4, 49.0)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 2




95.1% (289/304) of all residues were in favored (98%) regions.
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [2] 23 Gly (-25.1, 120.6)
- [2] 24 Thr (102.8, -25.5)
- [2] 71 Gly (57.7, -84.0)
- [2] 154 Tyr (40.3, -99.3)
- [2] 301 Ser (3.6, -92.3)
- [2] 303 Val (125.6, -159.9)
- [2] 305 Phe (121.7, -32.9)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 3









95.4% (290/304) of all residues were in favored (98%) regions.
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[3] 302 Gly (-58.4, -145.0)
[3] 303 Val (114.6, -97.0)
[3] 304 Thr (169.2, -71.3)
[3] 305 Phe (171.2, -76.6)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 4




95.7% (291/304) of all residues were in favored (98%) regions.
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [4] 23 Gly (24.8, 112.3)
- [4] 24 Thr (41.9, 11.1)
- [4] 154 Tyr (60.2, -91.2)
- [4] 303 Val (140.9, -56.1)
- [4] 304 Thr (-173.6, -148.7)
- [4] 305 Phe (-39.3, -91.0)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 5









93.8% (285/304) of all residues were in favored (98%) regions.
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [5] 52 Pro (-106.8, 129.3)
- [5] 71 Gly (45.7, -57.4)
- [5] 72 Asn (-165.0, 11.3)
- [5] 154 Tyr (52.4, -89.7)
- [5] 301 Ser (-47.8, 73.8)
- [5] 302 Gly (-160.9, -56.6)
- [5] 303 Val (97.2, -16.1)
- [5] 305 Phe (-39.6, -86.6)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 6









94.4% (287/304) of all residues were in favored (98%) regions.
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[6] 154 Tyr (31.0, -68.7)
[6] 301 Ser (-10.7, 63.5)
[6] 302 Gly (150.7, 26.6)
[6] 303 Val (6.8, -62.3)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 7








93.8% (285/304) of all residues were in favored (98%) regions.
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [7] 71 Gly (50.3, -70.1)
- [7] 154 Tyr (74.4, -120.7)
- [7] 278 Gly (156.4, 6.0)
- [7] 301 Ser (110.8, -57.4)
- [7] 303 Val (134.0, -78.1)
- [7] 304 Thr (-71.9, -133.2)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 8




94.4% (287/304) of all residues were in favored (98%) regions.
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[8] 46 Ser (-45.5, -16.1)
[8] 154 Tyr (39.2, -90.2)
[8] 277 Asn (34.1, 29.0)
[8] 301 Ser (131.8, -40.5)
[8] 303 Val (56.6, -46.2)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 9









95.4% (290/304) of all residues were in favored (98%) regions.
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [9] 154 Tyr (41.2, -90.4)
- [9] 155 Asp (-71.3, -129.9)
- [9] 156 Cys (42.2, 97.0)
- [9] 300 Cys (132.3, -25.7)
- [9] 304 Thr (-175.3, -73.7)
- [9] 305 Phe (169.2, -89.3)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 10




94.7% (288/304) of all residues were in favored (98%) regions.
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[10] 72 Asn (-51.0, -78.6)
[10] 195 Gly (-53.0, -165.7)
[10] 277 Asn (30.1, 38.9)
[10] 303 Val (90.9, 72.7)
[10] 304 Thr (27.7, -173.5)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 11









94.4% (287/304) of all residues were in favored (98%) regions.
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[11] 154 Tyr (67.7, -98.9)
[11] 191 Ala (-45.5, 95.5)
[11] 303 Val (90.7, 78.9)
[11] 304 Thr (30.9, -179.4)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 12



93.4% (284/304) of all residues were in favored (98%) regions.
99.0% (301/304) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):







[12] 71 Gly (-18.5, 100.0)

[12] 72 Asn (116.7, -58.6)

[12] 305 Phe (-60.7, 72.7)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 13




95.4% (290/304) of all residues were in favored (98%) regions.
99.0% (301/304) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[13] 71 Gly (3.7, 109.5)
[13] 303 Val (80.3, -19.8)
[13] 305 Phe (113.7, 35.2)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 14




92.1% (280/304) of all residues were in favored (98%) regions.
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[14] 45 Thr (-81.5, -116.7)	[14] 305 Phe (-13.2, 132.4)
[14] 59 Ile (-31.8, -74.6)	
[14] 72 Asn (176.7, -35.8)	
[14] 154 Tyr (-58.3, -96.9)	
[14] 191 Ala (-46.3, 101.4)	
[14] 301 Ser (-60.5, 82.0)	
[14] 303 Val (106.6, 112.8)	
[14] 304 Thr (161.2, 52.3)	

MolProbity Ramachandran analysis

7mhm.H.pdb, model 15








93.4% (284/304) of all residues were in favored (98%) regions.
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [15] 47 Glu (-22.2, 103.1)
- [15] 48 Asp (167.7, -9.1)
- [15] 73 Val (-178.0, 136.7)
- [15] 154 Tyr (33.5, -86.1)
- [15] 301 Ser (5.1, 48.0)
- [15] 303 Val (62.5, -1.6)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 16









94.4% (287/304) of all residues were in favored (98%) regions.
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[16] 48 Asp (71.3, 138.8)
[16] 49 Met (116.5, -8.4)
[16] 154 Tyr (20.9, -76.6)
[16] 302 Gly (28.6, -163.5)
[16] 303 Val (86.8, -74.8)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 17









94.1% (286/304) of all residues were in favored (98%) regions.
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[17] 23 Gly (33.2, -112.5)	[17] 305 Phe (54.9, 103.2)
[17] 49 Met (164.6, 22.3)	
[17] 50 Leu (165.3, 31.4)	
[17] 52 Pro (-111.7, 132.4)	
[17] 142 Asn (-2.6, 88.4)	
[17] 301 Ser (44.6, -72.9)	
[17] 303 Val (154.3, -55.0)	
[17] 304 Thr (-67.0, 35.5)	

MolProbity Ramachandran analysis

7mhm.H.pdb, model 18




93.1% (283/304) of all residues were in favored (98%) regions.
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [18] 49 Met (102.8, -8.2)
- [18] 52 Pro (-162.5, 140.5)
- [18] 71 Gly (52.2, -63.5)
- [18] 154 Tyr (70.7, -100.6)
- [18] 234 Ala (-26.1, -48.1)
- [18] 303 Val (66.2, -17.3)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 19




92.8% (282/304) of all residues were in favored (98%) regions.
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[19] 46 Ser (61.2, -90.1)	[19] 303 Val (101.0, -39.9)
[19] 48 Asp (153.2, -169.4)	[19] 305 Phe (94.5, -29.2)
[19] 49 Met (61.6, 99.0)	
[19] 50 Leu (145.5, -84.0)	
[19] 52 Pro (-165.6, 154.1)	
[19] 142 Asn (-32.9, 100.7)	
[19] 154 Tyr (85.5, -73.4)	
[19] 301 Ser (150.6, 31.9)	

MolProbity Ramachandran analysis

7mhm.H.pdb, model 20









92.8% (282/304) of all residues were in favored (98%) regions.
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[20] 3 Phe (169.2, 119.6)	[20] 303 Val (84.9, -1.7)
[20] 46 Ser (34.0, -99.3)	[20] 305 Phe (83.2, 49.3)
[20] 48 Asp (172.8, -163.7)	
[20] 49 Met (94.3, 104.6)	
[20] 50 Leu (153.2, -16.2)	
[20] 51 Asn (136.3, 133.4)	
[20] 52 Pro (-131.8, -149.1)	
[20] 154 Tyr (31.6, -79.6)	

MolProbity Ramachandran analysis

7mhm.H.pdb, model 21




94.1% (286/304) of all residues were in favored (98%) regions.
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

[21] 45 Thr (-55.0, -166.6)
[21] 51 Asn (-91.1, 69.9)
[21] 154 Tyr (73.3, -101.2)
[21] 302 Gly (50.7, 135.6)
[21] 304 Thr (131.1, -50.0)
[21] 305 Phe (99.4, 49.3)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 22




92.8% (282/304) of all residues were in favored (98%) regions.
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [22] 45 Thr (-51.8, -167.4)
- [22] 49 Met (2.5, 105.5)
- [22] 51 Asn (-98.5, 72.5)
- [22] 71 Gly (38.5, -95.2)
- [22] 305 Phe (132.7, 103.2)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 23




93.8% (285/304) of all residues were in favored (98%) regions.
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[23] 46 Ser (-5.7, -143.6)	[23] 195 Gly (-50.6, -171.1)
[23] 48 Asp (165.2, -11.2)	
[23] 49 Met (-44.9, 162.2)	
[23] 50 Leu (-152.5, -110.1)	
[23] 51 Asn (-74.2, 81.4)	
[23] 71 Gly (46.7, -84.3)	
[23] 155 Asp (112.7, -3.2)	
[23] 191 Ala (-49.0, 98.6)	

MolProbity Ramachandran analysis

7mhm.H.pdb, model 24




93.8% (285/304) of all residues were in favored (98%) regions.
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[24] 45 Thr (-53.8, 175.5)
[24] 46 Ser (14.4, -156.6)
[24] 50 Leu (141.2, -29.1)
[24] 71 Gly (7.1, -80.2)
[24] 302 Gly (54.5, 160.2)
[24] 304 Thr (176.5, -95.3)
[24] 305 Phe (100.0, 57.6)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 25









94.7% (288/304) of all residues were in favored (98%) regions.
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

[25] 46 Ser (-3.3, -147.2)
[25] 50 Leu (49.7, -61.3)
[25] 152 Ile (-147.8, -145.9)
[25] 153 Asp (112.0, -113.1)
[25] 154 Tyr (-29.7, -71.1)
[25] 155 Asp (149.1, 20.2)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 26









93.8% (285/304) of all residues were in favored (98%) regions.
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[26] 47 Glu (-50.5, -1.4)
[26] 49 Met (-6.1, -53.6)
[26] 50 Leu (-36.7, -20.5)
[26] 71 Gly (23.1, -103.5)
[26] 73 Val (178.6, 149.5)
[26] 153 Asp (14.6, -59.2)
[26] 305 Phe (82.5, 89.0)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 27









93.8% (285/304) of all residues were in favored (98%) regions.
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[27] 153 Asp (-15.3, -51.3)
[27] 154 Tyr (-50.1, 74.5)
[27] 193 Ala (-51.4, -179.5)
[27] 302 Gly (62.0, 130.4)
[27] 303 Val (-38.8, 98.2)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 28









92.4% (281/304) of all residues were in favored (98%) regions.
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[28] 45 Thr (-52.5, 97.7)
[28] 46 Ser (-65.1, 61.8)
[28] 48 Asp (-174.8, 47.2)
[28] 73 Val (-24.6, 99.3)
[28] 154 Tyr (-34.9, -30.4)
[28] 234 Ala (-33.4, -77.1)
[28] 303 Val (-3.3, 113.7)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 29




94.4% (287/304) of all residues were in favored (98%) regions.
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[29] 23 Gly (49.4, -99.7)
[29] 59 Ile (-48.4, -119.7)
[29] 60 Arg (-42.4, 33.0)
[29] 72 Asn (-69.7, 22.7)
[29] 153 Asp (35.1, -79.8)
[29] 302 Gly (144.1, 60.8)
[29] 303 Val (28.3, 91.2)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 30









92.8% (282/304) of all residues were in favored (98%) regions.
96.4% (293/304) of all residues were in allowed (>99.8%) regions.

There were 11 outliers (phi, psi):

[30] 70 Ala (-75.0, -115.1)	[30] 277 Asn (39.8, 83.0)
[30] 71 Gly (-65.2, -86.6)	[30] 302 Gly (-156.2, -68.8)
[30] 106 Ile (-79.3, -169.1)	[30] 303 Val (97.4, 65.7)
[30] 142 Asn (-14.5, 93.4)	
[30] 153 Asp (74.0, -140.4)	
[30] 155 Asp (19.5, 91.1)	
[30] 156 Cys (167.0, 100.4)	
[30] 191 Ala (-22.4, 125.6)	

MolProbity Ramachandran analysis

7mhm.H.pdb, model 31








94.7% (288/304) of all residues were in favored (98%) regions.
99.0% (301/304) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[31] 155 Asp (20.6, 33.8)
[31] 303 Val (62.3, 74.0)
[31] 305 Phe (-56.7, 58.0)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 32




93.8% (285/304) of all residues were in favored (98%) regions.
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

[32] 47 Glu (-29.5, -76.1)
[32] 70 Ala (-74.3, -123.5)
[32] 71 Gly (-51.6, -122.8)
[32] 154 Tyr (-18.8, 144.6)
[32] 155 Asp (57.9, -14.2)
[32] 301 Ser (153.7, 126.7)
[32] 302 Gly (-176.3, 7.0)
[32] 303 Val (-150.7, 63.4)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 33









95.4% (290/304) of all residues were in favored (98%) regions.
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [33] 72 Asn (169.3, -8.6)
- [33] 154 Tyr (35.2, 89.6)
- [33] 275 Gly (44.3, -163.5)
- [33] 277 Asn (27.2, 90.8)
- [33] 302 Gly (62.5, 123.9)
- [33] 304 Thr (-172.2, -107.0)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 34




93.4% (284/304) of all residues were in favored (98%) regions.
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[34] 52 Pro (-106.7, 137.1)	[34] 303 Val (105.0, 119.9)
[34] 71 Gly (136.9, 70.3)	
[34] 72 Asn (106.3, -4.0)	
[34] 153 Asp (-61.5, 77.2)	
[34] 154 Tyr (68.0, 118.0)	
[34] 155 Asp (-16.6, 52.1)	
[34] 300 Cys (-44.1, -84.6)	
[34] 301 Ser (-25.9, -91.7)	

MolProbity Ramachandran analysis

7mhm.H.pdb, model 35









93.1% (283/304) of all residues were in favored (98%) regions.
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[35] 71 Gly (180.0, 61.5)
[35] 72 Asn (116.1, -51.6)
[35] 155 Asp (128.9, -35.9)
[35] 165 Met (174.4, -159.7)
[35] 303 Val (61.5, 155.5)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 36



94.4% (287/304) of all residues were in favored (98%) regions.
99.3% (302/304) of all residues were in allowed (>99.8%) regions.







There were 2 outliers (phi, psi):

[36] 154 Tyr (40.2, 91.4)

[36] 302 Gly (39.8, -82.4)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 37









95.7% (291/304) of all residues were in favored (98%) regions.
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[37] 46 Ser (48.2, -83.8)
[37] 70 Ala (-53.8, -123.5)
[37] 301 Ser (-62.1, 21.8)
[37] 303 Val (-18.8, 77.8)
[37] 304 Thr (52.9, -74.3)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 38




93.4% (284/304) of all residues were in favored (98%) regions.
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[38] 23 Gly (26.5, -105.2)	[38] 304 Thr (121.9, -97.9)
[38] 71 Gly (-44.3, 65.8)	
[38] 72 Asn (41.1, 90.7)	
[38] 73 Val (159.0, 89.8)	
[38] 154 Tyr (91.1, 112.0)	
[38] 155 Asp (19.5, -56.3)	
[38] 156 Cys (-35.8, 144.2)	
[38] 191 Ala (-49.5, 98.6)	

MolProbity Ramachandran analysis

7mhm.H.pdb, model 39




93.4% (284/304) of all residues were in favored (98%) regions.
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[39] 71 Gly (-169.7, 81.9)
[39] 154 Tyr (57.3, 145.2)
[39] 155 Asp (-6.4, 139.4)
[39] 156 Cys (97.2, 118.4)
[39] 302 Gly (-20.6, -55.5)
[39] 303 Val (25.4, -62.1)
[39] 305 Phe (176.4, 65.0)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 40




94.1% (286/304) of all residues were in favored (98%) regions.
99.3% (302/304) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[40] 300 Cys (-41.7, -87.8)
[40] 305 Phe (161.6, -43.1)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 41









92.1% (280/304) of all residues were in favored (98%) regions.
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [41] 51 Asn (155.8, 58.9)
- [41] 154 Tyr (18.2, 66.2)
- [41] 155 Asp (108.1, -18.6)
- [41] 301 Ser (-54.7, 6.4)
- [41] 302 Gly (-146.4, -40.9)
- [41] 303 Val (33.9, 75.1)
- [41] 304 Thr (173.6, -100.7)
- [41] 305 Phe (147.3, -28.3)

MolProbity Ramachandran analysis

7mhm.H.pdb, model 42









92.8% (282/304) of all residues were in favored (98%) regions.
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[42] 47 Glu (-46.9, -76.5)	[42] 305 Phe (41.4, -44.3)
[42] 48 Asp (-27.4, -42.1)	
[42] 70 Ala (-73.8, -108.9)	
[42] 154 Tyr (41.6, 105.9)	
[42] 300 Cys (-164.6, -125.3)	
[42] 301 Ser (1.4, -62.6)	
[42] 303 Val (6.2, 58.9)	
[42] 304 Thr (-177.7, 27.2)	

MolProbity Ramachandran analysis

7mhm.H.pdb, model 43



93.4% (284/304) of all residues were in favored (98%) regions.
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[43] 72 Asn (90.1, -169.8)
[43] 73 Val (47.7, 129.1)
[43] 154 Tyr (131.1, 116.4)
[43] 303 Val (47.6, 92.8)