


MolProbity Ramachandran analysis

7mhn.H.pdb, all models



89.5% (4900/5472) of all residues were in favored (98%) regions.

96.7% (5294/5472) of all residues were in allowed (>99.8%) regions.

This list is truncated; use the MolProbity multi-chart.html for complete list.







There were 178 outliers (phi, psi):

[1] 71 Gly (63.9, -53.4)	[2] 73 Val (125.0, 111.0)	[3] 96 Pro (-58.6, -81.0)
[1] 73 Val (56.8, 102.6)	[2] 154 Tyr (-57.1, -83.9)	[3] 97 Lys (-61.0, 58.4)
[1] 154 Tyr (167.2, -76.1)	[2] 184 Pro (-66.9, -67.4)	[3] 154 Tyr (-161.5, -112.7)
[1] 301 Ser (-178.4, 35.4)	[2] 185 Phe (11.2, 135.3)	[3] 244 Gln (1.9, -40.3)
[2] 23 Gly (39.9, -90.7)	[2] 277 Asn (-12.2, 31.4)	[3] 275 Gly (56.7, 169.1)
[2] 44 Cys (-46.3, 175.5)	[2] 278 Gly (-172.9, -38.8)	[3] 277 Asn (-21.5, 94.2)
[2] 46 Ser (-16.3, -88.8)	[2] 305 Phe (106.2, -150.6)	[3] 303 Val (118.3, -166.3)
[2] 63 Asn (-46.7, 8.5)	[3] 73 Val (113.5, 144.0)	[3] 304 Thr (161.8, 6.8)
		[3] 305 Phe (26.5, -154.0)
		[4] 23 Gly (23.7, -113.3)
		[4] 24 Thr (-51.9, 2.7)
		[4] 48 Asp (-34.0, -15.4)

[4] 63 Asn (-32.5, -35.9)	[5] 154 Tyr (113.1, -49.8)	[6] 276 Met (-40.7, -93.1)
[4] 71 Gly (22.7, -101.8)	[5] 222 Arg (-34.1, -74.4)	[6] 278 Gly (38.9, -100.1)
[4] 73 Val (124.4, 158.3)	[5] 223 Phe (-39.0, 155.8)	[6] 279 Arg (-41.0, -160.1)
[4] 80 His (177.7, 121.5)	[5] 274 Asn (-167.2, -11.2)	[6] 300 Cys (141.0, 11.9)
[4] 153 Asp (-79.2, -103.8)	[5] 277 Asn (39.3, 30.1)	[6] 305 Phe (101.7, -133.1)
[4] 184 Pro (-100.4, 49.8)	[5] 279 Arg (-70.5, -137.1)	[7] 23 Gly (38.6, -105.9)
[4] 303 Val (113.3, -160.5)	[5] 299 Gln (-161.9, -18.3)	[7] 62 Ser (-37.4, 155.1)
[4] 304 Thr (-169.2, -63.1)	[5] 303 Val (-123.3, -139.3)	[7] 72 Asn (-169.5, 39.1)
[4] 305 Phe (77.7, -114.4)	[5] 305 Phe (107.5, -25.5)	[7] 154 Tyr (89.7, -67.5)
[5] 71 Gly (36.7, -79.7)	[6] 91 Val (-52.5, 174.4)	[7] 189 Gln (-60.9, 92.4)
[5] 106 Ile (-72.1, -155.2)	[6] 154 Tyr (53.5, -89.7)	[7] 277 Asn (-8.9, 108.0)
[5] 152 Ile (149.5, 115.9)	[6] 238 Asn (20.3, 66.1)	[7] 286 Leu (160.3, 123.7)

MolProbity Ramachandran analysis

7mhn.H.pdb, model 1









91.1% (277/304) of all residues were in favored (98%) regions.
98.7% (300/304) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [1] 71 Gly (63.9, -53.4)
- [1] 73 Val (56.8, 102.6)
- [1] 154 Tyr (167.2, -76.1)
- [1] 301 Ser (-178.4, 35.4)

MolProbity Ramachandran analysis

7mhn.H.pdb, model 2




89.5% (272/304) of all residues were in favored (98%) regions.
96.4% (293/304) of all residues were in allowed (>99.8%) regions.

There were 11 outliers (phi, psi):

[2] 23 Gly (39.9, -90.7)	[2] 277 Asn (-12.2, 31.4)
[2] 44 Cys (-46.3, 175.5)	[2] 278 Gly (-172.9, -38.8)
[2] 46 Ser (-16.3, -88.8)	[2] 305 Phe (106.2, -150.6)
[2] 63 Asn (-46.7, 8.5)	
[2] 73 Val (125.0, 111.0)	
[2] 154 Tyr (-57.1, -83.9)	
[2] 184 Pro (-66.9, -67.4)	
[2] 185 Phe (11.2, 135.3)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 3




89.8% (273/304) of all residues were in favored (98%) regions.
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[3] 73 Val (113.5, 144.0)	[3] 304 Thr (161.8, 6.8)
[3] 96 Pro (-58.6, -81.0)	[3] 305 Phe (26.5, -154.0)
[3] 97 Lys (-61.0, 58.4)	
[3] 154 Tyr (-161.5, -112.7)	
[3] 244 Gln (1.9, -40.3)	
[3] 275 Gly (56.7, 169.1)	
[3] 277 Asn (-21.5, 94.2)	
[3] 303 Val (118.3, -166.3)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 4









91.1% (277/304) of all residues were in favored (98%) regions.
96.1% (292/304) of all residues were in allowed (>99.8%) regions.

There were 12 outliers (phi, psi):

[4] 23 Gly (23.7, -113.3)	[4] 184 Pro (-100.4, 49.8)
[4] 24 Thr (-51.9, 2.7)	[4] 303 Val (113.3, -160.5)
[4] 48 Asp (-34.0, -15.4)	[4] 304 Thr (-169.2, -63.1)
[4] 63 Asn (-32.5, -35.9)	[4] 305 Phe (77.7, -114.4)
[4] 71 Gly (22.7, -101.8)	
[4] 73 Val (124.4, 158.3)	
[4] 80 His (177.7, 121.5)	
[4] 153 Asp (-79.2, -103.8)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 5









90.8% (276/304) of all residues were in favored (98%) regions.
96.1% (292/304) of all residues were in allowed (>99.8%) regions.

There were 12 outliers (phi, psi):

[5] 71 Gly (36.7, -79.7)	[5] 279 Arg (-70.5, -137.1)
[5] 106 Ile (-72.1, -155.2)	[5] 299 Gln (-161.9, -18.3)
[5] 152 Ile (149.5, 115.9)	[5] 303 Val (-123.3, -139.3)
[5] 154 Tyr (113.1, -49.8)	[5] 305 Phe (107.5, -25.5)
[5] 222 Arg (-34.1, -74.4)	
[5] 223 Phe (-39.0, 155.8)	
[5] 274 Asn (-167.2, -11.2)	
[5] 277 Asn (39.3, 30.1)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 6




90.8% (276/304) of all residues were in favored (98%) regions.
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [6] 91 Val (-52.5, 174.4)
- [6] 154 Tyr (53.5, -89.7)
- [6] 238 Asn (20.3, 66.1)
- [6] 276 Met (-40.7, -93.1)
- [6] 278 Gly (38.9, -100.1)
- [6] 279 Arg (-41.0, -160.1)
- [6] 300 Cys (141.0, 11.9)
- [6] 305 Phe (101.7, -133.1)

MolProbity Ramachandran analysis

7mhn.H.pdb, model 7









89.5% (272/304) of all residues were in favored (98%) regions.
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[7] 23 Gly (38.6, -105.9)	[7] 300 Cys (158.8, 42.1)
[7] 62 Ser (-37.4, 155.1)	[7] 305 Phe (72.0, -101.6)
[7] 72 Asn (-169.5, 39.1)	
[7] 154 Tyr (89.7, -67.5)	
[7] 189 Gln (-60.9, 92.4)	
[7] 277 Asn (-8.9, 108.0)	
[7] 286 Leu (160.3, 123.7)	
[7] 299 Gln (-59.5, 47.8)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 8




89.5% (272/304) of all residues were in favored (98%) regions.
96.1% (292/304) of all residues were in allowed (>99.8%) regions.

There were 12 outliers (phi, psi):

[8] 46 Ser (-41.7, -98.9)	[8] 276 Met (-37.6, -21.8)
[8] 71 Gly (58.7, -65.4)	[8] 303 Val (-162.9, -87.5)
[8] 80 His (172.3, 121.4)	[8] 304 Thr (-85.5, -108.7)
[8] 93 Thr (176.7, 111.4)	[8] 305 Phe (94.7, -128.2)
[8] 154 Tyr (24.0, -69.2)	
[8] 222 Arg (-24.4, 55.1)	
[8] 223 Phe (-61.9, -92.1)	
[8] 275 Gly (-25.6, -12.5)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 9









89.5% (272/304) of all residues were in favored (98%) regions.
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[9] 71 Gly (61.2, -77.5)	[9] 305 Phe (133.1, -105.7)
[9] 73 Val (-157.5, 76.4)	
[9] 81 Ser (159.9, 164.7)	
[9] 154 Tyr (62.9, -88.9)	
[9] 222 Arg (39.2, -46.0)	
[9] 229 Asp (171.3, -62.2)	
[9] 276 Met (-58.2, 58.1)	
[9] 303 Val (-169.4, 56.9)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 10









88.8% (270/304) of all residues were in favored (98%) regions.
96.1% (292/304) of all residues were in allowed (>99.8%) regions.

There were 12 outliers (phi, psi):

[10] 46 Ser (-155.3, -91.4)	[10] 276 Met (-9.7, 72.8)
[10] 47 Glu (-11.0, -47.6)	[10] 279 Arg (-37.5, 79.2)
[10] 49 Met (-32.3, -39.4)	[10] 301 Ser (94.2, 49.3)
[10] 71 Gly (53.0, -66.5)	[10] 305 Phe (35.1, -157.0)
[10] 73 Val (95.5, 124.1)	
[10] 106 Ile (-47.0, 170.7)	
[10] 154 Tyr (79.1, -116.7)	
[10] 275 Gly (175.1, 24.2)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 11




88.2% (268/304) of all residues were in favored (98%) regions.
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [11] 46 Ser (-22.5, -91.9)
- [11] 154 Tyr (110.5, -59.3)
- [11] 276 Met (62.2, 115.6)
- [11] 279 Arg (17.2, 158.8)
- [11] 303 Val (-162.3, -158.2)
- [11] 304 Thr (71.7, -92.7)

MolProbity Ramachandran analysis

7mhn.H.pdb, model 12




91.1% (277/304) of all residues were in favored (98%) regions.
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[12] 44 Cys (-34.3, -103.4)	[12] 304 Thr (22.2, -89.4)
[12] 45 Thr (139.9, 157.2)	
[12] 71 Gly (50.9, -92.4)	
[12] 80 His (163.6, 131.8)	
[12] 184 Pro (-98.0, 52.7)	
[12] 277 Asn (15.3, 96.0)	
[12] 300 Cys (-72.1, -159.0)	
[12] 301 Ser (-0.9, 155.0)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 13









89.5% (272/304) of all residues were in favored (98%) regions.
96.1% (292/304) of all residues were in allowed (>99.8%) regions.

There were 12 outliers (phi, psi):

[13] 47 Glu (160.5, -40.1)	[13] 299 Gln (-68.6, -83.5)
[13] 154 Tyr (173.5, -50.0)	[13] 301 Ser (122.8, 131.8)
[13] 238 Asn (29.1, 49.2)	[13] 304 Thr (-67.7, -124.6)
[13] 275 Gly (30.8, 106.3)	[13] 305 Phe (102.5, 3.3)
[13] 276 Met (-61.1, 21.0)	
[13] 277 Asn (4.0, 143.8)	
[13] 279 Arg (-173.3, -150.4)	
[13] 283 Gly (54.2, -4.8)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 14









88.8% (270/304) of all residues were in favored (98%) regions.
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [14] 23 Gly (34.4, -150.2)
- [14] 47 Glu (150.8, -40.8)
- [14] 71 Gly (46.2, -68.4)
- [14] 142 Asn (-14.6, 107.3)
- [14] 275 Gly (-66.3, -99.6)
- [14] 276 Met (-8.5, -72.9)
- [14] 277 Asn (127.8, 143.2)
- [14] 304 Thr (161.2, 90.2)

MolProbity Ramachandran analysis

7mhn.H.pdb, model 15









88.8% (270/304) of all residues were in favored (98%) regions.
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[15] 46 Ser (-19.3, 98.4)	[15] 304 Thr (143.9, 112.4)
[15] 47 Glu (129.3, -22.5)	[15] 305 Phe (-68.1, 46.4)
[15] 71 Gly (32.0, -96.4)	
[15] 154 Tyr (113.3, -129.3)	
[15] 278 Gly (-34.1, -32.9)	
[15] 279 Arg (-53.8, -147.8)	
[15] 301 Ser (-34.2, 92.2)	
[15] 303 Val (-57.8, 65.8)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 16




89.1% (271/304) of all residues were in favored (98%) regions.
96.4% (293/304) of all residues were in allowed (>99.8%) regions.

There were 11 outliers (phi, psi):

[16] 46 Ser (-36.9, 163.5)	[16] 301 Ser (13.3, 33.0)
[16] 59 Ile (-28.9, -53.4)	[16] 302 Gly (46.0, 85.9)
[16] 71 Gly (52.3, -41.5)	[16] 303 Val (-152.2, 25.8)
[16] 72 Asn (-162.6, -36.4)	
[16] 154 Tyr (117.6, -157.4)	
[16] 274 Asn (-15.2, -129.8)	
[16] 276 Met (-176.4, 68.3)	
[16] 278 Gly (35.4, -37.4)	

MolProbity Ramachandran analysis

7mhn.H.pdb, model 17




87.5% (266/304) of all residues were in favored (98%) regions.
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[17] 47 Glu (58.5, -32.8)
[17] 71 Gly (24.1, -88.3)
[17] 119 Asn (36.6, 35.5)
[17] 154 Tyr (150.8, -98.4)
[17] 274 Asn (19.2, -53.3)
[17] 304 Thr (130.9, -150.9)
[17] 305 Phe (87.2, 82.0)

MolProbity Ramachandran analysis

7mhn.H.pdb, model 18



88.5% (269/304) of all residues were in favored (98%) regions.
95.1% (289/304) of all residues were in allowed (>99.8%) regions.

There were 15 outliers (phi, psi):

[18] 25 Thr (173.0, 136.8)	[18] 271 Leu (-7.8, -43.7)
[18] 47 Glu (116.3, -71.0)	[18] 274 Asn (21.2, 36.0)
[18] 59 Ile (-71.1, 16.4)	[18] 275 Gly (1.3, -53.1)
[18] 71 Gly (28.3, -78.0)	[18] 276 Met (-9.0, 104.8)
[18] 73 Val (-38.1, 163.9)	[18] 278 Gly (40.3, -53.6)
[18] 106 Ile (-72.1, -170.7)	[18] 302 Gly (46.8, 87.4)
[18] 222 Arg (0.7, -90.9)	[18] 303 Val (-67.9, 71.5)
[18] 270 Glu (-57.7, -96.3)	