


MolProbity Ramachandran analysis

7mho.H.pdb, all models



90.5% (5505/6080) of all residues were in favored (98%) regions.

97.0% (5899/6080) of all residues were in allowed (>99.8%) regions.

This list is truncated; use the MolProbity multi-chart.html for complete list.

There were 181 outliers (phi, psi):

[1] 72 Asn (-75.7, -114.5)	[1] 303 Val (-147.6, -70.3)	[3] 68 Val (-69.3, -125.2)
[1] 73 Val (-13.1, 99.9)	[1] 304 Thr (91.7, 143.8)	[3] 69 Gln (120.3, 115.4)
[1] 153 Asp (31.4, 11.9)	[2] 73 Val (-31.1, 154.2)	[3] 71 Gly (48.7, -88.6)
[1] 191 Ala (-51.4, 102.9)	[2] 153 Asp (30.0, -0.5)	[3] 73 Val (-13.2, 118.2)
[1] 194 Ala (-36.8, -90.3)	[2] 154 Tyr (173.8, -92.1)	[3] 274 Asn (-171.6, -14.0)
[1] 222 Arg (-11.1, -63.6)	[2] 280 Thr (-26.3, 146.5)	[3] 277 Asn (51.5, 84.1)
[1] 273 Gln (-59.0, -76.3)	[2] 301 Ser (-27.3, 164.7)	[3] 303 Val (145.6, 43.2)
[1] 302 Gly (151.4, 35.6)	[2] 302 Gly (59.7, 90.5)	[3] 305 Phe (-31.7, -132.4)
		[4] 71 Gly (53.3, -93.1)
		[4] 155 Asp (158.7, -48.5)
		[4] 273 Gln (-73.9, -97.2)
		[4] 274 Asn (-19.1, -73.8)

[4] 277 Asn (37.5, -173.3)

[4] 301 Ser (176.1, 45.6)

[4] 303 Val (62.6, 93.5)

[4] 304 Thr (-175.9, 54.2)

[5] 72 Asn (-45.6, -100.7)

[5] 154 Tyr (88.5, -139.9)

[5] 191 Ala (-31.1, 94.5)

[5] 224 Thr (56.8, -179.7)

[5] 303 Val (-160.8, 112.5)

[5] 305 Phe (-30.6, 62.0)

[6] 73 Val (-87.2, -133.7)

[6] 154 Tyr (13.9, -19.8)

[6] 155 Asp (-173.6, 6.0)

[6] 194 Ala (-59.0, -94.5)

[6] 224 Thr (-44.3, -174.0)

[6] 232 Leu (-33.3, -34.5)

[6] 280 Thr (-46.4, 164.7)

[6] 300 Cys (157.1, -24.2)

[6] 303 Val (71.7, 32.1)

[6] 305 Phe (74.3, 131.3)

[7] 68 Val (-79.4, -124.5)

[7] 69 Gln (119.7, 103.5)

[7] 71 Gly (9.1, -79.9)

[7] 223 Phe (-20.9, 129.7)

[7] 274 Asn (-51.0, 49.1)

[7] 280 Thr (-8.4, 160.5)

[7] 303 Val (25.7, 55.0)

[7] 305 Phe (178.9, -66.3)

[8] 24 Thr (-172.3, -15.6)

[8] 108 Pro (-29.5, 126.8)

[8] 154 Tyr (33.3, -89.9)

[8] 223 Phe (-48.9, 174.4)

[8] 277 Asn (43.9, 143.3)







[8] 278 Gly (33.4, -76.8)

[8] 279 Arg (-46.7, 174.4)

[8] 301 Ser (-178.5, -99.8)

MolProbity Ramachandran analysis

7mho.H.pdb, model 1









86.8% (264/304) of all residues were in favored (98%) regions.
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[1] 72 Asn (-75.7, -114.5)	[1] 303 Val (-147.6, -70.3)
[1] 73 Val (-13.1, 99.9)	[1] 304 Thr (91.7, 143.8)
[1] 153 Asp (31.4, 11.9)	
[1] 191 Ala (-51.4, 102.9)	
[1] 194 Ala (-36.8, -90.3)	
[1] 222 Arg (-11.1, -63.6)	
[1] 273 Gln (-59.0, -76.3)	
[1] 302 Gly (151.4, 35.6)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 2









92.1% (280/304) of all residues were in favored (98%) regions.
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [2] 73 Val (-31.1, 154.2)
- [2] 153 Asp (30.0, -0.5)
- [2] 154 Tyr (173.8, -92.1)
- [2] 280 Thr (-26.3, 146.5)
- [2] 301 Ser (-27.3, 164.7)
- [2] 302 Gly (59.7, 90.5)

MolProbity Ramachandran analysis

7mho.H.pdb, model 3




90.8% (276/304) of all residues were in favored (98%) regions.
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [3] 68 Val (-69.3, -125.2)
- [3] 69 Gln (120.3, 115.4)
- [3] 71 Gly (48.7, -88.6)
- [3] 73 Val (-13.2, 118.2)
- [3] 274 Asn (-171.6, -14.0)
- [3] 277 Asn (51.5, 84.1)
- [3] 303 Val (145.6, 43.2)
- [3] 305 Phe (-31.7, -132.4)

MolProbity Ramachandran analysis

7mho.H.pdb, model 4









89.5% (272/304) of all residues were in favored (98%) regions.
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [4] 71 Gly (53.3, -93.1)
- [4] 155 Asp (158.7, -48.5)
- [4] 273 Gln (-73.9, -97.2)
- [4] 274 Asn (-19.1, -73.8)
- [4] 277 Asn (37.5, -173.3)
- [4] 301 Ser (176.1, 45.6)
- [4] 303 Val (62.6, 93.5)
- [4] 304 Thr (-175.9, 54.2)

MolProbity Ramachandran analysis

7mho.H.pdb, model 5









93.1% (283/304) of all residues were in favored (98%) regions.
98.0% (298/304) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [5] 72 Asn (-45.6, -100.7)
- [5] 154 Tyr (88.5, -139.9)
- [5] 191 Ala (-31.1, 94.5)
- [5] 224 Thr (56.8, -179.7)
- [5] 303 Val (-160.8, 112.5)
- [5] 305 Phe (-30.6, 62.0)

MolProbity Ramachandran analysis

7mho.H.pdb, model 6









91.1% (277/304) of all residues were in favored (98%) regions.
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[6] 73 Val (-87.2, -133.7)	[6] 303 Val (71.7, 32.1)
[6] 154 Tyr (13.9, -19.8)	[6] 305 Phe (74.3, 131.3)
[6] 155 Asp (-173.6, 6.0)	
[6] 194 Ala (-59.0, -94.5)	
[6] 224 Thr (-44.3, -174.0)	
[6] 232 Leu (-33.3, -34.5)	
[6] 280 Thr (-46.4, 164.7)	
[6] 300 Cys (157.1, -24.2)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 7









90.1% (274/304) of all residues were in favored (98%) regions.
97.4% (296/304) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

[7] 68 Val (-79.4, -124.5)
[7] 69 Gln (119.7, 103.5)
[7] 71 Gly (9.1, -79.9)
[7] 223 Phe (-20.9, 129.7)
[7] 274 Asn (-51.0, 49.1)
[7] 280 Thr (-8.4, 160.5)
[7] 303 Val (25.7, 55.0)
[7] 305 Phe (178.9, -66.3)

MolProbity Ramachandran analysis

7mho.H.pdb, model 8









92.1% (280/304) of all residues were in favored (98%) regions.
96.4% (293/304) of all residues were in allowed (>99.8%) regions.

There were 11 outliers (phi, psi):

[8] 24 Thr (-172.3, -15.6)	[8] 302 Gly (35.8, -67.6)
[8] 108 Pro (-29.5, 126.8)	[8] 303 Val (70.6, 10.8)
[8] 154 Tyr (33.3, -89.9)	[8] 304 Thr (156.4, 28.9)
[8] 223 Phe (-48.9, 174.4)	
[8] 277 Asn (43.9, 143.3)	
[8] 278 Gly (33.4, -76.8)	
[8] 279 Arg (-46.7, 174.4)	
[8] 301 Ser (-178.5, -99.8)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 9









88.2% (268/304) of all residues were in favored (98%) regions.
96.4% (293/304) of all residues were in allowed (>99.8%) regions.

There were 11 outliers (phi, psi):

[9] 45 Thr (-63.1, -173.1)	[9] 256 Gln (-17.2, -63.9)
[9] 68 Val (-65.5, -98.6)	[9] 301 Ser (178.7, -68.7)
[9] 69 Gln (103.0, 94.9)	[9] 302 Gly (11.8, 74.2)
[9] 71 Gly (60.1, -71.2)	
[9] 84 Asn (33.0, -103.5)	
[9] 154 Tyr (76.8, -116.0)	
[9] 221 Asn (-43.2, 170.3)	
[9] 224 Thr (-24.9, 178.8)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 10









88.8% (270/304) of all residues were in favored (98%) regions.
96.4% (293/304) of all residues were in allowed (>99.8%) regions.

There were 11 outliers (phi, psi):

[10] 46 Ser (-3.2, -131.4)	[10] 277 Asn (-16.1, 88.9)
[10] 48 Asp (-24.5, -40.3)	[10] 303 Val (-156.5, 52.6)
[10] 71 Gly (61.5, -75.0)	[10] 305 Phe (96.6, -62.6)
[10] 154 Tyr (46.3, -51.3)	
[10] 221 Asn (43.1, 120.8)	
[10] 274 Asn (28.7, 24.6)	
[10] 275 Gly (43.8, 108.1)	
[10] 276 Met (-57.8, 21.5)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 11




89.1% (271/304) of all residues were in favored (98%) regions.
95.7% (291/304) of all residues were in allowed (>99.8%) regions.

There were 13 outliers (phi, psi):

[11] 23 Gly (-22.4, -46.5)	[11] 273 Gln (-70.1, -154.0)
[11] 24 Thr (-41.8, -75.2)	[11] 274 Asn (30.1, -72.9)
[11] 71 Gly (58.1, -62.3)	[11] 277 Asn (4.7, 135.2)
[11] 72 Asn (-54.0, -94.6)	[11] 302 Gly (-22.9, 81.2)
[11] 73 Val (-152.7, -149.8)	[11] 303 Val (-10.9, 82.5)
[11] 84 Asn (38.6, -106.0)	
[11] 154 Tyr (77.2, -137.7)	
[11] 155 Asp (-21.5, -55.6)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 12









90.1% (274/304) of all residues were in favored (98%) regions.
96.1% (292/304) of all residues were in allowed (>99.8%) regions.

There were 12 outliers (phi, psi):

[12] 23 Gly (43.8, -93.3)	[12] 277 Asn (-50.4, 101.4)
[12] 24 Thr (-36.1, -173.9)	[12] 301 Ser (-9.9, -107.3)
[12] 25 Thr (-65.3, -165.9)	[12] 303 Val (57.4, 131.3)
[12] 154 Tyr (61.4, -70.1)	[12] 304 Thr (-65.0, -141.8)
[12] 191 Ala (-25.9, 112.8)	
[12] 221 Asn (173.6, 132.0)	
[12] 257 Thr (178.1, -13.9)	
[12] 276 Met (-59.8, 19.9)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 13




92.4% (281/304) of all residues were in favored (98%) regions.
98.4% (299/304) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[13] 15 Gly (-62.6, 13.9)
[13] 71 Gly (42.6, -76.2)
[13] 196 Thr (-22.3, 131.3)
[13] 301 Ser (-64.0, -163.8)
[13] 303 Val (56.5, 140.8)

MolProbity Ramachandran analysis

7mho.H.pdb, model 14









91.4% (278/304) of all residues were in favored (98%) regions.
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[14] 72 Asn (-59.8, 4.4)	[14] 304 Thr (94.7, 160.6)
[14] 154 Tyr (155.7, -96.5)	
[14] 191 Ala (-40.4, 105.3)	
[14] 193 Ala (57.9, 142.1)	
[14] 222 Arg (-33.2, -2.9)	
[14] 223 Phe (-50.2, -160.5)	
[14] 277 Asn (28.6, 18.1)	
[14] 303 Val (92.4, -38.5)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 15




92.8% (282/304) of all residues were in favored (98%) regions.
97.0% (295/304) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[15] 71 Gly (59.3, -19.5)	[15] 304 Thr (110.2, 124.0)
[15] 154 Tyr (-40.1, -12.8)	
[15] 155 Asp (-69.6, -146.5)	
[15] 156 Cys (-6.5, 118.3)	
[15] 276 Met (-58.2, -76.1)	
[15] 277 Asn (65.6, -84.2)	
[15] 301 Ser (-58.8, -149.8)	
[15] 303 Val (-177.4, 14.8)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 16









90.8% (276/304) of all residues were in favored (98%) regions.
96.4% (293/304) of all residues were in allowed (>99.8%) regions.

There were 11 outliers (phi, psi):

[16] 71 Gly (30.4, -80.6)	[16] 301 Ser (-51.3, -98.0)
[16] 97 Lys (-162.8, 3.7)	[16] 303 Val (52.9, -53.1)
[16] 154 Tyr (175.6, 114.1)	[16] 304 Thr (-166.5, 27.2)
[16] 274 Asn (-169.0, -29.4)	
[16] 278 Gly (64.9, -60.4)	
[16] 279 Arg (-21.8, 55.8)	
[16] 280 Thr (-37.2, 154.9)	
[16] 300 Cys (-79.9, -127.9)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 17









91.8% (279/304) of all residues were in favored (98%) regions.
97.7% (297/304) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [17] 23 Gly (38.0, -103.6)
- [17] 154 Tyr (31.0, 97.2)
- [17] 276 Met (-46.4, -80.5)
- [17] 277 Asn (-30.1, 66.4)
- [17] 279 Arg (96.6, 174.2)
- [17] 303 Val (71.9, -144.5)
- [17] 305 Phe (127.1, 93.1)

MolProbity Ramachandran analysis

7mho.H.pdb, model 18




88.8% (270/304) of all residues were in favored (98%) regions.
95.7% (291/304) of all residues were in allowed (>99.8%) regions.

There were 13 outliers (phi, psi):

[18] 68 Val (-72.9, -156.3)	[18] 277 Asn (-53.0, 68.8)
[18] 69 Gln (156.9, 136.2)	[18] 301 Ser (85.8, -130.8)
[18] 71 Gly (39.4, -83.9)	[18] 303 Val (98.6, 160.8)
[18] 78 Ile (-153.3, 27.6)	[18] 304 Thr (-21.7, -139.9)
[18] 155 Asp (46.0, -50.5)	[18] 305 Phe (-8.9, -72.1)
[18] 223 Phe (-59.5, -169.9)	
[18] 273 Gln (-70.1, -86.4)	
[18] 274 Asn (-46.6, -162.7)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 19









90.5% (275/304) of all residues were in favored (98%) regions.
96.7% (294/304) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[19] 154 Tyr (41.8, 108.0)	[19] 304 Thr (38.8, 82.1)
[19] 155 Asp (111.9, -57.7)	[19] 305 Phe (119.7, -15.9)
[19] 165 Met (176.8, -167.5)	
[19] 275 Gly (-30.7, -175.9)	
[19] 299 Gln (-64.0, 74.0)	
[19] 300 Cys (-163.7, -20.0)	
[19] 301 Ser (146.1, 1.7)	
[19] 303 Val (-5.6, 108.5)	

MolProbity Ramachandran analysis

7mho.H.pdb, model 20



90.5% (275/304) of all residues were in favored (98%) regions.
99.0% (301/304) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):
[20] 154 Tyr (63.0, -75.6)
[20] 223 Phe (-40.4, -169.8)
[20] 277 Asn (53.4, 119.0)