







MolProbity Ramachandran analysis

2k7x.H.pdb, all models









90.8% (2251/2478) of all residues were in favored (98%) regions.
99.0% (2452/2478) of all residues were in allowed (>99.8%) regions.

There were 26 outliers (phi, psi):

[1] 188 Arg (40.7, -90.1)	[6] 251 Gly (-25.0, -60.8)	[11] 303 Val (-74.7, 39.3)
[1] 189 Gln (47.7, -84.0)	[6] 304 Thr (55.8, -175.7)	[13] 251 Gly (-26.1, -59.6)
[1] 277 Asn (60.2, -24.9)	[7] 302 Gly (-177.8, 63.5)	[15] 303 Val (-75.7, 40.6)
[1] 301 Ser (-65.6, 21.7)	[8] 217 Arg (-72.0, 27.6)	[18] 278 Gly (-175.9, -48.1)
[1] 303 Val (105.2, -24.1)	[8] 278 Gly (-176.4, -39.7)	[19] 217 Arg (-71.4, 22.7)
[3] 217 Arg (-67.9, 21.3)	[9] 300 Cys (30.3, 48.0)	[19] 251 Gly (-23.8, -60.7)
[4] 190 Thr (-69.2, 25.3)	[10] 194 Ala (-71.2, 32.2)	[19] 303 Val (-78.1, 28.5)
[6] 189 Gln (29.7, -142.3)	[10] 278 Gly (164.3, 12.7)	[20] 251 Gly (-26.7, -58.7)
		[20] 303 Val (-78.8, 36.2)
		[21] 278 Gly (179.8, -48.0)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 1









83.1% (98/118) of all residues were in favored (98%) regions.
95.8% (113/118) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [1] 188 Arg (40.7, -90.1)
- [1] 189 Gln (47.7, -84.0)
- [1] 277 Asn (60.2, -24.9)
- [1] 301 Ser (-65.6, 21.7)
- [1] 303 Val (105.2, -24.1)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 2









93.2% (110/118) of all residues were in favored (98%) regions.
100.0% (118/118) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2k7x.H.pdb, model 3









89.0% (105/118) of all residues were in favored (98%) regions.
99.2% (117/118) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[3] 217 Arg (-67.9, 21.3)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 4









90.7% (107/118) of all residues were in favored (98%) regions.
99.2% (117/118) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[4] 190 Thr (-69.2, 25.3)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 5









90.7% (107/118) of all residues were in favored (98%) regions.
100.0% (118/118) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2k7x.H.pdb, model 6









89.8% (106/118) of all residues were in favored (98%) regions.
97.5% (115/118) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[6] 189 Gln (29.7, -142.3)
[6] 251 Gly (-25.0, -60.8)
[6] 304 Thr (55.8, -175.7)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 7









92.4% (109/118) of all residues were in favored (98%) regions.
99.2% (117/118) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[7] 302 Gly (-177.8, 63.5)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 8









92.4% (109/118) of all residues were in favored (98%) regions.
98.3% (116/118) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[8] 217 Arg (-72.0, 27.6)
[8] 278 Gly (-176.4, -39.7)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 9









92.4% (109/118) of all residues were in favored (98%) regions.
99.2% (117/118) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[9] 300 Cys (30.3, 48.0)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 10









91.5% (108/118) of all residues were in favored (98%) regions.
98.3% (116/118) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[10] 194 Ala (-71.2, 32.2)
[10] 278 Gly (164.3, 12.7)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 11









90.7% (107/118) of all residues were in favored (98%) regions.
99.2% (117/118) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[11] 303 Val (-74.7, 39.3)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 12









93.2% (110/118) of all residues were in favored (98%) regions.
100.0% (118/118) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2k7x.H.pdb, model 13









89.8% (106/118) of all residues were in favored (98%) regions.
99.2% (117/118) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[13] 251 Gly (-26.1, -59.6)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 14









90.7% (107/118) of all residues were in favored (98%) regions.
100.0% (118/118) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2k7x.H.pdb, model 15









89.8% (106/118) of all residues were in favored (98%) regions.
99.2% (117/118) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[15] 303 Val (-75.7, 40.6)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 16









92.4% (109/118) of all residues were in favored (98%) regions.
100.0% (118/118) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2k7x.H.pdb, model 17









90.7% (107/118) of all residues were in favored (98%) regions.
100.0% (118/118) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2k7x.H.pdb, model 18









90.7% (107/118) of all residues were in favored (98%) regions.
99.2% (117/118) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[18] 278 Gly (-175.9, -48.1)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 19









91.5% (108/118) of all residues were in favored (98%) regions.
97.5% (115/118) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[19] 217 Arg (-71.4, 22.7)
[19] 251 Gly (-23.8, -60.7)
[19] 303 Val (-78.1, 28.5)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 20









91.5% (108/118) of all residues were in favored (98%) regions.
98.3% (116/118) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[20] 251 Gly (-26.7, -58.7)
[20] 303 Val (-78.8, 36.2)

MolProbity Ramachandran analysis

2k7x.H.pdb, model 21



91.5% (108/118) of all residues were in favored (98%) regions.
99.2% (117/118) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[21] 278 Gly (179.8, -48.0)