


MolProbity Ramachandran analysis

1yo4.H.pdb, all models



78.0% (663/850) of all residues were in favored (98%) regions.
95.8% (814/850) of all residues were in allowed (>99.8%) regions.







There were 36 outliers (phi, psi):

[1] 71 Leu (63.0, -86.0)	[3] 69 Pro (-69.6, -61.8)	[5] 10 Arg (-59.3, 98.4)
[2] 22 Ser (174.6, 61.6)	[3] 80 Gln (-175.1, 36.4)	[5] 21 Pro (-60.4, 83.3)
[2] 36 Asp (-77.5, -88.0)	[4] 29 Ser (72.4, 101.7)	[5] 69 Pro (-81.6, -69.8)
[2] 74 Arg (-175.0, 1.0)	[4] 36 Asp (176.6, 24.0)	[5] 72 Phe (168.5, -177.3)
[2] 75 Gln (-166.8, -5.6)	[4] 64 Ala (-57.2, 97.8)	[5] 77 Glu (-162.7, -35.6)
[2] 78 Val (-160.3, -13.3)	[4] 72 Phe (66.3, -88.4)	[6] 71 Leu (67.5, -88.4)
[3] 31 Phe (68.9, 126.6)	[4] 73 Ile (60.3, -82.1)	[6] 78 Val (78.8, -29.1)
[3] 67 Val (88.5, -50.8)	[4] 83 Tyr (61.4, -84.5)	[7] 23 Gly (64.0, 150.2)
		[8] 36 Asp (-63.6, -87.4)
		[8] 67 Val (57.7, -144.7)
		[8] 78 Val (66.6, 89.0)
		[8] 82 Leu (-177.5, -153.8)

[9] 67 Val (-150.3, 7.8)
[9] 68 Ser (56.6, 94.0)
[9] 71 Leu (178.5, -143.7)
[9] 82 Leu (-84.6, -128.4)
[10] 22 Ser (76.0, 122.3)
[10] 73 Ile (62.5, -76.0)
[10] 74 Arg (-163.8, -28.8)
[10] 78 Val (-165.6, 113.2)

MolProbity Ramachandran analysis

1yo4.H.pdb, model 1




75.3% (64/85) of all residues were in favored (98%) regions.
98.8% (84/85) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[1] 71 Leu (63.0, -86.0)

MolProbity Ramachandran analysis

1yo4.H.pdb, model 2



75.3% (64/85) of all residues were in favored (98%) regions.







94.1% (80/85) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [2] 22 Ser (174.6, 61.6)
- [2] 36 Asp (-77.5, -88.0)
- [2] 74 Arg (-175.0, 1.0)
- [2] 75 Gln (-166.8, -5.6)
- [2] 78 Val (-160.3, -13.3)

MolProbity Ramachandran analysis

1yo4.H.pdb, model 3









84.7% (72/85) of all residues were in favored (98%) regions.
95.3% (81/85) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [3] 31 Phe (68.9, 126.6)
- [3] 67 Val (88.5, -50.8)
- [3] 69 Pro (-69.6, -61.8)
- [3] 80 Gln (-175.1, 36.4)

MolProbity Ramachandran analysis

1yo4.H.pdb, model 4




81.2% (69/85) of all residues were in favored (98%) regions.
92.9% (79/85) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [4] 29 Ser (72.4, 101.7)
- [4] 36 Asp (176.6, 24.0)
- [4] 64 Ala (-57.2, 97.8)
- [4] 72 Phe (66.3, -88.4)
- [4] 73 Ile (60.3, -82.1)
- [4] 83 Tyr (61.4, -84.5)

MolProbity Ramachandran analysis

1yo4.H.pdb, model 5









77.6% (66/85) of all residues were in favored (98%) regions.
94.1% (80/85) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [5] 10 Arg (-59.3, 98.4)
- [5] 21 Pro (-60.4, 83.3)
- [5] 69 Pro (-81.6, -69.8)
- [5] 72 Phe (168.5, -177.3)
- [5] 77 Glu (-162.7, -35.6)

MolProbity Ramachandran analysis

1yo4.H.pdb, model 6









76.5% (65/85) of all residues were in favored (98%) regions.
97.6% (83/85) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[6] 71 Leu (67.5, -88.4)
[6] 78 Val (78.8, -29.1)

MolProbity Ramachandran analysis

1yo4.H.pdb, model 7








82.4% (70/85) of all residues were in favored (98%) regions.
98.8% (84/85) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[7] 23 Gly (64.0, 150.2)

MolProbity Ramachandran analysis

1yo4.H.pdb, model 8



76.5% (65/85) of all residues were in favored (98%) regions.

95.3% (81/85) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[8] 36 Asp (-63.6, -87.4)







[8] 67 Val (57.7, -144.7)

[8] 78 Val (66.6, 89.0)

[8] 82 Leu (-177.5, -153.8)

MolProbity Ramachandran analysis

1yo4.H.pdb, model 9









75.3% (64/85) of all residues were in favored (98%) regions.
95.3% (81/85) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [9] 67 Val (-150.3, 7.8)
- [9] 68 Ser (56.6, 94.0)
- [9] 71 Leu (178.5, -143.7)
- [9] 82 Leu (-84.6, -128.4)

MolProbity Ramachandran analysis

1yo4.H.pdb, model 10



75.3% (64/85) of all residues were in favored (98%) regions.
95.3% (81/85) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[10] 22 Ser (76.0, 122.3)
[10] 73 Ile (62.5, -76.0)
[10] 74 Arg (-163.8, -28.8)
[10] 78 Val (-165.6, 113.2)