







MolProbity Ramachandran analysis

2gdt.H.pdb, all models



78.1% (1780/2280) of all residues were in favored (98%) regions.
94.2% (2148/2280) of all residues were in allowed (>99.8%) regions.
This list is truncated; use the MolProbity multi-chart.html for complete list.







There were 132 outliers (phi, psi):

[1] 15 Val (-65.4, 65.0)	[2] 63 Ser (52.1, -176.4)	[3] 103 Thr (61.0, 105.6)
[1] 55 Gln (-156.4, -59.6)	[2] 74 Val (52.5, 177.7)	[4] 15 Val (-59.6, 65.4)
[1] 63 Ser (43.0, 78.9)	[2] 84 Ile (-28.2, -67.6)	[4] 72 His (61.7, 150.1)
[1] 84 Ile (-33.7, -56.8)	[2] 102 Glu (-55.6, -100.4)	[4] 103 Thr (72.3, 105.8)
[1] 91 Ile (-65.7, -93.3)	[3] 15 Val (-70.1, 64.0)	[5] 15 Val (-63.1, 65.4)
[1] 92 Thr (21.6, 116.4)	[3] 16 Leu (-49.8, -8.4)	[5] 55 Gln (-164.6, -56.2)
[1] 103 Thr (69.2, 102.0)	[3] 54 Glu (-72.1, -89.2)	[5] 63 Ser (23.9, 84.6)
[2] 55 Gln (-158.7, -56.4)	[3] 55 Gln (-160.5, -62.1)	[5] 71 Gly (47.9, -90.0)

[5] 84 Ile (-39.8, 66.5)	[6] 102 Glu (-75.2, -101.5)	[8] 68 Thr (36.5, 88.8)
[5] 85 Gln (90.0, -142.6)	[6] 114 Lys (28.5, 95.0)	[8] 70 His (25.2, 89.1)
[5] 86 Tyr (60.1, -81.1)	[7] 15 Val (-72.8, 62.5)	[8] 102 Glu (-65.6, -101.4)
[5] 91 Ile (-178.4, -42.7)	[7] 54 Glu (-71.9, -90.0)	[9] 15 Val (-66.3, 64.4)
[5] 103 Thr (76.3, 100.9)	[7] 55 Gln (-155.8, -57.4)	[9] 55 Gln (-171.0, -59.4)
[6] 15 Val (-68.3, 63.4)	[7] 62 Arg (-67.1, 18.1)	[9] 70 His (93.7, 113.0)
[6] 16 Leu (-34.9, -33.8)	[7] 66 Leu (24.3, 93.2)	[9] 84 Ile (-25.3, -49.2)
[6] 54 Glu (-74.7, -87.1)	[7] 82 Asp (-166.8, -38.0)	[9] 102 Glu (-82.4, -101.1)
[6] 55 Gln (-168.4, -51.6)	[7] 103 Thr (72.0, 103.1)	[10] 15 Val (-58.6, 63.8)
[6] 84 Ile (-48.4, 65.7)	[8] 15 Val (-62.8, 65.1)	[10] 54 Glu (-74.5, -93.5)
[6] 85 Gln (96.2, -86.1)	[8] 54 Glu (-71.4, -89.7)	[10] 55 Gln (-155.7, -62.5)
[6] 86 Tyr (23.2, -85.4)	[8] 55 Gln (-166.0, -65.2)	[10] 64 Asp (91.7, 156.8)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 1









78.9% (90/114) of all residues were in favored (98%) regions.
93.9% (107/114) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [1] 15 Val (-65.4, 65.0)
- [1] 55 Gln (-156.4, -59.6)
- [1] 63 Ser (43.0, 78.9)
- [1] 84 Ile (-33.7, -56.8)
- [1] 91 Ile (-65.7, -93.3)
- [1] 92 Thr (21.6, 116.4)
- [1] 103 Thr (69.2, 102.0)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 2









75.4% (86/114) of all residues were in favored (98%) regions.
95.6% (109/114) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [2] 55 Gln (-158.7, -56.4)
- [2] 63 Ser (52.1, -176.4)
- [2] 74 Val (52.5, 177.7)
- [2] 84 Ile (-28.2, -67.6)
- [2] 102 Glu (-55.6, -100.4)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 3









77.2% (88/114) of all residues were in favored (98%) regions.
93.0% (106/114) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [3] 15 Val (-70.1, 64.0)
- [3] 16 Leu (-49.8, -8.4)
- [3] 54 Glu (-72.1, -89.2)
- [3] 55 Gln (-160.5, -62.1)
- [3] 68 Thr (43.0, -147.7)
- [3] 73 Lys (-49.4, 96.3)
- [3] 84 Ile (-25.6, -70.5)
- [3] 103 Thr (61.0, 105.6)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 4









79.8% (91/114) of all residues were in favored (98%) regions.
96.5% (110/114) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [4] 15 Val (-59.6, 65.4)
- [4] 55 Gln (-169.6, -57.8)
- [4] 72 His (61.7, 150.1)
- [4] 103 Thr (72.3, 105.8)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 5









78.1% (89/114) of all residues were in favored (98%) regions.
92.1% (105/114) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[5] 15 Val (-63.1, 65.4)	[5] 103 Thr (76.3, 100.9)
[5] 55 Gln (-164.6, -56.2)	
[5] 63 Ser (23.9, 84.6)	
[5] 71 Gly (47.9, -90.0)	
[5] 84 Ile (-39.8, 66.5)	
[5] 85 Gln (90.0, -142.6)	
[5] 86 Tyr (60.1, -81.1)	
[5] 91 Ile (-178.4, -42.7)	

MolProbity Ramachandran analysis

2gdt.H.pdb, model 6









74.6% (85/114) of all residues were in favored (98%) regions.
92.1% (105/114) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[6] 15 Val (-68.3, 63.4)	[6] 114 Lys (28.5, 95.0)
[6] 16 Leu (-34.9, -33.8)	
[6] 54 Glu (-74.7, -87.1)	
[6] 55 Gln (-168.4, -51.6)	
[6] 84 Ile (-48.4, 65.7)	
[6] 85 Gln (96.2, -86.1)	
[6] 86 Tyr (23.2, -85.4)	
[6] 102 Glu (-75.2, -101.5)	

MolProbity Ramachandran analysis

2gdt.H.pdb, model 7









77.2% (88/114) of all residues were in favored (98%) regions.
93.9% (107/114) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [7] 15 Val (-72.8, 62.5)
- [7] 54 Glu (-71.9, -90.0)
- [7] 55 Gln (-155.8, -57.4)
- [7] 62 Arg (-67.1, 18.1)
- [7] 66 Leu (24.3, 93.2)
- [7] 82 Asp (-166.8, -38.0)
- [7] 103 Thr (72.0, 103.1)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 8









78.1% (89/114) of all residues were in favored (98%) regions.
94.7% (108/114) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [8] 15 Val (-62.8, 65.1)
- [8] 54 Glu (-71.4, -89.7)
- [8] 55 Gln (-166.0, -65.2)
- [8] 68 Thr (36.5, 88.8)
- [8] 70 His (25.2, 89.1)
- [8] 102 Glu (-65.6, -101.4)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 9









78.9% (90/114) of all residues were in favored (98%) regions.
95.6% (109/114) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [9] 15 Val (-66.3, 64.4)
- [9] 55 Gln (-171.0, -59.4)
- [9] 70 His (93.7, 113.0)
- [9] 84 Ile (-25.3, -49.2)
- [9] 102 Glu (-82.4, -101.1)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 10









78.9% (90/114) of all residues were in favored (98%) regions.
94.7% (108/114) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [10] 15 Val (-58.6, 63.8)
- [10] 54 Glu (-74.5, -93.5)
- [10] 55 Gln (-155.7, -62.5)
- [10] 64 Asp (91.7, 156.8)
- [10] 92 Thr (95.5, 150.1)
- [10] 103 Thr (71.2, 105.8)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 11









78.9% (90/114) of all residues were in favored (98%) regions.
93.9% (107/114) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [11] 15 Val (-61.1, 65.2)
- [11] 16 Leu (-32.9, -27.7)
- [11] 54 Glu (-73.8, -101.2)
- [11] 55 Gln (-152.0, -63.7)
- [11] 63 Ser (54.6, 170.0)
- [11] 73 Lys (-57.9, 94.0)
- [11] 103 Thr (73.4, 101.5)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 12









80.7% (92/114) of all residues were in favored (98%) regions.
94.7% (108/114) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [12] 15 Val (-72.6, 63.8)
- [12] 54 Glu (-68.4, -93.6)
- [12] 55 Gln (-168.6, -65.7)
- [12] 65 Ala (36.9, 79.3)
- [12] 100 Val (-93.5, 38.7)
- [12] 102 Glu (-42.4, -91.3)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 13









78.9% (90/114) of all residues were in favored (98%) regions.
93.0% (106/114) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [13] 15 Val (-68.9, 66.1)
- [13] 16 Leu (-25.1, -44.0)
- [13] 55 Gln (-167.6, -57.1)
- [13] 64 Asp (-164.3, -48.3)
- [13] 68 Thr (29.5, -84.7)
- [13] 72 His (56.6, 175.8)
- [13] 84 Ile (-31.9, -62.9)
- [13] 103 Thr (68.2, 96.0)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 14









82.5% (94/114) of all residues were in favored (98%) regions.
92.1% (105/114) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[14] 15 Val (-64.4, 63.9)	[14] 102 Glu (-60.1, -100.0)
[14] 54 Glu (-66.2, -94.3)	
[14] 55 Gln (-161.4, -64.6)	
[14] 63 Ser (55.6, 174.3)	
[14] 66 Leu (43.0, -90.6)	
[14] 68 Thr (46.6, 178.3)	
[14] 73 Lys (-52.1, 100.0)	
[14] 100 Val (-79.6, 38.9)	

MolProbity Ramachandran analysis

2gdt.H.pdb, model 15








76.3% (87/114) of all residues were in favored (98%) regions.
93.9% (107/114) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [15] 15 Val (-61.5, 64.5)
- [15] 54 Glu (-76.8, -93.8)
- [15] 55 Gln (-157.4, -53.8)
- [15] 58 Val (-57.3, 86.1)
- [15] 64 Asp (-165.2, -39.1)
- [15] 84 Ile (-25.0, -71.0)
- [15] 102 Glu (-47.3, -91.9)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 16









77.2% (88/114) of all residues were in favored (98%) regions.
95.6% (109/114) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[16] 16 Leu (-39.3, -21.7)
[16] 54 Glu (-69.3, -104.6)
[16] 55 Gln (-160.6, -64.6)
[16] 62 Arg (-69.2, 25.2)
[16] 89 Ser (95.7, 176.0)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 17









78.1% (89/114) of all residues were in favored (98%) regions.
94.7% (108/114) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

[17] 15 Val (-69.9, 63.9)
[17] 55 Gln (-167.6, -50.7)
[17] 70 His (24.1, 69.4)
[17] 71 Gly (-173.2, -28.2)
[17] 84 Ile (-25.6, -68.7)
[17] 102 Glu (-69.9, -101.1)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 18









74.6% (85/114) of all residues were in favored (98%) regions.
93.9% (107/114) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[18] 15 Val (-56.5, 65.8)
[18] 16 Leu (-29.5, -43.5)
[18] 54 Glu (-75.7, -96.1)
[18] 55 Gln (-159.0, -54.6)
[18] 91 Ile (-155.3, -32.3)
[18] 103 Thr (69.4, 116.7)
[18] 113 Arg (-66.5, 30.3)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 19









82.5% (94/114) of all residues were in favored (98%) regions.
96.5% (110/114) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[19] 15 Val (-61.9, 64.4)
[19] 55 Gln (-162.1, -54.1)
[19] 63 Ser (35.6, 88.9)
[19] 102 Glu (-81.5, -101.2)

MolProbity Ramachandran analysis

2gdt.H.pdb, model 20



74.6% (85/114) of all residues were in favored (98%) regions.
93.9% (107/114) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [20] 15 Val (-57.5, 67.1)
- [20] 54 Glu (-62.2, -86.4)
- [20] 55 Gln (-161.6, -62.6)
- [20] 62 Arg (-69.6, 38.8)
- [20] 69 Asn (-32.0, -73.5)
- [20] 84 Ile (-23.2, -83.5)
- [20] 103 Thr (65.6, 98.0)