




7pku.H.pdb, all models









[1] A 31 Ile (57.4, 12.9)	[2] A 26 Glu (60.7, -80.6)	[4] B 239 Gln (62.1, 118.0)
[1] A 94 Gly (164.5, 54.8)	[2] B 211 Ala (62.7, 124.0)	[4] B 245 Thr (54.1, 111.6)
[1] B 107 Tyr (-127.3, -45.4)	[2] B 257 Lys (-126.0, -49.7)	[4] B 257 Lys (-153.4, -48.5)
[1] B 196 Asn (64.6, 143.6)	[3] A 17 Gly (-172.7, 8.2)	[5] A 27 Leu (61.1, 157.1)
[1] B 197 Ser (63.0, 124.4)	[3] A 26 Glu (47.6, -74.1)	[5] A 76 Gly (-170.9, -76.4)
[1] B 214 Gly (-174.0, -55.4)	[3] A 31 Ile (57.3, -1.9)	[5] A 99 Ala (56.5, 174.0)
[1] B 217 Ala (63.4, 151.6)	[3] A 94 Gly (177.9, 52.5)	[5] B 197 Ser (62.6, 125.7)
[1] B 236 Gly (-171.9, -57.6)	[3] A 110 Asp (63.6, 146.4)	[5] B 243 Gly (145.9, 55.2)

[5] B 259 Arg (62.3, 123.9)	[7] A 26 Glu (43.9, 127.6)	[8] A 27 Leu (61.2, 133.9)
[6] A 17 Gly (139.1, 69.4)	[7] A 67 Pro (-35.2, -29.4)	[8] A 95 Glu (-168.9, -42.2)
[6] A 28 Asp (-163.2, -47.6)	[7] A 107 Thr (-134.3, -51.4)	[8] B 197 Ser (63.0, 126.6)
[6] A 31 Ile (56.9, 9.1)	[7] B 194 Ser (62.0, 119.7)	[8] B 210 Met (58.3, 111.6)
[6] A 86 Thr (55.3, 115.6)	[7] B 200 Gly (168.9, 87.9)	[8] B 211 Ala (61.8, 123.7)
[6] B 193 Ser (63.5, 132.3)	[7] B 203 Arg (62.8, 126.2)	[8] B 217 Ala (63.4, 124.2)
[6] B 198 Thr (57.2, 157.7)	[7] B 216 Asp (63.1, 127.6)	[8] B 237 Lys (63.0, 123.9)
[6] B 203 Arg (63.1, 126.8)	[7] B 232 Ser (53.2, 112.9)	[8] B 245 Thr (51.7, 6.3)
[6] B 204 Gly (-177.6, -54.9)	[7] B 233 Lys (62.2, 152.5)	[9] A 28 Asp (61.4, 122.0)
[6] B 211 Ala (62.6, 125.5)	[7] B 236 Gly (-148.1, -89.0)	[9] A 31 Ile (52.9, 13.7)
[6] B 212 Gly (-174.3, 75.0)	[8] A 17 Gly (-142.9, -71.1)	[9] A 67 Pro (-54.0, 1.2)
[6] B 245 Thr (55.8, 121.2)	[8] A 26 Glu (62.9, -82.9)	[9] B 193 Ser (-58.8, 112.3)

MolProbity Ramachandran analysis

7pku.H.pdb, model 1









79.4% (131/165) of all residues were in favored (98%) regions.
95.2% (157/165) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [1] A 31 Ile (57.4, 12.9)
- [1] A 94 Gly (164.5, 54.8)
- [1] A 107 Tyr (-127.3, -45.4)
- [1] B 196 Asn (64.6, 143.6)
- [1] B 197 Ser (63.0, 124.4)
- [1] B 214 Gly (-174.0, -55.4)
- [1] B 217 Ala (63.4, 151.6)
- [1] B 236 Gly (-171.9, -57.6)

MolProbity Ramachandran analysis

7pku.H.pdb, model 2









83.6% (138/165) of all residues were in favored (98%) regions.
98.2% (162/165) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):
[2] A 26 Glu (60.7, -80.6)
[2] B 211 Ala (62.7, 124.0)
[2] B 257 Lys (-126.0, -49.7)

MolProbity Ramachandran analysis

7pku.H.pdb, model 3









83.0% (137/165) of all residues were in favored (98%) regions.
95.8% (158/165) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [3] A 17 Gly (-172.7, 8.2)
- [3] A 26 Glu (47.6, -74.1)
- [3] A 31 Ile (57.3, -1.9)
- [3] A 94 Gly (177.9, 52.5)
- [3] A 110 Asp (63.6, 146.4)
- [3] B 206 Ser (-128.1, -43.8)
- [3] B 257 Lys (103.4, -61.4)

MolProbity Ramachandran analysis

7pku.H.pdb, model 4









82.4% (136/165) of all residues were in favored (98%) regions.
97.0% (160/165) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[4] A 94 Gly (169.6, 53.9)
[4] A 96 Phe (60.5, 93.4)
[4] B 239 Gln (62.1, 118.0)
[4] B 245 Thr (54.1, 111.6)
[4] B 257 Lys (-153.4, -48.5)

MolProbity Ramachandran analysis

7pku.H.pdb, model 5









81.2% (134/165) of all residues were in favored (98%) regions.
96.4% (159/165) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [5] A 27 Leu (61.1, 157.1)
- [5] A 76 Gly (-170.9, -76.4)
- [5] A 99 Ala (56.5, 174.0)
- [5] B 197 Ser (62.6, 125.7)
- [5] B 243 Gly (145.9, 55.2)
- [5] B 259 Arg (62.3, 123.9)

MolProbity Ramachandran analysis

7pku.H.pdb, model 6









80.6% (133/165) of all residues were in favored (98%) regions.
93.3% (154/165) of all residues were in allowed (>99.8%) regions.

There were 11 outliers (phi, psi):

[6] A 17 Gly (139.1, 69.4)	[6] B 211 Ala (62.6, 125.5)
[6] A 28 Asp (-163.2, -47.6)	[6] B 212 Gly (-174.3, 75.0)
[6] A 31 Ile (56.9, 9.1)	[6] B 245 Thr (55.8, 121.2)
[6] A 86 Thr (55.3, 115.6)	
[6] B 193 Ser (63.5, 132.3)	
[6] B 198 Thr (57.2, 157.7)	
[6] B 203 Arg (63.1, 126.8)	
[6] B 204 Gly (-177.6, -54.9)	

MolProbity Ramachandran analysis

7pku.H.pdb, model 7









78.2% (129/165) of all residues were in favored (98%) regions.
93.9% (155/165) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[7] A 26 Glu (43.9, 127.6)	[7] B 233 Lys (62.2, 152.5)
[7] A 67 Pro (-35.2, -29.4)	[7] B 236 Gly (-148.1, -89.0)
[7] A 107 Tyr (-134.3, -51.4)	
[7] B 194 Ser (62.0, 119.7)	
[7] B 200 Gly (168.9, 87.9)	
[7] B 203 Arg (62.8, 126.2)	
[7] B 216 Asp (63.1, 127.6)	
[7] B 232 Ser (53.2, 112.9)	

MolProbity Ramachandran analysis

7pku.H.pdb, model 8









81.8% (135/165) of all residues were in favored (98%) regions.
93.9% (155/165) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[8] A 17 Gly (-142.9, -71.1)	[8] B 237 Lys (63.0, 123.9)
[8] A 26 Glu (62.9, -82.9)	[8] B 245 Thr (51.7, 6.3)
[8] A 27 Leu (61.2, 133.9)	
[8] A 95 Glu (-168.9, -42.2)	
[8] B 197 Ser (63.0, 126.6)	
[8] B 210 Met (58.3, 111.6)	
[8] B 211 Ala (61.8, 123.7)	
[8] B 217 Ala (63.4, 124.2)	

MolProbity Ramachandran analysis

7pku.H.pdb, model 9









80.0% (132/165) of all residues were in favored (98%) regions.
96.4% (159/165) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [9] A 28 Asp (61.4, 122.0)
- [9] A 31 Ile (52.9, 13.7)
- [9] A 67 Pro (-54.0, 1.2)
- [9] B 193 Ser (58.8, 112.3)
- [9] B 242 Gln (64.1, 126.3)
- [9] B 257 Lys (157.7, -50.9)

MolProbity Ramachandran analysis

7pku.H.pdb, model 10



78.8% (130/165) of all residues were in favored (98%) regions.
95.8% (158/165) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[10] A 26 Glu (55.1, -71.5)
[10] B 206 Ser (-127.8, -43.1)
[10] B 216 Asp (62.3, 118.4)
[10] B 235 Ser (54.1, 113.0)
[10] B 237 Lys (63.2, 126.0)
[10] B 238 Gly (-173.3, -54.7)
[10] B 243 Gly (-179.7, 55.8)