


MolProbity Ramachandran analysis

2gri.H.pdb, all models



76.3% (1679/2200) of all residues were in favored (98%) regions.
94.3% (2074/2200) of all residues were in allowed (>99.8%) regions.
This list is truncated; use the MolProbity multi-chart.html for complete list.


There were 126 outliers (phi, psi):

[1] 4 Lys (-58.7, -171.8)	[2] 5 Gly (47.1, -99.0)	[3] 27 Glu (179.1, 110.8)
[1] 5 Gly (-48.6, -81.8)	[2] 40 Cys (-178.3, 92.8)	[3] 40 Cys (-175.9, 80.7)
[1] 11 Asp (-171.0, -54.6)	[2] 46 Glu (-61.0, -169.7)	[3] 46 Glu (-60.9, -170.0)
[1] 40 Cys (-179.8, 87.0)	[2] 63 Val (-55.3, -73.9)	[3] 63 Val (-64.0, -75.4)
[1] 46 Glu (-59.3, -174.4)	[2] 69 Val (-153.4, 66.7)	[3] 69 Val (-152.7, 61.3)
[1] 69 Val (-153.6, 68.3)	[2] 111 Asp (179.7, -33.8)	[3] 111 Asp (-179.2, -34.4)
[1] 111 Asp (179.9, -34.1)	[3] 5 Gly (50.5, -99.4)	[4] 28 Leu (-161.7, -56.3)
[2] 4 Lys (-48.8, 103.0)	[3] 11 Asp (-166.4, -51.5)	[4] 40 Cys (-179.8, 89.1)
		[4] 46 Glu (-57.5, -179.4)
		[4] 69 Val (-153.4, 67.5)
		[4] 111 Asp (-179.8, -34.2)
		[4] 69 Val (-157.0, 62.3)
		[5] 4 Lys (-57.2, -171.7)

[5] 5 Gly (-46.2, -84.1)	[7] 16 Val (-157.6, -34.9)	[9] 69 Val (-154.7, 65.8)
[5] 9 Gly (-44.5, 100.7)	[7] 40 Cys (-178.3, 84.6)	[9] 111 Asp (172.6, -56.3)
[5] 11 Asp (-167.5, -59.8)	[7] 69 Val (-153.5, 63.0)	[10] 7 Thr (67.6, 145.8)
[5] 40 Cys (179.9, 85.2)	[7] 111 Asp (-179.8, -34.2)	[10] 27 Glu (176.5, 123.5)
[5] 46 Glu (-57.4, -179.9)	[8] 4 Lys (41.6, -154.9)	[10] 40 Cys (-179.3, 84.8)
[5] 69 Val (-148.4, 62.9)	[8] 46 Glu (-61.2, -172.3)	[10] 69 Val (-159.0, 68.2)
[5] 111 Asp (-179.8, -34.2)	[8] 69 Val (-154.4, 66.8)	[10] 111 Asp (-179.9, -34.2)
[6] 5 Gly (64.3, -91.2)	[8] 111 Asp (-179.9, -34.1)	[11] 3 Ile (54.2, 179.7)
[6] 6 Val (-168.0, 26.9)	[9] 5 Gly (46.6, -99.9)	[11] 7 Thr (65.5, 149.1)
[6] 40 Cys (-179.9, 86.4)	[9] 11 Asp (-169.7, -65.6)	[11] 27 Glu (174.5, 138.2)
[6] 69 Val (-157.0, 62.3)	[9] 28 Leu (-172.0, -53.7)	[11] 40 Cys (178.7, 94.7)
[6] 111 Asp (-179.9, -34.3)	[9] 40 Cys (-177.1, 86.2)	[11] 69 Val (-159.0, 67.6)

MolProbity Ramachandran analysis

2gri.H.pdb, model 1




75.5% (83/110) of all residues were in favored (98%) regions.
93.6% (103/110) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [1] 4 Lys (-58.7, -171.8)
- [1] 5 Gly (-48.6, -81.8)
- [1] 11 Asp (-171.0, -54.6)
- [1] 40 Cys (-179.8, 87.0)
- [1] 46 Glu (-59.3, -174.4)
- [1] 69 Val (-153.6, 68.3)
- [1] 111 Asp (179.9, -34.1)

MolProbity Ramachandran analysis

2gri.H.pdb, model 2




75.5% (83/110) of all residues were in favored (98%) regions.
93.6% (103/110) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [2] 4 Lys (-48.8, 103.0)
- [2] 5 Gly (47.1, -99.0)
- [2] 40 Cys (-178.3, 92.8)
- [2] 46 Glu (-61.0, -169.7)
- [2] 63 Val (-55.3, -73.9)
- [2] 69 Val (-153.4, 66.7)
- [2] 111 Asp (179.7, -33.8)

MolProbity Ramachandran analysis

2gri.H.pdb, model 3









75.5% (83/110) of all residues were in favored (98%) regions.
92.7% (102/110) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [3] 5 Gly (50.5, -99.4)
- [3] 11 Asp (-166.4, -51.5)
- [3] 27 Glu (179.1, 110.8)
- [3] 40 Cys (-175.9, 80.7)
- [3] 46 Glu (-60.9, -170.0)
- [3] 63 Val (-64.0, -75.4)
- [3] 69 Val (-152.7, 61.3)
- [3] 111 Asp (-179.2, -34.4)

MolProbity Ramachandran analysis

2gri.H.pdb, model 4




75.5% (83/110) of all residues were in favored (98%) regions.
95.5% (105/110) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [4] 28 Leu (-161.7, -56.3)
- [4] 40 Cys (-179.8, 89.1)
- [4] 46 Glu (-57.5, -179.4)
- [4] 69 Val (-153.4, 67.5)
- [4] 111 Asp (-179.8, -34.3)

MolProbity Ramachandran analysis

2gri.H.pdb, model 5









77.3% (85/110) of all residues were in favored (98%) regions.
92.7% (102/110) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [5] 4 Lys (-57.2, -171.7)
- [5] 5 Gly (-46.2, -84.1)
- [5] 9 Gly (-44.5, 100.7)
- [5] 11 Asp (-167.5, -59.8)
- [5] 40 Cys (179.9, 85.2)
- [5] 46 Glu (-57.4, -179.9)
- [5] 69 Val (-148.4, 62.9)
- [5] 111 Asp (-179.8, -34.2)

MolProbity Ramachandran analysis

2gri.H.pdb, model 6









76.4% (84/110) of all residues were in favored (98%) regions.
95.5% (105/110) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [6] 5 Gly (64.3, -91.2)
- [6] 6 Val (-168.0, 26.9)
- [6] 40 Cys (-179.9, 86.4)
- [6] 69 Val (-157.0, 62.3)
- [6] 111 Asp (-179.9, -34.3)

MolProbity Ramachandran analysis

2gri.H.pdb, model 7








79.1% (87/110) of all residues were in favored (98%) regions.
96.4% (106/110) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [7] 16 Val (-157.6, -34.9)
- [7] 40 Cys (-178.3, 84.6)
- [7] 69 Val (-153.5, 63.0)
- [7] 111 Asp (-179.8, -34.2)

MolProbity Ramachandran analysis

2gri.H.pdb, model 8




76.4% (84/110) of all residues were in favored (98%) regions.
96.4% (106/110) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [8] 4 Lys (41.6, -154.9)
- [8] 46 Glu (-61.2, -172.3)
- [8] 69 Val (-154.4, 66.8)
- [8] 111 Asp (-179.9, -34.1)

MolProbity Ramachandran analysis

2gri.H.pdb, model 9




75.5% (83/110) of all residues were in favored (98%) regions.
94.5% (104/110) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [9] 5 Gly (46.6, -99.9)
- [9] 11 Asp (-169.7, -65.6)
- [9] 28 Leu (-172.0, -53.7)
- [9] 40 Cys (-177.1, 86.2)
- [9] 69 Val (-154.7, 65.8)
- [9] 111 Asp (172.6, -56.3)

MolProbity Ramachandran analysis

2gri.H.pdb, model 10




75.5% (83/110) of all residues were in favored (98%) regions.
95.5% (105/110) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[10] 7 Thr (67.6, 145.8)
[10] 27 Glu (176.5, 123.5)
[10] 40 Cys (-179.3, 84.8)
[10] 69 Val (-159.0, 68.2)
[10] 111 Asp (-179.9, -34.2)

MolProbity Ramachandran analysis

2gri.H.pdb, model 11




79.1% (87/110) of all residues were in favored (98%) regions.
94.5% (104/110) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [11] 3 Ile (54.2, 179.7)
- [11] 7 Thr (65.5, 149.1)
- [11] 27 Glu (174.5, 138.2)
- [11] 40 Cys (178.7, 94.7)
- [11] 69 Val (-159.0, 67.6)
- [11] 111 Asp (-179.8, -34.2)

MolProbity Ramachandran analysis

2gri.H.pdb, model 12




76.4% (84/110) of all residues were in favored (98%) regions.
96.4% (106/110) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [12] 40 Cys (-178.3, 49.0)
- [12] 46 Glu (-62.5, -169.8)
- [12] 69 Val (-148.3, 70.8)
- [12] 111 Asp (178.1, -33.9)

MolProbity Ramachandran analysis

2gri.H.pdb, model 13




75.5% (83/110) of all residues were in favored (98%) regions.
91.8% (101/110) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[13] 4 Lys (-58.6, -172.5)	[13] 111 Asp (178.2, -33.9)
[13] 5 Gly (-50.3, -78.9)	
[13] 11 Asp (-170.9, -65.8)	
[13] 25 Thr (-179.0, 93.7)	
[13] 27 Glu (174.5, 114.3)	
[13] 40 Cys (-179.5, 57.4)	
[13] 46 Glu (-62.3, -169.9)	
[13] 69 Val (-154.9, 65.5)	

MolProbity Ramachandran analysis

2gri.H.pdb, model 14




74.5% (82/110) of all residues were in favored (98%) regions.
92.7% (102/110) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [14] 4 Lys (-59.4, -167.4)
- [14] 5 Gly (-62.2, 92.5)
- [14] 6 Val (53.0, 91.3)
- [14] 11 Asp (-171.4, -63.2)
- [14] 40 Cys (176.1, 87.0)
- [14] 46 Glu (-59.9, -174.2)
- [14] 69 Val (-152.5, 67.3)
- [14] 111 Asp (-179.8, -34.2)

MolProbity Ramachandran analysis

2gri.H.pdb, model 15




76.4% (84/110) of all residues were in favored (98%) regions.
92.7% (102/110) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

[15] 3 Ile (64.7, 154.0)
[15] 4 Lys (-62.3, -162.0)
[15] 5 Gly (-44.9, -85.8)
[15] 9 Gly (-44.6, 101.7)
[15] 40 Cys (179.7, 88.6)
[15] 46 Glu (-58.1, -177.9)
[15] 69 Val (-153.8, 60.0)
[15] 111 Asp (172.6, -56.3)

MolProbity Ramachandran analysis

2gri.H.pdb, model 16




77.3% (85/110) of all residues were in favored (98%) regions.
95.5% (105/110) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[16] 4 Lys (43.0, -156.9)
[16] 27 Glu (179.1, 105.3)
[16] 40 Cys (177.9, 89.9)
[16] 69 Val (-156.0, 66.9)
[16] 111 Asp (180.0, -34.1)

MolProbity Ramachandran analysis

2gri.H.pdb, model 17






74.5% (82/110) of all residues were in favored (98%) regions.
94.5% (104/110) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

[17] 4 Lys (-58.4, -172.3)
[17] 5 Gly (-58.4, 96.5)
[17] 6 Val (51.4, 88.6)
[17] 40 Cys (179.4, 85.0)
[17] 69 Val (-154.6, 60.8)
[17] 111 Asp (172.5, -56.3)

2gri.H.pdb, model 18









There were 6 outliers (phi, psi):

[18]	6 Val (-168.2, 27.1)
[18]	11 Asp (-160.8, -46.8)
[18]	28 Leu (-172.0, -48.1)
[18]	40 Cys (-177.0, 86.3)
[18]	69 Val (-153.9, 53.8)
[18]	111 Asp (-180.0, -34.2)

MolProbity Ramachandran analysis

2gri.H.pdb, model 19









75.5% (83/110) of all residues were in favored (98%) regions.
93.6% (103/110) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [19] 3 Ile (64.6, 153.0)
- [19] 11 Asp (-164.4, -64.9)
- [19] 32 Val (-162.3, 38.2)
- [19] 40 Cys (179.8, 85.4)
- [19] 46 Glu (-54.3, 176.1)
- [19] 69 Val (-158.1, 70.6)
- [19] 111 Asp (-179.8, -34.2)

MolProbity Ramachandran analysis

2gri.H.pdb, model 20



75.5% (83/110) of all residues were in favored (98%) regions.
92.7% (102/110) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

[20] 4 Lys (-58.3, -172.6)
[20] 5 Gly (-59.7, 95.4)
[20] 6 Val (53.8, 91.2)
[20] 11 Asp (-170.7, -66.2)
[20] 28 Leu (-163.1, -54.3)
[20] 40 Cys (-176.6, 84.5)
[20] 69 Val (-156.0, 74.5)
[20] 111 Asp (-179.7, -34.2)