


MolProbity Ramachandran analysis

2k87.H.pdb, all models



86.9% (1981/2280) of all residues were in favored (98%) regions.
 97.0% (2212/2280) of all residues were in allowed (>99.8%) regions.
 This list is truncated; use the MolProbity multi-chart.html for complete list.







There were 68 outliers (phi, psi):

[1] 8 Asp (27.9, 34.4)	[3] 7 Ile (-46.9, 98.4)	[5] 8 Asp (24.8, 42.9)
[1] 17 Asn (49.1, 4.7)	[3] 8 Asp (33.3, 27.1)	[5] 22 Asn (-54.4, -3.3)
[1] 115 Asp (-61.8, 13.3)	[3] 67 Tyr (-45.5, -17.1)	[6] 18 Ala (40.7, 166.5)
[2] 22 Asn (-68.0, 21.3)	[3] 90 Asn (148.3, 40.3)	[6] 112 Lys (41.2, 101.5)
[2] 112 Lys (57.9, 127.0)	[3] 114 Val (48.1, 18.2)	[7] 8 Asp (45.2, 14.5)
[2] 114 Val (85.6, 93.2)	[4] 8 Asp (37.7, 23.5)	[7] 56 Pro (-83.1, 92.4)
[3] 4 Glu (63.7, 95.1)	[4] 112 Lys (57.2, 140.6)	[8] 5 Gln (57.5, -179.4)
[3] 5 Gln (66.8, 128.3)	[5] 5 Gln (53.7, 128.3)	[8] 56 Pro (-79.0, 92.4)
		[8] 112 Lys (58.0, 120.3)
		[9] 56 Pro (-78.1, 94.4)
		[13] 90 Asn (152.2, 45.7)
		[10] 22 Asn (-71.3, 21.3)

[10] 56 Pro (-75.0, 94.3)	[14] 22 Asn (-69.0, 27.7)	[17] 22 Asn (-60.6, 6.7)
[10] 81 Leu (-24.9, -72.1)	[14] 67 Tyr (-44.1, -20.6)	[17] 56 Pro (-78.1, 94.9)
[11] 22 Asn (-53.9, 12.9)	[15] 8 Asp (41.9, 24.4)	[17] 67 Tyr (-34.7, -26.9)
[11] 72 Ala (60.1, -19.4)	[15] 17 Asn (46.5, 9.1)	[17] 72 Ala (59.2, -13.9)
[11] 112 Lys (59.8, 130.3)	[16] 5 Gln (60.7, 165.7)	[17] 112 Lys (-34.6, 95.3)
[11] 114 Val (-154.6, 31.1)	[16] 8 Asp (37.0, 19.5)	[18] 4 Glu (49.9, -165.3)
[12] 8 Asp (38.4, 26.6)	[16] 17 Asn (60.2, -20.1)	[18] 22 Asn (-55.2, 13.8)
[12] 56 Pro (-72.6, 93.8)	[16] 22 Asn (-66.5, 15.3)	[18] 56 Pro (-83.0, 93.8)
[13] 8 Asp (33.1, 33.0)	[16] 56 Pro (-74.6, 94.8)	[19] 8 Asp (35.3, 33.3)
[13] 56 Pro (-72.6, 93.3)	[16] 67 Tyr (-43.6, -22.2)	[19] 17 Asn (45.5, 8.0)
[13] 90 Asn (152.2, 45.7)	[17] 16 Pro (-73.6, -149.2)	[19] 56 Pro (-80.2, 93.0)
[13] 112 Lys (56.7, 139.2)	[17] 17 Asn (-37.4, 36.0)	[19] 90 Asn (143.8, 27.8)

MolProbity Ramachandran analysis

2k87.H.pdb, model 1









89.5% (102/114) of all residues were in favored (98%) regions.
97.4% (111/114) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

- [1] 8 Asp (27.9, 34.4)
- [1] 17 Asn (49.1, 4.7)
- [1] 115 Asp (-61.8, 13.3)

MolProbity Ramachandran analysis

2k87.H.pdb, model 2









90.4% (103/114) of all residues were in favored (98%) regions.
97.4% (111/114) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[2] 22 Asn (-68.0, 21.3)
[2] 112 Lys (57.9, 127.0)
[2] 114 Val (85.6, 93.2)

MolProbity Ramachandran analysis

2k87.H.pdb, model 3









85.1% (97/114) of all residues were in favored (98%) regions.
93.9% (107/114) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [3] 4 Glu (63.7, 95.1)
- [3] 5 Gln (66.8, 128.3)
- [3] 7 Ile (-46.9, 98.4)
- [3] 8 Asp (33.3, 27.1)
- [3] 67 Tyr (-45.5, -17.1)
- [3] 90 Asn (148.3, 40.3)
- [3] 114 Val (48.1, 18.2)

MolProbity Ramachandran analysis

2k87.H.pdb, model 4









88.6% (101/114) of all residues were in favored (98%) regions.
98.2% (112/114) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[4] 8 Asp (37.7, 23.5)
[4] 112 Lys (57.2, 140.6)

MolProbity Ramachandran analysis

2k87.H.pdb, model 5









82.5% (94/114) of all residues were in favored (98%) regions.
97.4% (111/114) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[5] 5 Gln (53.7, 128.3)
[5] 8 Asp (24.8, 42.9)
[5] 22 Asn (-54.4, -3.3)

MolProbity Ramachandran analysis

2k87.H.pdb, model 6









80.7% (92/114) of all residues were in favored (98%) regions.
98.2% (112/114) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):

[6] 18 Ala (40.7, 166.5)
[6] 112 Lys (41.2, 101.5)

MolProbity Ramachandran analysis

2k87.H.pdb, model 7









91.2% (104/114) of all residues were in favored (98%) regions.
98.2% (112/114) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[7] 8 Asp (45.2, 14.5)
[7] 56 Pro (-83.1, 92.4)

MolProbity Ramachandran analysis

2k87.H.pdb, model 8









84.2% (96/114) of all residues were in favored (98%) regions.
97.4% (111/114) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[8] 5 Gln (57.5, -179.4)
[8] 56 Pro (-79.0, 92.4)
[8] 112 Lys (58.0, 120.3)

MolProbity Ramachandran analysis

2k87.H.pdb, model 9









83.3% (95/114) of all residues were in favored (98%) regions.
98.2% (112/114) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[9] 56 Pro (-78.1, 94.4)
[9] 67 Tyr (-38.5, -27.2)

MolProbity Ramachandran analysis

2k87.H.pdb, model 10









86.8% (99/114) of all residues were in favored (98%) regions.
97.4% (111/114) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[10] 22 Asn (-71.3, 21.3)
[10] 56 Pro (-75.0, 94.3)
[10] 81 Leu (-24.9, -72.1)

MolProbity Ramachandran analysis

2k87.H.pdb, model 11









89.5% (102/114) of all residues were in favored (98%) regions.
96.5% (110/114) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[11] 22 Asn (-53.9, 12.9)
[11] 72 Ala (60.1, -19.4)
[11] 112 Lys (59.8, 130.3)
[11] 114 Val (-154.6, 31.1)

MolProbity Ramachandran analysis

2k87.H.pdb, model 12




89.5% (102/114) of all residues were in favored (98%) regions.
98.2% (112/114) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[12] 8 Asp (38.4, 26.6)
[12] 56 Pro (-72.6, 93.8)

MolProbity Ramachandran analysis

2k87.H.pdb, model 13









87.7% (100/114) of all residues were in favored (98%) regions.
96.5% (110/114) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[13] 8 Asp (33.1, 33.0)
[13] 56 Pro (-72.6, 93.3)
[13] 90 Asn (152.2, 45.7)
[13] 112 Lys (56.7, 139.2)

MolProbity Ramachandran analysis

2k87.H.pdb, model 14









88.6% (101/114) of all residues were in favored (98%) regions.
98.2% (112/114) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[14] 22 Asn (-69.0, 27.7)
[14] 67 Tyr (-44.1, -20.6)

MolProbity Ramachandran analysis

2k87.H.pdb, model 15









86.0% (98/114) of all residues were in favored (98%) regions.
98.2% (112/114) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[15] 8 Asp (41.9, 24.4)
[15] 17 Asn (46.5, 9.1)

MolProbity Ramachandran analysis

2k87.H.pdb, model 16









88.6% (101/114) of all residues were in favored (98%) regions.
94.7% (108/114) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

[16] 5 Gln (60.7, 165.7)
[16] 8 Asp (37.0, 19.5)
[16] 17 Asn (60.2, -20.1)
[16] 22 Asn (-66.5, 15.3)
[16] 56 Pro (-74.6, 94.8)
[16] 67 Tyr (-43.6, -22.2)

MolProbity Ramachandran analysis

2k87.H.pdb, model 17




86.0% (98/114) of all residues were in favored (98%) regions.
93.9% (107/114) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [17] 16 Pro (-73.6, -149.2)
- [17] 17 Asn (-37.4, 36.0)
- [17] 22 Asn (-60.6, 6.7)
- [17] 56 Pro (-78.1, 94.9)
- [17] 67 Tyr (-34.7, -26.9)
- [17] 72 Ala (59.2, -13.9)
- [17] 112 Lys (-34.6, 95.3)

MolProbity Ramachandran analysis

2k87.H.pdb, model 18









85.1% (97/114) of all residues were in favored (98%) regions.
97.4% (111/114) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[18] 4 Glu (49.9, -165.3)
[18] 22 Asn (-55.2, 13.8)
[18] 56 Pro (-83.0, 93.8)

MolProbity Ramachandran analysis

2k87.H.pdb, model 19









87.7% (100/114) of all residues were in favored (98%) regions.
95.6% (109/114) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[19] 8 Asp (35.3, 33.3)
[19] 17 Asn (45.5, 8.0)
[19] 56 Pro (-80.2, 93.0)
[19] 90 Asn (143.8, 27.8)
[19] 114 Val (45.0, 22.8)

MolProbity Ramachandran analysis

2k87.H.pdb, model 20



86.8% (99/114) of all residues were in favored (98%) regions.
97.4% (111/114) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[20] 2 Tyr (-73.8, -95.1)
[20] 17 Asn (52.8, -5.5)
[20] 56 Pro (-72.2, 94.7)