







MolProbity Ramachandran analysis

2kaf.H.pdb, all models



91.2% (1186/1300) of all residues were in favored (98%) regions.
97.5% (1268/1300) of all residues were in allowed (>99.8%) regions.







There were 32 outliers (phi, psi):

[1] 17 Arg (51.2, -82.0)	[6] 17 Arg (54.7, -85.4)	[12] 17 Arg (60.6, -84.2)
[2] 17 Arg (53.8, -81.1)	[7] 17 Arg (56.6, -85.3)	[13] 17 Arg (56.9, -83.6)
[2] 32 Phe (-81.8, -112.8)	[7] 32 Phe (-85.3, -112.9)	[13] 32 Phe (-80.9, -112.7)
[3] 17 Arg (57.8, -78.3)	[8] 17 Arg (59.7, -85.4)	[13] 54 Gly (173.9, -9.4)
[4] 17 Arg (50.8, -85.8)	[8] 32 Phe (-85.6, -112.9)	[14] 17 Arg (62.9, -82.4)
[4] 36 Gly (-45.0, 105.4)	[9] 17 Arg (55.3, -85.0)	[15] 17 Arg (53.7, -85.7)
[4] 54 Gly (-176.4, 17.8)	[10] 17 Arg (54.7, -86.8)	[16] 17 Arg (59.9, -82.5)
[5] 17 Arg (45.9, -87.0)	[11] 17 Arg (52.3, -87.0)	[16] 32 Phe (-83.3, -112.7)
		[16] 54 Gly (-178.1, 9.6)
		[17] 17 Arg (57.1, -84.5)
		[17] 54 Gly (163.9, -9.8)

[18] 17 Arg (52.9, -85.9)
[19] 17 Arg (54.6, -85.6)
[20] 32 Phe (-80.4, -111.8)
[20] 36 Gly (-56.3, 98.2)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 1



92.3% (60/65) of all residues were in favored (98%) regions.







98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):

[1] 17 Arg (51.2, -82.0)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 2









93.8% (61/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[2] 17 Arg (53.8, -81.1)
[2] 32 Phe (-81.8, -112.8)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 3



92.3% (60/65) of all residues were in favored (98%) regions.







98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):

[3] 17 Arg (57.8, -78.3)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 4









89.2% (58/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

- [4] 17 Arg (50.8, -85.8)
- [4] 36 Gly (-45.0, 105.4)
- [4] 54 Gly (-176.4, 17.8)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 5









93.8% (61/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[5] 17 Arg (45.9, -87.0)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 6









92.3% (60/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[6] 17 Arg (54.7, -85.4)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 7









92.3% (60/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[7] 17 Arg (56.6, -85.3)
[7] 32 Phe (-85.3, -112.9)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 8









92.3% (60/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[8] 17 Arg (59.7, -85.4)
[8] 32 Phe (-85.6, -112.9)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 9









93.8% (61/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[9] 17 Arg (55.3, -85.0)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 10









92.3% (60/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[10] 17 Arg (54.7, -86.8)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 11









87.7% (57/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[11] 17 Arg (52.3, -87.0)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 12



93.8% (61/65) of all residues were in favored (98%) regions.







98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):

[12] 17 Arg (60.6, -84.2)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 13









90.8% (59/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[13] 17 Arg (56.9, -83.6)
[13] 32 Phe (-80.9, -112.7)
[13] 54 Gly (173.9, -9.4)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 14









87.7% (57/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[14] 17 Arg (62.9, -82.4)
[14] 54 Gly (-166.8, 10.5)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 15









89.2% (58/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[15] 17 Arg (53.7, -85.7)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 16









89.2% (58/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

- [16] 17 Arg (59.9, -82.5)
- [16] 32 Phe (-83.3, -112.7)
- [16] 54 Gly (-178.1, 9.6)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 17









87.7% (57/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[17] 17 Arg (57.1, -84.5)
[17] 54 Gly (163.9, -9.8)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 18









90.8% (59/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[18] 17 Arg (52.9, -85.9)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 19









93.8% (61/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[19] 17 Arg (54.6, -85.6)

MolProbity Ramachandran analysis

2kaf.H.pdb, model 20



89.2% (58/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[20] 32 Phe (-80.4, -111.8)
[20] 36 Gly (-56.3, 98.2)