




2kqv.H.pdb, all models









[13] 114 Ser (-74.2, -84.0)
[19] 92 Gly (143.7, 82.4)
[20] 4 Met (57.8, 172.2)
[20] 128 Ser (55.2, -176.9)

[2] 127 Thr (178.3, -54.0)	[5] 53 Asp (-65.9, 14.3)
[3] 60 Phe (44.1, 153.3)	[5] 60 Phe (52.3, -173.4)
[4] 4 Met (-57.1, -79.3)	[5] 127 Thr (-165.2, -60.8)
[4] 56 Val (-149.7, 51.8)	[6] 127 Thr (174.0, -42.4)
[4] 60 Phe (46.1, 175.9)	[6] 60 Phe (44.5, 174.4)
[4] 86 Ile (-75.5, -79.9)	[7] 78 Asn (59.1, -12.1)
[5] 3 His (56.4, -66.2)	[8] 3 His (65.7, 100.7)
[5] 4 Met (45.9, -169.3)	[8] 6 Thr (49.8, 162.1)

MolProbity Ramachandran analysis

2kqv.H.pdb, model 1









87.4% (111/127) of all residues were in favored (98%) regions.
100.0% (127/127) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2kqv.H.pdb, model 2









86.6% (110/127) of all residues were in favored (98%) regions.
99.2% (126/127) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[2] 127 Thr (178.3, -54.0)

MolProbity Ramachandran analysis

2kqv.H.pdb, model 3









89.8% (114/127) of all residues were in favored (98%) regions.
99.2% (126/127) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[3] 60 Phe (44.1, 153.3)

MolProbity Ramachandran analysis

2kqv.H.pdb, model 4









88.2% (112/127) of all residues were in favored (98%) regions.
96.9% (123/127) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [4] 4 Met (-57.1, -79.3)
- [4] 56 Val (-149.7, 51.8)
- [4] 60 Phe (46.1, 175.9)
- [4] 86 Ile (-75.5, -79.9)

MolProbity Ramachandran analysis

2kqv.H.pdb, model 5




88.2% (112/127) of all residues were in favored (98%) regions.
96.1% (122/127) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [5] 3 His (56.4, -66.2)
- [5] 4 Met (45.9, -169.3)
- [5] 53 Asp (-65.9, 14.3)
- [5] 60 Phe (52.3, -173.4)
- [5] 127 Thr (-165.2, -60.8)

MolProbity Ramachandran analysis

2kqv.H.pdb, model 6









89.8% (114/127) of all residues were in favored (98%) regions.
99.2% (126/127) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[6] 127 Thr (174.0, -42.4)

MolProbity Ramachandran analysis




2kqv.H.pdb, model 7



87.4% (111/127) of all residues were in favored (98%) regions.
98.4% (125/127) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[7] 60 Phe (44.5, 174.4)
[7] 78 Asn (59.1, -12.1)

2kqv.H.pdb, model 8









There were 2 outliers (phi, psi):

```
[8] 3 His (65.7, 100.7)
[8] 6 Thr (49.8, 162.1)
```

MolProbity Ramachandran analysis

2kqv.H.pdb, model 9









82.7% (105/127) of all residues were in favored (98%) regions.
97.6% (124/127) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[9] 4 Met (44.9, 175.6)
[9] 6 Thr (64.6, 129.6)
[9] 60 Phe (47.6, -176.0)

MolProbity Ramachandran analysis

2kqv.H.pdb, model 10









82.7% (105/127) of all residues were in favored (98%) regions.
98.4% (125/127) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[10] 114 Ser (-74.4, -85.5)
[10] 127 Thr (-176.9, 93.1)

MolProbity Ramachandran analysis

2kqv.H.pdb, model 11









87.4% (111/127) of all residues were in favored (98%) regions.
99.2% (126/127) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[11] 4 Met (48.2, -163.0)

MolProbity Ramachandran analysis

2kqv.H.pdb, model 12









85.0% (108/127) of all residues were in favored (98%) regions.
96.9% (123/127) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[12] 7 Val (-150.4, -165.9)
[12] 60 Phe (46.0, 167.6)
[12] 86 Ile (-61.2, -77.1)
[12] 128 Ser (50.9, -169.9)

MolProbity Ramachandran analysis

2kqv.H.pdb, model 13









87.4% (111/127) of all residues were in favored (98%) regions.
97.6% (124/127) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[13] 4 Met (45.1, -171.0)
[13] 57 Arg (-55.0, 94.2)
[13] 114 Ser (-74.2, -84.0)

MolProbity Ramachandran analysis

2kqv.H.pdb, model 14









85.8% (109/127) of all residues were in favored (98%) regions.
100.0% (127/127) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2kqv.H.pdb, model 15








87.4% (111/127) of all residues were in favored (98%) regions.
100.0% (127/127) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2kqv.H.pdb, model 16









90.6% (115/127) of all residues were in favored (98%) regions.
100.0% (127/127) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2kqv.H.pdb, model 17









85.8% (109/127) of all residues were in favored (98%) regions.
100.0% (127/127) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2kqv.H.pdb, model 18









82.7% (105/127) of all residues were in favored (98%) regions.
100.0% (127/127) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

2kqv.H.pdb, model 19









86.6% (110/127) of all residues were in favored (98%) regions.
99.2% (126/127) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[19] 92 Gly (143.7, 82.4)

MolProbity Ramachandran analysis

2kqv.H.pdb, model 20



87.4% (111/127) of all residues were in favored (98%) regions.
98.4% (125/127) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[20] 4 Met (57.8, 172.2)
[20] 128 Ser (55.2, -176.9)