







MolProbity Ramachandran analysis

2kqw.H.pdb, all models



84.5% (1099/1300) of all residues were in favored (98%) regions.
96.4% (1253/1300) of all residues were in allowed (>99.8%) regions.







There were 47 outliers (phi, psi):

[1] 177 Ser (150.5, -44.3)	[6] 177 Ser (157.9, -43.6)	[9] 179 Val (45.1, 128.8)
[2] 177 Ser (146.2, -50.6)	[6] 179 Val (52.3, 81.3)	[10] 148 Arg (-53.8, -76.3)
[3] 177 Ser (151.6, -42.0)	[7] 177 Ser (153.7, -44.1)	[10] 177 Ser (155.8, -41.7)
[3] 179 Val (43.4, 83.9)	[8] 174 Thr (43.5, -165.0)	[10] 179 Val (48.3, 135.5)
[4] 174 Thr (43.2, -164.8)	[8] 177 Ser (151.8, -47.3)	[11] 177 Ser (164.1, -48.6)
[4] 177 Ser (150.7, -39.8)	[8] 179 Val (65.0, 93.8)	[11] 179 Val (58.1, 84.4)
[4] 179 Val (50.6, 115.2)	[9] 153 Ser (-163.4, -23.5)	[12] 168 Asp (54.3, -20.3)
[5] 177 Ser (157.9, -40.7)	[9] 177 Ser (158.2, -41.0)	[12] 177 Ser (156.8, -33.2)
		[12] 179 Val (45.4, 119.5)
		[13] 177 Ser (158.1, -36.5)
		[13] 179 Val (44.3, 124.0)
		[14] 174 Thr (44.9, -163.7)

[14] 177 Ser (163.7, -47.1)	[18] 189 Ser (-60.2, -172.8)
[14] 179 Val (45.7, 141.0)	[19] 160 Gly (168.4, -86.2)
[15] 134 Glu (-51.6, -76.9)	[19] 177 Ser (154.3, -50.8)
[15] 174 Thr (42.8, -167.3)	[19] 185 Gly (163.3, -4.6)
[15] 177 Ser (150.3, -41.7)	[20] 148 Arg (-44.7, -78.4)
[15] 179 Val (59.5, 89.5)	[20] 177 Ser (160.6, -45.9)
[16] 177 Ser (161.3, -46.7)	[20] 179 Val (62.6, 100.2)
[16] 179 Val (51.2, 85.1)	
[17] 177 Ser (153.2, -43.9)	
[17] 179 Val (59.7, 95.5)	
[18] 177 Ser (155.0, -39.8)	
[18] 179 Val (44.2, 118.3)	

MolProbity Ramachandran analysis

2kqw.H.pdb, model 1









89.2% (58/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[1] 177 Ser (150.5, -44.3)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 2









87.7% (57/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[2] 177 Ser (146.2, -50.6)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 3









78.5% (51/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):

[3] 177 Ser (151.6, -42.0)
[3] 179 Val (43.4, 83.9)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 4









83.1% (54/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[4] 174 Thr (43.2, -164.8)
[4] 177 Ser (150.7, -39.8)
[4] 179 Val (50.6, 115.2)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 5









84.6% (55/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[5] 177 Ser (157.9, -40.7)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 6









86.2% (56/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):

[6] 177 Ser (157.9, -43.6)
[6] 179 Val (52.3, 81.3)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 7









83.1% (54/65) of all residues were in favored (98%) regions.
98.5% (64/65) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[7] 177 Ser (153.7, -44.1)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 8









78.5% (51/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[8] 174 Thr (43.5, -165.0)
[8] 177 Ser (151.8, -47.3)
[8] 179 Val (65.0, 93.8)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 9









87.7% (57/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[9] 153 Ser (-163.4, -23.5)
[9] 177 Ser (158.2, -41.0)
[9] 179 Val (45.1, 128.8)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 10









87.7% (57/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):
[10] 148 Arg (-53.8, -76.3)
[10] 177 Ser (155.8, -41.7)
[10] 179 Val (48.3, 135.5)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 11









84.6% (55/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[11] 177 Ser (164.1, -48.6)
[11] 179 Val (58.1, 84.4)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 12









80.0% (52/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[12] 168 Asp (54.3, -20.3)
[12] 177 Ser (156.8, -33.2)
[12] 179 Val (45.4, 119.5)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 13









81.5% (53/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[13] 177 Ser (158.1, -36.5)
[13] 179 Val (44.3, 124.0)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 14









84.6% (55/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[14] 174 Thr (44.9, -163.7)
[14] 177 Ser (163.7, -47.1)
[14] 179 Val (45.7, 141.0)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 15



84.6% (55/65) of all residues were in favored (98%) regions.

93.8% (61/65) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[15] 134 Glu (-51.6, -76.9)







[15] 174 Thr (42.8, -167.3)

[15] 177 Ser (150.3, -41.7)

[15] 179 Val (59.5, 89.5)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 16









86.2% (56/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[16] 177 Ser (161.3, -46.7)
[16] 179 Val (51.2, 85.1)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 17




90.8% (59/65) of all residues were in favored (98%) regions.
96.9% (63/65) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[17] 177 Ser (153.2, -43.9)
[17] 179 Val (59.7, 95.5)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 18



87.7% (57/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):







[18] 177 Ser (155.0, -39.8)

[18] 179 Val (44.2, 118.3)

[18] 189 Ser (-60.2, -172.8)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 19









80.0% (52/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[19] 160 Gly (168.4, -86.2)
[19] 177 Ser (154.3, -50.8)
[19] 185 Gly (163.3, -4.6)

MolProbity Ramachandran analysis

2kqw.H.pdb, model 20



84.6% (55/65) of all residues were in favored (98%) regions.
95.4% (62/65) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[20] 148 Arg (-44.7, -78.4)
[20] 177 Ser (160.6, -45.9)
[20] 179 Val (62.6, 100.2)