


MolProbity Ramachandran analysis

2rnk.H.pdb, all models



83.0% (2340/2820) of all residues were in favored (98%) regions.
97.3% (2745/2820) of all residues were in allowed (>99.8%) regions.
This list is truncated; use the MolProbity multi-chart.html for complete list.

There were 75 outliers (phi, psi):







[1] 569 Ile (51.0, 142.5)	[2] 608 Ile (29.3, -89.0)	[6] -2 His (53.8, 91.5)
[1] 608 Ile (34.1, -88.4)	[3] 569 Ile (56.4, 142.5)	[6] 513 Val (44.9, -164.6)
[2] -1 Met (50.7, 86.4)	[3] 608 Ile (45.3, -92.0)	[6] 517 Glu (159.4, 166.9)
[2] 517 Glu (59.2, -64.3)	[4] -3 Ser (95.0, 117.4)	[6] 518 Ala (154.8, 104.5)
[2] 565 Lys (41.1, -157.4)	[4] 523 Glu (47.6, -167.1)	[7] 523 Glu (52.6, -168.9)
[2] 566 Gly (-44.3, 42.9)	[4] 565 Lys (-27.6, -30.9)	[7] 569 Ile (53.1, 144.7)
[2] 572 Gly (61.9, 109.6)	[5] 608 Ile (24.1, -90.9)	[7] 608 Ile (54.7, -80.6)
[2] 573 Ile (56.3, 103.6)	[5] 612 Thr (-55.2, -3.2)	[8] 513 Val (77.0, 106.7)
		[8] 569 Ile (56.5, 137.4)
		[8] 608 Ile (35.6, -90.7)
		[9] 544 Arg (31.0, 42.1)

[9] 569 Ile (57.8, 135.8)	[12] 569 Ile (54.9, 145.3)	[16] 517 Glu (36.3, 107.8)
[9] 608 Ile (30.4, -98.4)	[12] 608 Ile (43.3, -94.0)	[16] 569 Ile (46.7, 114.1)
[10] 517 Glu (47.3, 101.7)	[13] -1 Met (46.1, 80.8)	[16] 572 Gly (71.1, 107.5)
[10] 569 Ile (44.8, 132.5)	[13] 522 Lys (54.6, 118.8)	[16] 573 Ile (41.2, 101.0)
[10] 650 Ser (47.9, 85.1)	[13] 569 Ile (49.8, 141.9)	[17] -2 His (56.0, -179.9)
[11] 513 Val (57.8, 179.0)	[13] 608 Ile (26.9, -94.8)	[17] -1 Met (164.9, -31.6)
[11] 532 Asn (-155.9, -95.9)	[13] 611 Val (-147.7, -62.9)	[17] 523 Glu (41.3, -161.6)
[11] 569 Ile (54.4, 138.6)	[14] 516 Ser (55.4, 167.2)	[17] 532 Asn (-170.5, -105.4)
[11] 608 Ile (29.8, -94.6)	[14] 569 Ile (-37.2, 145.5)	[17] 569 Ile (52.2, 146.4)
[12] 513 Val (61.5, 161.9)	[15] 523 Glu (44.4, -158.1)	[17] 578 Val (-79.4, 31.2)
[12] 520 Asn (96.6, 136.8)	[15] 608 Ile (-42.1, -76.6)	[17] 579 Arg (24.2, 71.4)
[12] 532 Asn (-164.3, -86.5)	[15] 650 Ser (63.8, 102.5)	[17] 608 Ile (23.8, -90.6)

[12] 569 Ile (54.9, 145.3)	[16] 517 Glu (36.3, 107.8)
[12] 608 Ile (43.3, -94.0)	[16] 569 Ile (46.7, 114.1)
[13] -1 Met (46.1, 80.8)	[16] 572 Gly (71.1, 107.5)
[13] 522 Lys (54.6, 118.8)	[16] 573 Ile (41.2, 101.0)
[13] 569 Ile (49.8, 141.9)	[17] -2 His (56.0, -179.9)
[13] 608 Ile (26.9, -94.8)	[17] -1 Met (164.9, -31.6)
[13] 611 Val (-147.7, -62.9)	[17] 523 Glu (41.3, -161.6)
[14] 516 Ser (55.4, 167.2)	[17] 532 Asn (-170.5, -105.4)
[14] 569 Ile (-37.2, 145.5)	[17] 569 Ile (52.2, 146.4)
[15] 523 Glu (44.4, -158.1)	[17] 578 Val (-79.4, 31.2)
[15] 608 Ile (-42.1, -76.6)	[17] 579 Arg (24.2, 71.4)
[15] 650 Ser (63.8, 102.5)	[17] 608 Ile (23.8, -90.6)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 1









86.5% (122/141) of all residues were in favored (98%) regions.
98.6% (139/141) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[1] 569 Ile (51.0, 142.5)
[1] 608 Ile (34.1, -88.4)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 2









80.1% (113/141) of all residues were in favored (98%) regions.
95.0% (134/141) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [2] -1 Met (50.7, 86.4)
- [2] 517 Glu (59.2, -64.3)
- [2] 565 Lys (41.1, -157.4)
- [2] 566 Gly (-44.3, 42.9)
- [2] 572 Gly (61.9, 109.6)
- [2] 573 Ile (56.3, 103.6)
- [2] 608 Ile (29.3, -89.0)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 3









83.0% (117/141) of all residues were in favored (98%) regions.
98.6% (139/141) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[3] 569 Ile (56.4, 142.5)
[3] 608 Ile (45.3, -92.0)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 4









80.1% (113/141) of all residues were in favored (98%) regions.
97.9% (138/141) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

- [4] -3 Ser (95.0, 117.4)
- [4] 523 Glu (47.6, -167.1)
- [4] 565 Lys (-27.6, -30.9)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 5









83.7% (118/141) of all residues were in favored (98%) regions.
98.6% (139/141) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[5] 608 Ile (24.1, -90.9)
[5] 612 Thr (-55.2, -3.2)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 6









79.4% (112/141) of all residues were in favored (98%) regions.
96.5% (136/141) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [6] -2 His (53.8, 91.5)
- [6] 513 Val (44.9, -164.6)
- [6] 517 Glu (159.4, 166.9)
- [6] 518 Ala (154.8, 104.5)
- [6] 608 Ile (-45.2, -69.0)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 7








82.3% (116/141) of all residues were in favored (98%) regions.
97.9% (138/141) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[7] 523 Glu (52.6, -168.9)
[7] 569 Ile (53.1, 144.7)
[7] 608 Ile (54.7, -80.6)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 8









80.9% (114/141) of all residues were in favored (98%) regions.
97.9% (138/141) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[8] 513 Val (77.0, 106.7)
[8] 569 Ile (56.5, 137.4)
[8] 608 Ile (35.6, -90.7)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 9









83.0% (117/141) of all residues were in favored (98%) regions.
97.9% (138/141) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[9] 544 Arg (31.0, 42.1)
[9] 569 Ile (57.8, 135.8)
[9] 608 Ile (30.4, -98.4)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 10









84.4% (119/141) of all residues were in favored (98%) regions.
97.9% (138/141) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[10] 517 Glu (47.3, 101.7)
[10] 569 Ile (44.8, 132.5)
[10] 650 Ser (47.9, 85.1)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 11









84.4% (119/141) of all residues were in favored (98%) regions.
97.2% (137/141) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [11] 513 Val (57.8, 179.0)
- [11] 532 Asn (-155.9, -95.9)
- [11] 569 Ile (54.4, 138.6)
- [11] 608 Ile (29.8, -94.6)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 12









80.9% (114/141) of all residues were in favored (98%) regions.
96.5% (136/141) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [12] 513 Val (61.5, 161.9)
- [12] 520 Asn (96.6, 136.8)
- [12] 532 Asn (-164.3, -86.5)
- [12] 569 Ile (54.9, 145.3)
- [12] 608 Ile (43.3, -94.0)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 13









83.7% (118/141) of all residues were in favored (98%) regions.
96.5% (136/141) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[13] -1 Met (46.1, 80.8)
[13] 522 Lys (54.6, 118.8)
[13] 569 Ile (49.8, 141.9)
[13] 608 Ile (26.9, -94.8)
[13] 611 Val (-147.7, -62.9)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 14









87.2% (123/141) of all residues were in favored (98%) regions.
98.6% (139/141) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):

[14] 516 Ser (55.4, 167.2)
[14] 569 Ile (-37.2, 145.5)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 15








80.9% (114/141) of all residues were in favored (98%) regions.
97.9% (138/141) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[15] 523 Glu (44.4, -158.1)
[15] 608 Ile (-42.1, -76.6)
[15] 650 Ser (63.8, 102.5)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 16




85.1% (120/141) of all residues were in favored (98%) regions.
97.2% (137/141) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[16] 517 Glu (36.3, 107.8)
[16] 569 Ile (46.7, 114.1)
[16] 572 Gly (71.1, 107.5)
[16] 573 Ile (41.2, 101.0)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 17









83.0% (117/141) of all residues were in favored (98%) regions.
94.3% (133/141) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

[17] -2 His (56.0, -179.9)
[17] -1 Met (164.9, -31.6)
[17] 523 Glu (41.3, -161.6)
[17] 532 Asn (-170.5, -105.4)
[17] 569 Ile (52.2, 146.4)
[17] 578 Val (-79.4, 31.2)
[17] 579 Arg (24.2, 71.4)
[17] 608 Ile (23.8, -90.6)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 18









85.8% (121/141) of all residues were in favored (98%) regions.
98.6% (139/141) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[18] 565 Lys (35.6, -157.7)
[18] 566 Gly (-41.8, 13.0)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 19









83.0% (117/141) of all residues were in favored (98%) regions.
97.2% (137/141) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[19] -3 Ser (62.8, -90.5)
[19] 513 Val (58.2, 97.7)
[19] 569 Ile (-37.6, 145.8)
[19] 608 Ile (39.9, -101.4)

MolProbity Ramachandran analysis

2rnk.H.pdb, model 20



82.3% (116/141) of all residues were in favored (98%) regions.
96.5% (136/141) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[20] -1 Met (-171.1, -51.7)
[20] 513 Val (65.0, 179.8)
[20] 518 Ala (64.8, 111.3)
[20] 523 Glu (38.7, -158.4)
[20] 569 Ile (56.5, 153.8)