


MolProbity Ramachandran analysis

7lhq.H.pdb, all models




95.7% (1569/1640) of all residues were in favored (98%) regions.
98.3% (1612/1640) of all residues were in allowed (>99.8%) regions.

There were 28 outliers (phi, psi):

[1] 10 Thr (52.3, 96.7)	[7] 6 Asp (53.4, -178.0)	[12] 7 Val (51.0, 92.7)
[1] 68 Asp (168.9, -29.3)	[7] 68 Asp (164.1, -25.2)	[12] 68 Asp (168.6, -30.0)
[2] 68 Asp (169.0, -29.3)	[8] 68 Asp (169.0, -30.3)	[13] 3 Lys (54.3, -177.5)
[2] 82 Thr (52.1, 94.1)	[9] 68 Asp (169.1, -30.5)	[13] 68 Asp (168.7, -32.3)
[3] 68 Asp (168.4, -25.4)	[10] 68 Asp (169.9, -30.9)	[14] 68 Asp (169.0, -30.8)
[4] 68 Asp (167.3, -30.4)	[11] 2 Ser (65.2, -76.3)	[15] 10 Thr (55.7, 99.4)
[5] 68 Asp (168.9, -28.9)	[11] 64 Gln (-161.7, -30.9)	[16] 64 Gln (-161.8, -30.7)
[6] 68 Asp (168.6, -25.1)	[11] 68 Asp (168.8, -31.0)	[16] 68 Asp (172.7, -30.7)
		[17] 68 Asp (168.9, -31.8)
		[18] 68 Asp (165.2, -26.1)
		[19] 68 Asp (165.0, -24.9)
		[20] 68 Asp (168.7, -29.7)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 1



95.1% (78/82) of all residues were in favored (98%) regions.

97.6% (80/82) of all residues were in allowed (>99.8%) regions.


There were 2 outliers (phi, psi):

[1] 10 Thr (52.3, 96.7)

[1] 68 Asp (168.9, -29.3)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 2



95.1% (78/82) of all residues were in favored (98%) regions.

97.6% (80/82) of all residues were in allowed (>99.8%) regions.


There were 2 outliers (phi, psi):

[2] 68 Asp (169.0, -29.3)

[2] 82 Thr (52.1, 94.1)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 3









95.1% (78/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[3] 68 Asp (168.4, -25.4)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 4




96.3% (79/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[4] 68 Asp (167.3, -30.4)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 5




96.3% (79/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[5] 68 Asp (168.9, -28.9)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 6









96.3% (79/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[6] 68 Asp (168.6, -25.1)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 7




95.1% (78/82) of all residues were in favored (98%) regions.
97.6% (80/82) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[7] 6 Asp (53.4, -178.0)
[7] 68 Asp (164.1, -25.2)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 8









96.3% (79/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[8] 68 Asp (169.0, -30.3)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 9




96.3% (79/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[9] 68 Asp (169.1, -30.5)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 10




96.3% (79/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[10] 68 Asp (169.9, -30.9)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 11




95.1% (78/82) of all residues were in favored (98%) regions.
96.3% (79/82) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[11] 2 Ser (65.2, -76.3)
[11] 64 Gln (-161.7, -30.9)
[11] 68 Asp (168.8, -31.0)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 12



93.9% (77/82) of all residues were in favored (98%) regions.

97.6% (80/82) of all residues were in allowed (>99.8%) regions.


There were 2 outliers (phi, psi):

[12] 7 Val (51.0, 92.7)

[12] 68 Asp (168.6, -30.0)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 13



95.1% (78/82) of all residues were in favored (98%) regions.

97.6% (80/82) of all residues were in allowed (>99.8%) regions.


There were 2 outliers (phi, psi):

[13] 3 Lys (54.3, -177.5)

[13] 68 Asp (168.7, -32.3)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 14









96.3% (79/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[14] 68 Asp (169.0, -30.8)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 15




93.9% (77/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[15] 10 Thr (55.7, 99.4)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 16



96.3% (79/82) of all residues were in favored (98%) regions.

97.6% (80/82) of all residues were in allowed (>99.8%) regions.


There were 2 outliers (phi, psi):

[16] 64 Gln (-161.8, -30.7)

[16] 68 Asp (172.7, -30.7)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 17









95.1% (78/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[17] 68 Asp (168.9, -31.8)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 18









96.3% (79/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[18] 68 Asp (165.2, -26.1)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 19




96.3% (79/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[19] 68 Asp (165.0, -24.9)

MolProbity Ramachandran analysis

7lhq.H.pdb, model 20



96.3% (79/82) of all residues were in favored (98%) regions.
98.8% (81/82) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[20] 68 Asp (168.7, -29.7)