


MolProbity Ramachandran analysis

1ysy.H.pdb, all models



66.7% (1107/1660) of all residues were in favored (98%) regions.

92.5% (1536/1660) of all residues were in allowed (>99.8%) regions.

This list is truncated; use the MolProbity multi-chart.html for complete list.







There were 124 outliers (phi, psi):

[1] 7 Asp (44.9, 104.2)	[2] 30 Leu (-44.8, -74.4)	[3] 50 Ala (-58.5, -76.4)
[1] 11 Thr (-47.3, -74.1)	[2] 66 Gly (161.3, 31.0)	[3] 70 Ile (46.9, 17.8)
[1] 27 Ser (35.7, 86.6)	[2] 67 Ala (-80.3, -139.6)	[4] 3 Ser (-54.2, 101.7)
[1] 50 Ala (-59.5, -76.2)	[2] 68 Val (38.8, -62.7)	[4] 27 Ser (47.1, -161.4)
[1] 66 Gly (150.2, 14.5)	[3] 2 His (-69.7, -98.6)	[4] 50 Ala (-59.7, -77.3)
[1] 73 Leu (-54.6, 0.7)	[3] 3 Ser (-26.0, 100.3)	[4] 70 Ile (-150.3, 12.7)
[2] 4 Lys (-168.2, -44.4)	[3] 10 Cys (102.9, -11.3)	[5] 7 Asp (95.7, 133.7)
[2] 11 Thr (-60.4, -82.6)	[3] 28 Ser (95.5, 122.4)	[5] 21 Gln (-23.4, -54.8)
		[5] 50 Ala (-58.2, -78.2)
		[5] 66 Gly (143.6, 50.9)
		[5] 70 Ile (-150.7, 36.1)
		[5] 82 Ala (-162.8, 5.1)

[6] 11 Thr (-30.5, -78.3)	[7] 50 Ala (-50.5, -77.5)	[9] 11 Thr (45.1, -91.2)
[6] 50 Ala (-58.2, -85.0)	[8] 14 Val (-23.8, -63.3)	[9] 48 Thr (-68.4, -80.3)
[6] 66 Gly (138.6, 42.3)	[8] 28 Ser (-168.5, 17.9)	[9] 50 Ala (-56.6, -84.5)
[6] 84 Leu (-162.4, -36.6)	[8] 50 Ala (-55.1, -80.1)	[9] 66 Gly (149.2, 26.7)
[7] 3 Ser (36.7, 23.6)	[8] 66 Gly (145.1, 49.5)	[9] 82 Ala (-179.5, 17.4)
[7] 6 Ser (-54.4, 100.4)	[8] 70 Ile (-155.7, 17.4)	[10] 3 Ser (42.1, 80.9)
[7] 7 Asp (44.4, 102.7)	[8] 75 Glu (-26.5, -40.9)	[10] 6 Ser (-68.1, -80.7)
[7] 24 Val (-98.8, -103.3)	[9] 2 His (57.2, 166.9)	[10] 14 Val (-69.2, -75.7)
[7] 26 Ser (143.1, -115.7)	[9] 3 Ser (44.1, 100.6)	[10] 26 Ser (-58.9, 4.3)
[7] 29 Lys (-80.4, -100.1)	[9] 5 Met (-22.9, 83.8)	[10] 45 Lys (-167.7, -65.7)
[7] 41 Ile (-48.4, -12.2)	[9] 8 Val (-148.9, 5.3)	[10] 50 Ala (-58.0, -79.3)
[7] 45 Lys (63.7, -81.9)	[9] 10 Cys (-81.1, -96.7)	[11] 2 His (178.9, -63.9)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 1









63.9% (53/83) of all residues were in favored (98%) regions.
92.8% (77/83) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [1] 7 Asp (44.9, 104.2)
- [1] 11 Thr (-47.3, -74.1)
- [1] 27 Ser (35.7, 86.6)
- [1] 50 Ala (-59.5, -76.2)
- [1] 66 Gly (150.2, 14.5)
- [1] 73 Leu (-54.6, 0.7)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 2









62.7% (52/83) of all residues were in favored (98%) regions.
92.8% (77/83) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [2] 4 Lys (-168.2, -44.4)
- [2] 11 Thr (-60.4, -82.6)
- [2] 30 Leu (-44.8, -74.4)
- [2] 66 Gly (161.3, 31.0)
- [2] 67 Ala (-80.3, -139.6)
- [2] 68 Val (38.8, -62.7)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 3









62.7% (52/83) of all residues were in favored (98%) regions.
92.8% (77/83) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [3] 2 His (-69.7, -98.6)
- [3] 3 Ser (-26.0, 100.3)
- [3] 10 Cys (102.9, -11.3)
- [3] 28 Ser (95.5, 122.4)
- [3] 50 Ala (-58.5, -76.4)
- [3] 70 Ile (46.9, 17.8)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 4









72.3% (60/83) of all residues were in favored (98%) regions.
95.2% (79/83) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [4] 3 Ser (-54.2, 101.7)
- [4] 27 Ser (47.1, -161.4)
- [4] 50 Ala (-59.7, -77.3)
- [4] 70 Ile (-150.3, 12.7)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 5




73.5% (61/83) of all residues were in favored (98%) regions.
92.8% (77/83) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [5] 7 Asp (95.7, 133.7)
- [5] 21 Gln (-23.4, -54.8)
- [5] 50 Ala (-58.2, -78.2)
- [5] 66 Gly (143.6, 50.9)
- [5] 70 Ile (-150.7, 36.1)
- [5] 82 Ala (-162.8, 5.1)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 6









68.7% (57/83) of all residues were in favored (98%) regions.
95.2% (79/83) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [6] 11 Thr (-30.5, -78.3)
- [6] 50 Ala (-58.2, -85.0)
- [6] 66 Gly (138.6, 42.3)
- [6] 84 Leu (-162.4, -36.6)

MolProbity Ramachandran analysis

1lisy.H.pdb, model 7








68.7% (57/83) of all residues were in favored (98%) regions.
89.2% (74/83) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[7] 3 Ser (36.7, 23.6)	[7] 50 Ala (-50.5, -77.5)
[7] 6 Ser (-54.4, 100.4)	
[7] 7 Asp (44.4, 102.7)	
[7] 24 Val (-98.8, -103.3)	
[7] 26 Ser (143.1, -115.7)	
[7] 29 Lys (-80.4, -100.1)	
[7] 41 Ile (-48.4, -12.2)	
[7] 45 Lys (63.7, -81.9)	

MolProbity Ramachandran analysis

1tsy.H.pdb, model 8









65.1% (54/83) of all residues were in favored (98%) regions.
92.8% (77/83) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [8] 14 Val (-23.8, -63.3)
- [8] 28 Ser (-168.5, 17.9)
- [8] 50 Ala (-55.1, -80.1)
- [8] 66 Gly (145.1, 49.5)
- [8] 70 Ile (-155.7, 17.4)
- [8] 75 Glu (-26.5, -40.9)

MolProbity Ramachandran analysis

1tsy.H.pdb, model 9









63.9% (53/83) of all residues were in favored (98%) regions.
88.0% (73/83) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[9] 2 His (57.2, 166.9)	[9] 66 Gly (149.2, 26.7)
[9] 3 Ser (44.1, 100.6)	[9] 82 Ala (-179.5, 17.4)
[9] 5 Met (-22.9, 83.8)	
[9] 8 Val (-148.9, 5.3)	
[9] 10 Cys (-81.1, -96.7)	
[9] 11 Thr (45.1, -91.2)	
[9] 48 Thr (-68.4, -80.3)	
[9] 50 Ala (-56.6, -84.5)	

MolProbity Ramachandran analysis

1ysy.H.pdb, model 10




68.7% (57/83) of all residues were in favored (98%) regions.
92.8% (77/83) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [10] 3 Ser (42.1, 80.9)
- [10] 6 Ser (-68.1, -80.7)
- [10] 14 Val (-69.2, -75.7)
- [10] 26 Ser (-58.9, 4.3)
- [10] 45 Lys (-167.7, -65.7)
- [10] 50 Ala (-58.0, -79.3)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 11









61.4% (51/83) of all residues were in favored (98%) regions.
91.6% (76/83) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [11] 2 His (178.9, -63.9)
- [11] 3 Ser (59.1, 115.4)
- [11] 10 Cys (-69.4, -107.1)
- [11] 26 Ser (153.8, 163.5)
- [11] 50 Ala (-56.5, -75.3)
- [11] 77 Met (-58.1, 2.6)
- [11] 82 Ala (-164.9, 15.4)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 12









66.3% (55/83) of all residues were in favored (98%) regions.
95.2% (79/83) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [12] 3 Ser (-165.1, 17.4)
- [12] 8 Val (-150.3, 9.8)
- [12] 66 Gly (149.8, 68.2)
- [12] 68 Val (57.7, -30.7)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 13









65.1% (54/83) of all residues were in favored (98%) regions.
97.6% (81/83) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[13] 6 Ser (-66.2, 65.0)
[13] 50 Ala (-56.3, -86.2)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 14









65.1% (54/83) of all residues were in favored (98%) regions.
95.2% (79/83) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [14] 23 Arg (-42.2, -76.5)
- [14] 33 Gln (-54.7, 0.3)
- [14] 50 Ala (-50.3, -86.6)
- [14] 66 Gly (137.3, 68.2)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 15








61.4% (51/83) of all residues were in favored (98%) regions.
89.2% (74/83) of all residues were in allowed (>99.8%) regions.

There were 9 outliers (phi, psi):

[15] 11 Thr (-75.2, -92.8)	[15] 71 Asn (-66.3, -80.1)
[15] 26 Ser (-51.9, 75.8)	
[15] 33 Gln (-56.1, 0.1)	
[15] 48 Thr (-61.0, -81.1)	
[15] 50 Ala (-58.6, -83.0)	
[15] 56 Ser (-53.6, -76.3)	
[15] 68 Val (-80.6, -101.1)	
[15] 70 Ile (-149.5, 1.7)	

MolProbity Ramachandran analysis

1ysy.H.pdb, model 16









67.5% (56/83) of all residues were in favored (98%) regions.
92.8% (77/83) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [16] 8 Val (-148.9, -1.6)
- [16] 9 Lys (69.9, 138.3)
- [16] 11 Thr (-66.2, -85.4)
- [16] 24 Val (-123.6, -104.5)
- [16] 48 Thr (-71.5, -84.5)
- [16] 50 Ala (-61.1, -82.1)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 17



67.5% (56/83) of all residues were in favored (98%) regions.

94.0% (78/83) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[17] 4 Lys (-162.0, -29.6)

[17] 7 Asp (47.2, 115.1)







[17] 11 Thr (-24.9, -74.3)

[17] 25 Glu (95.6, 111.2)

[17] 50 Ala (-64.9, -81.6)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 18









68.7% (57/83) of all residues were in favored (98%) regions.
91.6% (76/83) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[18] 7 Asp (46.2, 104.4)
[18] 10 Cys (49.1, -17.7)
[18] 24 Val (-85.0, -160.3)
[18] 28 Ser (64.7, 135.5)
[18] 48 Thr (-60.9, -76.8)
[18] 50 Ala (-56.4, -83.5)
[18] 70 Ile (-79.5, 23.9)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 19









75.9% (63/83) of all residues were in favored (98%) regions.
92.8% (77/83) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

[19] 2 His (58.2, -88.0)
[19] 7 Asp (65.1, 151.5)
[19] 8 Val (-154.4, -4.6)
[19] 23 Arg (-68.5, -80.5)
[19] 28 Ser (24.4, 73.4)
[19] 66 Gly (138.6, 41.2)

MolProbity Ramachandran analysis

1ysy.H.pdb, model 20



65.1% (54/83) of all residues were in favored (98%) regions.
86.7% (72/83) of all residues were in allowed (>99.8%) regions.

There were 11 outliers (phi, psi):

[20] 2 His (64.6, 98.7)	[20] 50 Ala (-58.4, -81.3)
[20] 3 Ser (66.2, 149.3)	[20] 66 Gly (136.3, 54.3)
[20] 6 Ser (47.6, -161.2)	[20] 70 Ile (-160.2, 18.0)
[20] 14 Val (-47.1, -70.8)	
[20] 22 Leu (-163.3, -28.9)	
[20] 26 Ser (91.8, 56.2)	
[20] 28 Ser (44.4, -167.0)	
[20] 48 Thr (-67.7, -86.5)	