


MolProbity Ramachandran analysis

2kys.H.pdb, all models



79.7% (1323/1660) of all residues were in favored (98%) regions.

95.9% (1592/1660) of all residues were in allowed (>99.8%) regions.

This list is truncated; use the MolProbity multi-chart.html for complete list.

There were 68 outliers (phi, psi):

[1] 3 Ser (43.8, -86.4)	[3] 10 Cys (52.2, 88.6)	[5] 23 Arg (175.3, 27.7)
[1] 5 Met (-57.1, 96.9)	[3] 12 Ser (44.1, 99.1)	[6] 10 Cys (50.9, 177.5)
[1] 23 Arg (-169.4, 21.1)	[3] 23 Arg (-168.0, 15.7)	[6] 12 Ser (43.8, 84.7)
[1] 69 Asp (-65.3, 16.5)	[4] 23 Arg (-175.1, -41.0)	[6] 23 Arg (-171.9, 26.0)
[2] 4 Lys (46.1, 165.6)	[4] 24 Val (42.0, 129.1)	[7] 4 Lys (178.7, -34.7)
[2] 23 Arg (-179.0, 25.4)	[5] 3 Ser (45.9, -77.5)	[7] 5 Met (176.1, 122.0)
[3] 2 His (-161.4, -39.1)	[5] 7 Asp (48.5, 83.5)	[7] 12 Ser (44.5, 107.6)
[3] 5 Met (46.3, -62.6)	[5] 10 Cys (55.7, -66.4)	[7] 23 Arg (-174.9, 26.6)
		[7] 27 Ser (-70.9, 23.5)
		[8] 12 Ser (49.4, 97.8)
		[15] 23 Arg (-178.5, 39.1)
		[9] 11 Thr (-173.0, -67.4)

[9] 23 Arg (178.6, 35.8)

[9] 69 Asp (-67.0, 20.7)

[10] 4 Lys (57.4, 116.0)

[10] 12 Ser (54.7, 110.8)

[10] 23 Arg (174.9, 28.8)

[10] 27 Ser (-70.2, 31.9)

[11] 23 Arg (178.2, 24.8)

[11] 25 Glu (-60.7, 89.2)

[12] 5 Met (46.6, -163.8)

[12] 10 Cys (59.4, 148.7)

[12] 23 Arg (-168.1, -22.2)

[12] 24 Val (43.9, -174.3)

[12] 69 Asp (-70.1, 19.0)

[13] 10 Cys (58.5, 150.0)

[13] 12 Ser (49.6, 174.6)

[13] 44 Ala (44.3, -153.8)

[14] 11 Thr (43.7, -86.5)

[14] 23 Arg (-172.6, 18.8)

[14] 26 Ser (-69.5, 30.6)

[14] 76 Glu (-63.7, -85.2)

[15] 5 Met (44.7, 21.2)

[15] 6 Ser (43.3, 24.2)

[15] 11 Thr (-163.1, -53.7)

[15] 23 Arg (-169.9, 6.1)

[16] 3 Ser (51.9, -176.5)

[16] 4 Lys (-162.2, -28.3)

[16] 12 Ser (52.2, -172.4)

[16] 24 Val (-44.1, 161.8)

[17] 3 Ser (63.2, -85.1)

[17] 10 Cys (67.3, 102.1)

[17] 23 Arg (179.9, 32.2)

[17] 24 Val (-44.3, 155.0)

[18] 8 Val (-76.1, 25.1)







[18] 18 Val (-44.8, -17.5)

[18] 23 Arg (-173.5, 25.6)

[19] 6 Ser (52.6, 172.3)

MolProbity Ramachandran analysis

2kys.H.pdb, model 1









80.7% (67/83) of all residues were in favored (98%) regions.
95.2% (79/83) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [1] 3 Ser (43.8, -86.4)
- [1] 5 Met (-57.1, 96.9)
- [1] 23 Arg (-169.4, 21.1)
- [1] 69 Asp (-65.3, 16.5)

MolProbity Ramachandran analysis

2kys.H.pdb, model 2



83.1% (69/83) of all residues were in favored (98%) regions.
97.6% (81/83) of all residues were in allowed (>99.8%) regions.







There were 2 outliers (phi, psi):

[2] 4 Lys (46.1, 165.6)

[2] 23 Arg (-179.0, 25.4)

MolProbity Ramachandran analysis

2kys.H.pdb, model 3









80.7% (67/83) of all residues were in favored (98%) regions.
94.0% (78/83) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[3] 2 His (-161.4, -39.1)
[3] 5 Met (46.3, -62.6)
[3] 10 Cys (52.2, 88.6)
[3] 12 Ser (44.1, 99.1)
[3] 23 Arg (-168.0, 15.7)

MolProbity Ramachandran analysis

2kys.H.pdb, model 4









84.3% (70/83) of all residues were in favored (98%) regions.
97.6% (81/83) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[4] 23 Arg (-175.1, -41.0)
[4] 24 Val (42.0, 129.1)

MolProbity Ramachandran analysis

2kys.H.pdb, model 5




69.9% (58/83) of all residues were in favored (98%) regions.
95.2% (79/83) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [5] 3 Ser (45.9, -77.5)
- [5] 7 Asp (48.5, 83.5)
- [5] 10 Cys (55.7, -66.4)
- [5] 23 Arg (175.3, 27.7)

MolProbity Ramachandran analysis

2kys.H.pdb, model 6




74.7% (62/83) of all residues were in favored (98%) regions.
96.4% (80/83) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[6] 10 Cys (50.9, 177.5)
[6] 12 Ser (43.8, 84.7)
[6] 23 Arg (-171.9, 26.0)

MolProbity Ramachandran analysis

2kys.H.pdb, model 7



74.7% (62/83) of all residues were in favored (98%) regions.







94.0% (78/83) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [7] 4 Lys (178.7, -34.7)
- [7] 5 Met (176.1, 122.0)
- [7] 12 Ser (44.5, 107.6)
- [7] 23 Arg (-174.9, 26.6)
- [7] 27 Ser (-70.9, 23.5)

MolProbity Ramachandran analysis

2kys.H.pdb, model 8




81.9% (68/83) of all residues were in favored (98%) regions.
97.6% (81/83) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[8] 12 Ser (49.4, 97.8)
[8] 23 Arg (-178.5, 39.1)

MolProbity Ramachandran analysis

2kys.H.pdb, model 9




83.1% (69/83) of all residues were in favored (98%) regions.
96.4% (80/83) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[9] 11 Thr (-173.0, -67.4)
[9] 23 Arg (178.6, 35.8)
[9] 69 Asp (-67.0, 20.7)

MolProbity Ramachandran analysis

2kys.H.pdb, model 10




78.3% (65/83) of all residues were in favored (98%) regions.
95.2% (79/83) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [10] 4 Lys (57.4, 116.0)
- [10] 12 Ser (54.7, 110.8)
- [10] 23 Arg (174.9, 28.8)
- [10] 27 Ser (-70.2, 31.9)

MolProbity Ramachandran analysis

2kys.H.pdb, model 11









81.9% (68/83) of all residues were in favored (98%) regions.
97.6% (81/83) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):

[11] 23 Arg (178.2, 24.8)
[11] 25 Glu (-60.7, 89.2)

MolProbity Ramachandran analysis

2kys.H.pdb, model 12









77.1% (64/83) of all residues were in favored (98%) regions.
94.0% (78/83) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[12] 5 Met (46.6, -163.8)
[12] 10 Cys (59.4, 148.7)
[12] 23 Arg (-168.1, -22.2)
[12] 24 Val (43.9, -174.3)
[12] 69 Asp (-70.1, 19.0)

MolProbity Ramachandran analysis

2kys.H.pdb, model 13









78.3% (65/83) of all residues were in favored (98%) regions.
96.4% (80/83) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[13] 10 Cys (58.5, 150.0)
[13] 12 Ser (49.6, 174.6)
[13] 44 Ala (44.3, -153.8)

MolProbity Ramachandran analysis

2kys.H.pdb, model 14









81.9% (68/83) of all residues were in favored (98%) regions.
95.2% (79/83) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [14] 11 Thr (43.7, -86.5)
- [14] 23 Arg (-172.6, 18.8)
- [14] 26 Ser (-69.5, 30.6)
- [14] 76 Glu (-63.7, -85.2)

MolProbity Ramachandran analysis

2kys.H.pdb, model 15









80.7% (67/83) of all residues were in favored (98%) regions.
95.2% (79/83) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[15] 5 Met (44.7, 21.2)
[15] 6 Ser (43.3, 24.2)
[15] 11 Thr (-163.1, -53.7)
[15] 23 Arg (-169.9, 6.1)

MolProbity Ramachandran analysis

2kys.H.pdb, model 16









81.9% (68/83) of all residues were in favored (98%) regions.
95.2% (79/83) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[16] 3 Ser (51.9, -176.5)
[16] 4 Lys (-162.2, -28.3)
[16] 12 Ser (52.2, -172.4)
[16] 24 Val (-44.1, 161.8)

MolProbity Ramachandran analysis

2kys.H.pdb, model 17









78.3% (65/83) of all residues were in favored (98%) regions.
95.2% (79/83) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [17] 3 Ser (63.2, -85.1)
- [17] 10 Cys (67.3, 102.1)
- [17] 23 Arg (179.9, 32.2)
- [17] 24 Val (-44.3, 155.0)

MolProbity Ramachandran analysis

2kys.H.pdb, model 18









79.5% (66/83) of all residues were in favored (98%) regions.
96.4% (80/83) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[18] 8 Val (-76.1, 25.1)
[18] 18 Val (-44.8, -17.5)
[18] 23 Arg (-173.5, 25.6)

MolProbity Ramachandran analysis

2kys.H.pdb, model 19









83.1% (69/83) of all residues were in favored (98%) regions.
96.4% (80/83) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[19] 6 Ser (52.6, 172.3)
[19] 23 Arg (-172.5, 41.0)
[19] 69 Asp (-65.9, 19.9)

MolProbity Ramachandran analysis

2kys.H.pdb, model 20



79.5% (66/83) of all residues were in favored (98%) regions.
97.6% (81/83) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[20] 23 Arg (-179.8, 36.4)
[20] 69 Asp (-63.6, 17.0)