


6yi3.H.pdb, all models



There were 52 outliers (phi, psi):


[1] 9 Thr (35.1, 77.1)	[8] 137 Arg (60.3, 93.0)	[15] 24 Leu (64.2, 141.5)
[1] 24 Leu (173.8, 136.0)	[9] 8 Asn (-158.6, -98.3)	[17] 64 Leu (52.9, 178.7)
[5] 4 Gly (-175.2, 78.1)	[9] 24 Leu (165.7, 137.4)	[18] 9 Thr (-37.2, 94.9)
[5] 24 Leu (165.3, 141.9)	[10] 9 Thr (-17.7, 97.4)	[18] 24 Leu (157.2, 144.8)
[6] 2 Ala (61.9, 154.2)	[11] 4 Gly (-174.2, -79.0)	[18] 63 Asp (47.0, -161.5)
[6] 24 Leu (167.5, 137.4)	[11] 24 Leu (172.6, 144.8)	[18] 64 Leu (53.5, -176.7)
[8] 4 Gly (-166.3, -34.7)	[13] 9 Thr (28.3, 81.1)	[19] 63 Asp (26.6, 86.3)
[8] 24 Leu (173.7, 144.5)	[13] 24 Leu (161.0, 141.8)	[20] 24 Leu (173.7, 145.6)

[21]	8 Asn (-168.2, -73.7)	[31]	9 Thr (35.7, 78.9)
[21]	24 Leu (174.5, 134.2)	[32]	24 Leu (176.0, 135.2)
[22]	24 Leu (168.6, 136.7)	[33]	8 Asn (-156.1, -102.5)
[23]	24 Leu (165.3, 142.2)	[35]	8 Asn (-158.4, -91.8)
[24]	8 Asn (-164.7, -80.6)	[35]	24 Leu (172.7, 141.6)
[25]	24 Leu (173.0, 142.7)	[36]	8 Asn (-163.1, -86.8)
[26]	24 Leu (62.5, 146.3)	[36]	9 Thr (39.4, 79.0)
[27]	2 Ala (62.0, -23.2)	[36]	24 Leu (173.9, 141.0)
[28]	9 Thr (24.6, 78.8)	[37]	24 Leu (171.3, 142.2)
[28]	24 Leu (66.5, 144.2)	[39]	7 Asn (-168.7, -55.3)
[30]	24 Leu (153.7, 143.4)	[39]	24 Leu (172.3, 141.7)
[30]	63 Asp (30.7, -121.6)	[40]	63 Asp (38.5, -140.8)

[31] 9 Thr (35.7, 78.9)
[32] 24 Leu (176.0, 135.2)
[33] 8 Asn (-156.1, -102.5)
[35] 8 Asn (-158.4, -91.8)
[35] 24 Leu (172.7, 141.6)
[36] 8 Asn (-163.1, -86.8)
[36] 9 Thr (39.4, 79.0)
[36] 24 Leu (173.9, 141.0)
[37] 24 Leu (171.3, 142.2)
[39] 7 Asn (-168.7, -55.3)
[39] 24 Leu (172.3, 141.7)
[40] 63 Asp (38.5, -140.8)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 1









95.7% (132/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):

[1] 9 Thr (35.1, 77.1)
[1] 24 Leu (173.8, 136.0)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 2









92.0% (127/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

6yi3.H.pdb, model 3









97.1% (134/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

6yi3.H.pdb, model 4









94.2% (130/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

6yi3.H.pdb, model 5




93.5% (129/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[5] 4 Gly (-175.2, 78.1)
[5] 24 Leu (165.3, 141.9)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 6









94.9% (131/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[6] 2 Ala (61.9, 154.2)
[6] 24 Leu (167.5, 137.4)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 7









94.2% (130/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

6yi3.H.pdb, model 8




94.9% (131/138) of all residues were in favored (98%) regions.
97.8% (135/138) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[8] 4 Gly (-166.3, -34.7)
[8] 24 Leu (173.7, 144.5)
[8] 137 Arg (60.3, 93.0)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 9









92.8% (128/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[9] 8 Asn (-158.6, -98.3)
[9] 24 Leu (165.7, 137.4)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 10




94.2% (130/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[10] 9 Thr (-17.7, 97.4)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 11









93.5% (129/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[11] 4 Gly (-174.2, -79.0)
[11] 24 Leu (172.6, 144.8)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 12









97.1% (134/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

6yi3.H.pdb, model 13









94.9% (131/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[13] 9 Thr (28.3, 81.1)
[13] 24 Leu (161.0, 141.8)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 14









92.0% (127/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[14] 9 Thr (10.2, 83.8)
[14] 24 Leu (171.6, 144.2)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 15








92.0% (127/138) of all residues were in favored (98%) regions.
97.8% (135/138) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[15] 9 Thr (40.5, 79.6)
[15] 20 Gly (-42.1, 103.0)
[15] 24 Leu (64.2, 141.5)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 16









96.4% (133/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

6yi3.H.pdb, model 17









96.4% (133/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[17] 64 Leu (52.9, 178.7)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 18









93.5% (129/138) of all residues were in favored (98%) regions.
97.1% (134/138) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[18] 9 Thr (-37.2, 94.9)
[18] 24 Leu (157.2, 144.8)
[18] 63 Asp (47.0, -161.5)
[18] 64 Leu (53.5, -176.7)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 19




94.9% (131/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[19] 63 Asp (26.6, 86.3)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 20




94.2% (130/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[20] 24 Leu (173.7, 145.6)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 21









92.8% (128/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[21] 8 Asn (-168.2, -73.7)
[21] 24 Leu (174.5, 134.2)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 22









94.2% (130/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[22] 24 Leu (168.6, 136.7)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 23








94.2% (130/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[23] 24 Leu (165.3, 142.2)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 24









93.5% (129/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[24] 8 Asn (-164.7, -80.6)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 25









94.2% (130/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[25] 24 Leu (173.0, 142.7)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 26









94.2% (130/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[26] 24 Leu (62.5, 146.3)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 27









92.8% (128/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[27] 2 Ala (62.0, -23.2)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 28









92.8% (128/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[28] 9 Thr (24.6, 78.8)
[28] 24 Leu (66.5, 144.2)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 29









93.5% (129/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

6yi3.H.pdb, model 30









92.8% (128/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[30] 24 Leu (153.7, 143.4)
[30] 63 Asp (30.7, -121.6)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 31




94.9% (131/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[31] 9 Thr (35.7, 78.9)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 32









96.4% (133/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[32] 24 Leu (176.0, 135.2)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 33









95.7% (132/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[33] 8 Asn (-156.1, -102.5)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 34









95.7% (132/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

6yi3.H.pdb, model 35




92.0% (127/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[35] 8 Asn (-158.4, -91.8)
[35] 24 Leu (172.7, 141.6)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 36









91.3% (126/138) of all residues were in favored (98%) regions.
97.8% (135/138) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[36] 8 Asn (-163.1, -86.8)
[36] 9 Thr (39.4, 79.0)
[36] 24 Leu (173.9, 141.0)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 37









92.0% (127/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[37] 24 Leu (171.3, 142.2)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 38




94.2% (130/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

6yi3.H.pdb, model 39








90.6% (125/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[39] 7 Asn (-168.7, -55.3)
[39] 24 Leu (172.3, 141.7)

MolProbity Ramachandran analysis

6yi3.H.pdb, model 40



94.2% (130/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[40] 63 Asp (38.5, -140.8)