










7acs.H.pdb, all models



[24] 63 Asp (65.8, 99.1)
[25] 66 Pro (-52.9, 102.6)
[25] 123 Gln (-57.7, 98.3)
[26] 14 Thr (-78.8, -142.1)
[26] 63 Asp (50.1, 82.9)
[27] 24 Leu (169.8, 137.5)
[27] 63 Asp (63.4, 105.2)
[28] 63 Asp (64.4, 99.3)
[29] 63 Asp (48.7, 82.8)
[30] 37 Asn (-173.2, -28.2)

MolProbity Ramachandran analysis

7acs.H.pdb, model 1









85.5% (118/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):

[1] 9 Thr (48.8, 83.8)
[1] 63 Asp (61.4, 107.7)

MolProbity Ramachandran analysis

7acs.H.pdb, model 2









92.8% (128/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[2] 37 Asn (-161.7, -38.6)
[2] 139 Gly (168.8, -90.5)

MolProbity Ramachandran analysis

7acs.H.pdb, model 3









87.0% (120/138) of all residues were in favored (98%) regions.
97.8% (135/138) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[3] 9 Thr (65.7, 117.7)
[3] 24 Leu (172.5, 131.2)
[3] 63 Asp (56.0, 95.6)

MolProbity Ramachandran analysis

7acs.H.pdb, model 4









91.3% (126/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[4] 63 Asp (62.8, 101.0)

MolProbity Ramachandran analysis

7acs.H.pdb, model 5









88.4% (122/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

7acs.H.pdb, model 6









89.9% (124/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[6] 63 Asp (55.4, 97.1)

MolProbity Ramachandran analysis

7acs.H.pdb, model 7








89.1% (123/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[7] 63 Asp (63.9, 115.2)

MolProbity Ramachandran analysis

7acs.H.pdb, model 8









91.3% (126/138) of all residues were in favored (98%) regions.
97.8% (135/138) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[8] 24 Leu (178.3, 125.1)
[8] 63 Asp (52.4, 88.9)
[8] 139 Gly (164.3, 36.9)

MolProbity Ramachandran analysis

7acs.H.pdb, model 9




91.3% (126/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[9] 63 Asp (63.9, 115.9)

MolProbity Ramachandran analysis

7acs.H.pdb, model 10









93.5% (129/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[10] 24 Leu (173.1, 122.3)
[10] 63 Asp (56.8, 94.4)

MolProbity Ramachandran analysis

7acs.H.pdb, model 11









87.0% (120/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[11] 63 Asp (59.3, 91.1)

MolProbity Ramachandran analysis

7acs.H.pdb, model 12









89.1% (123/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[12] 63 Asp (65.9, 99.3)
[12] 113 Asn (-170.1, -45.6)

MolProbity Ramachandran analysis

7acs.H.pdb, model 13









87.0% (120/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[13] 63 Asp (41.0, 87.2)

MolProbity Ramachandran analysis

7acs.H.pdb, model 14









90.6% (125/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):

[14] 37 Asn (-176.7, -27.8)
[14] 63 Asp (56.8, 89.3)

MolProbity Ramachandran analysis

7acs.H.pdb, model 15








90.6% (125/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[15] 66 Pro (-51.6, 103.0)

MolProbity Ramachandran analysis

7acs.H.pdb, model 16









90.6% (125/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[16] 63 Asp (59.0, 91.2)

MolProbity Ramachandran analysis

7acs.H.pdb, model 17









88.4% (122/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

7acs.H.pdb, model 18









88.4% (122/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

7acs.H.pdb, model 19









89.9% (124/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[19] 24 Leu (175.3, 133.5)
[19] 63 Asp (48.5, 90.6)

MolProbity Ramachandran analysis

7acs.H.pdb, model 20









89.9% (124/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[20] 63 Asp (46.8, 85.7)

MolProbity Ramachandran analysis

7acs.H.pdb, model 21









87.7% (121/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

7acs.H.pdb, model 22









91.3% (126/138) of all residues were in favored (98%) regions.
100.0% (138/138) of all residues were in allowed (>99.8%) regions.

There were no outliers.

MolProbity Ramachandran analysis

7acs.H.pdb, model 23








91.3% (126/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[23] 63 Asp (52.1, 86.2)

MolProbity Ramachandran analysis

7acs.H.pdb, model 24









91.3% (126/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[24] 63 Asp (65.8, 99.1)

MolProbity Ramachandran analysis

7acs.H.pdb, model 25









86.2% (119/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[25] 66 Pro (-52.9, 102.6)
[25] 123 Gln (-57.7, 98.3)

MolProbity Ramachandran analysis

7acs.H.pdb, model 26




89.1% (123/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[26] 14 Thr (-78.8, -142.1)
[26] 63 Asp (50.1, 82.9)

MolProbity Ramachandran analysis

7acs.H.pdb, model 27









85.5% (118/138) of all residues were in favored (98%) regions.
98.6% (136/138) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):

[27] 24 Leu (169.8, 137.5)
[27] 63 Asp (63.4, 105.2)

MolProbity Ramachandran analysis

7acs.H.pdb, model 28









87.7% (121/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[28] 63 Asp (64.4, 99.3)

MolProbity Ramachandran analysis

7acs.H.pdb, model 29









83.3% (115/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[29] 63 Asp (48.7, 82.8)

MolProbity Ramachandran analysis

7acs.H.pdb, model 30



91.3% (126/138) of all residues were in favored (98%) regions.
99.3% (137/138) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[30] 37 Asn (-173.2, -28.2)