






MolProbity Ramachandran analysis

7act.H.pdb, all models



85.5% (1154/1350) of all residues were in favored (98%) regions.
96.8% (1307/1350) of all residues were in allowed (>99.8%) regions.







There were 43 outliers (phi, psi):

[1] 49 Thr (58.5, -31.6)	[2] 162 Pro (-37.7, 169.6)
[1] 68 Arg (54.6, -6.0)	[3] 49 Thr (61.3, -21.8)
[1] 71 Gly (-179.3, 21.5)	[3] 71 Gly (-169.2, 20.1)
[1] 94 Ile (61.3, 124.9)	[3] 94 Ile (68.4, 136.8)
[1] 103 Asp (56.0, -0.3)	[4] 60 Gly (-66.0, -126.8)
[1] 140 Asn (-64.5, 24.4)	[4] 68 Arg (42.9, 16.5)
[2] 71 Gly (-167.4, -1.3)	[4] 71 Gly (161.9, 10.8)
[2] 94 Ile (65.6, 92.1)	[4] 94 Ile (70.8, 156.3)
	[5] 94 Ile (60.2, 97.0)
	[5] 96 Gly (66.9, 117.4)
	[5] 155 Ala (-32.5, 150.3)
	[6] 60 Gly (-73.0, -126.9)
	[6] 64 Leu (-65.7, 42.9)
	[6] 71 Gly (170.1, 35.0)
	[6] 94 Ile (70.5, 145.0)
	[6] 96 Gly (176.2, 85.7)
	[6] 158 Val (-52.8, 106.4)

[7] 71 Gly (174.9, 25.7)	[10] 71 Gly (159.2, 28.9)
[7] 94 Ile (68.1, 160.9)	[10] 83 Gln (-65.0, 73.8)
[7] 158 Val (-53.5, 97.8)	[10] 94 Ile (57.9, 145.1)
[7] 162 Pro (-43.3, 160.4)	
[8] 49 Thr (46.2, -50.2)	
[8] 70 Gln (-65.4, 72.4)	
[8] 71 Gly (-175.2, 34.0)	
[8] 94 Ile (64.6, 157.8)	
[8] 140 Asn (-64.5, 59.6)	
[9] 49 Thr (58.5, -29.3)	
[9] 71 Gly (157.1, 38.2)	
[9] 94 Ile (65.9, 143.7)	

MolProbity Ramachandran analysis

7act.H.pdb, model 1









87.4% (118/135) of all residues were in favored (98%) regions.
95.6% (129/135) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [1] 49 Thr (58.5, -31.6)
- [1] 68 Arg (54.6, -6.0)
- [1] 71 Gly (-179.3, 21.5)
- [1] 94 Ile (61.3, 124.9)
- [1] 103 Asp (56.0, -0.3)
- [1] 140 Asn (-64.5, 24.4)

MolProbity Ramachandran analysis

7act.H.pdb, model 2









87.4% (118/135) of all residues were in favored (98%) regions.
97.8% (132/135) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

- [2] 71 Gly (-167.4, -1.3)
- [2] 94 Ile (65.6, 92.1)
- [2] 162 Pro (-37.7, 169.6)

MolProbity Ramachandran analysis

7act.H.pdb, model 3









87.4% (118/135) of all residues were in favored (98%) regions.
97.8% (132/135) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[3] 49 Thr (61.3, -21.8)
[3] 71 Gly (-169.2, 20.1)
[3] 94 Ile (68.4, 136.8)

MolProbity Ramachandran analysis

7act.H.pdb, model 4









83.7% (113/135) of all residues were in favored (98%) regions.
96.3% (130/135) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [4] 60 Gly (-66.0, -126.8)
- [4] 68 Arg (42.9, 16.5)
- [4] 71 Gly (161.9, 10.8)
- [4] 94 Ile (70.8, 156.3)
- [4] 140 Asn (-48.0, -12.5)

MolProbity Ramachandran analysis

7act.H.pdb, model 5









83.0% (112/135) of all residues were in favored (98%) regions.
96.3% (130/135) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[5] 49 Thr (57.2, -29.8)
[5] 71 Gly (174.8, 13.7)
[5] 94 Ile (60.2, 97.0)
[5] 96 Gly (66.9, 117.4)
[5] 155 Ala (-32.5, 150.3)

MolProbity Ramachandran analysis

7act.H.pdb, model 6









83.7% (113/135) of all residues were in favored (98%) regions.
95.6% (129/135) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

[6] 60 Gly (-73.0, -126.9)
[6] 64 Leu (-65.7, 42.9)
[6] 71 Gly (170.1, 35.0)
[6] 94 Ile (70.5, 145.0)
[6] 96 Gly (176.2, 85.7)
[6] 158 Val (-52.8, 106.4)

MolProbity Ramachandran analysis

7act.H.pdb, model 7








85.9% (116/135) of all residues were in favored (98%) regions.
97.0% (131/135) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [7] 71 Gly (174.9, 25.7)
- [7] 94 Ile (68.1, 160.9)
- [7] 158 Val (-53.5, 97.8)
- [7] 162 Pro (-43.3, 160.4)

MolProbity Ramachandran analysis

7act.H.pdb, model 8









83.0% (112/135) of all residues were in favored (98%) regions.
96.3% (130/135) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [8] 49 Thr (46.2, -50.2)
- [8] 70 Gln (-65.4, 72.4)
- [8] 71 Gly (-175.2, 34.0)
- [8] 94 Ile (64.6, 157.8)
- [8] 140 Asn (-64.5, 59.6)

MolProbity Ramachandran analysis

7act.H.pdb, model 9









88.1% (119/135) of all residues were in favored (98%) regions.
97.8% (132/135) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

[9] 49 Thr (58.5, -29.3)
[9] 71 Gly (157.1, 38.2)
[9] 94 Ile (65.9, 143.7)

MolProbity Ramachandran analysis

7act.H.pdb, model 10



85.2% (115/135) of all residues were in favored (98%) regions.
97.8% (132/135) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

- [10] 71 Gly (159.2, 28.9)
- [10] 83 Gln (-65.0, 73.8)
- [10] 94 Ile (57.9, 145.1)