


MolProbity Ramachandran analysis

2jw8.H.pdb, all models



89.1% (4133/4640) of all residues were in favored (98%) regions.
98.3% (4563/4640) of all residues were in allowed (>99.8%) regions.
This list is truncated; use the MolProbity multi-chart.html for complete list.







There were 77 outliers (phi, psi):

[1] A 301 His (58.2, -10.2)	[2] B 342 Asp (-61.8, 17.3)	[4] B 280 Pro (-65.8, -62.0)
[1] A 317 Gly (-148.8, -51.5)	[3] A 288 Gly (-176.4, 75.7)	[5] A 249 Lys (-40.5, 98.8)
[1] A 319 Ser (-70.0, 27.6)	[3] A 319 Ser (-63.2, 28.3)	[5] A 252 Ala (-71.3, 32.9)
[1] B 342 Asp (-59.3, 8.8)	[3] A 342 Asp (-48.7, 73.1)	[5] B 342 Asp (34.3, 77.3)
[2] A 288 Gly (154.9, 80.8)	[3] B 278 Arg (-70.3, -87.9)	[6] A 278 Arg (-46.2, 100.0)
[2] A 296 Gly (44.9, -104.6)	[3] B 342 Asp (-32.6, 6.5)	[6] A 317 Gly (-179.9, -83.5)
[2] B 288 Gly (170.2, 81.5)	[4] A 317 Gly (172.9, -78.5)	[6] A 319 Ser (-67.9, 28.2)
[2] B 318 Met (-67.5, 37.0)	[4] B 256 Ser (169.1, 157.8)	[6] B 288 Gly (142.2, 75.9)
		[7] A 255 Ala (-65.7, -81.8)
		[7] A 257 Lys (-174.2, -60.9)
		[7] A 267 Lys (-47.8, 3.0)

[7] A 288 Gly (168.8, 81.8)	[11] A 253 Ala (-51.2, -78.2)	[14] B 263 Arg (-20.1, 101.5)
[7] A 319 Ser (-65.3, 24.5)	[11] A 319 Ser (-65.0, 23.2)	[14] B 288 Gly (145.6, 82.1)
[7] A 342 Asp (-54.1, 2.5)	[11] B 317 Gly (-163.6, -74.3)	[14] B 317 Gly (-164.2, -70.3)
[8] A 317 Gly (-178.6, -61.7)	[12] A 249 Lys (122.7, 88.2)	[15] A 267 Lys (-56.4, 0.3)
[8] B 288 Gly (-179.6, 80.2)	[12] A 288 Gly (163.9, 93.4)	[15] A 318 Met (-59.9, 6.2)
[9] A 288 Gly (173.2, 73.0)	[12] A 317 Gly (-150.8, -75.4)	[15] B 280 Pro (-79.4, -52.1)
[9] A 319 Ser (-65.6, 28.2)	[12] A 319 Ser (-67.2, 26.7)	[16] A 288 Gly (164.2, 87.0)
[10] A 288 Gly (169.5, 91.4)	[12] B 286 Asn (51.3, -16.5)	[16] B 249 Lys (52.2, 5.6)
[10] A 319 Ser (-65.9, 27.7)	[12] B 319 Ser (-65.6, 27.0)	[16] B 261 Gln (-163.3, -12.7)
[10] B 253 Ala (-56.0, -78.4)	[13] B 320 Arg (30.8, 79.5)	[16] B 342 Asp (-68.3, 42.9)
[10] B 285 Gly (149.6, 78.9)	[14] A 285 Gly (149.6, 85.6)	[17] A 317 Gly (-168.7, -53.9)
[10] B 288 Gly (-176.3, 79.7)	[14] A 288 Gly (172.6, 81.0)	[17] A 342 Asp (-63.8, 52.2)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 1









88.4% (205/232) of all residues were in favored (98%) regions.
98.3% (228/232) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [1] A 301 His (58.2, -10.2)
- [1] A 317 Gly (-148.8, -51.5)
- [1] A 319 Ser (-70.0, 27.6)
- [1] B 342 Asp (-59.3, 8.8)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 2









89.2% (207/232) of all residues were in favored (98%) regions.
97.8% (227/232) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [2] A 288 Gly (154.9, 80.8)
- [2] A 296 Gly (44.9, -104.6)
- [2] B 288 Gly (170.2, 81.5)
- [2] B 318 Met (-67.5, 37.0)
- [2] B 342 Asp (-61.8, 17.3)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 3









86.6% (201/232) of all residues were in favored (98%) regions.
97.8% (227/232) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [3] A 288 Gly (-176.4, 75.7)
- [3] A 319 Ser (-63.2, 28.3)
- [3] A 342 Asp (-48.7, 73.1)
- [3] B 278 Arg (-70.3, -87.9)
- [3] B 342 Asp (-32.6, 6.5)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 4









90.9% (211/232) of all residues were in favored (98%) regions.
98.7% (229/232) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):
[4] A 317 Gly (172.9, -78.5)
[4] B 256 Ser (169.1, 157.8)
[4] B 280 Pro (-65.8, -62.0)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 5









90.9% (211/232) of all residues were in favored (98%) regions.
98.3% (228/232) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [5] A 249 Lys (-40.5, 98.8)
- [5] A 252 Ala (-71.3, 32.9)
- [5] A 288 Gly (-176.8, 69.2)
- [5] B 342 Asp (34.3, 77.3)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 6









87.9% (204/232) of all residues were in favored (98%) regions.
98.3% (228/232) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):
[6] A 278 Arg (-46.2, 100.0)
[6] A 317 Gly (-179.9, -83.5)
[6] A 319 Ser (-67.9, 28.2)
[6] B 288 Gly (142.2, 75.9)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 7








89.7% (208/232) of all residues were in favored (98%) regions.
97.4% (226/232) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [7] A 255 Ala (-65.7, -81.8)
- [7] A 257 Lys (-174.2, -60.9)
- [7] A 267 Lys (-47.8, 3.0)
- [7] A 288 Gly (168.8, 81.8)
- [7] A 319 Ser (-65.3, 24.5)
- [7] A 342 Asp (-54.1, 2.5)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 8









90.9% (211/232) of all residues were in favored (98%) regions.
99.1% (230/232) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[8] A 317 Gly (-178.6, -61.7)
[8] B 288 Gly (-179.6, 80.2)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 9









91.8% (213/232) of all residues were in favored (98%) regions.
99.1% (230/232) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[9] A 288 Gly (173.2, 73.0)
[9] A 319 Ser (-65.6, 28.2)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 10









87.1% (202/232) of all residues were in favored (98%) regions.
97.8% (227/232) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [10] A 288 Gly (169.5, 91.4)
- [10] A 319 Ser (-65.9, 27.7)
- [10] B 253 Ala (-56.0, -78.4)
- [10] B 285 Gly (149.6, 78.9)
- [10] B 288 Gly (-176.3, 79.7)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 11









87.5% (203/232) of all residues were in favored (98%) regions.
98.7% (229/232) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):

- [11] A 253 Ala (-51.2, -78.2)
- [11] A 319 Ser (-65.0, 23.2)
- [11] B 317 Gly (-163.6, -74.3)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 12









88.8% (206/232) of all residues were in favored (98%) regions.
97.4% (226/232) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [12] A 249 Lys (122.7, 88.2)
- [12] A 288 Gly (163.9, 93.4)
- [12] A 317 Gly (-150.8, -75.4)
- [12] A 319 Ser (-67.2, 26.7)
- [12] B 286 Asn (51.3, -16.5)
- [12] B 319 Ser (-65.6, 27.0)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 13









90.1% (209/232) of all residues were in favored (98%) regions.
99.6% (231/232) of all residues were in allowed (>99.8%) regions.

There were 1 outliers (phi, psi):
[13] B 320 Arg (30.8, 79.5)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 14









89.7% (208/232) of all residues were in favored (98%) regions.
97.8% (227/232) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [14] A 285 Gly (149.6, 85.6)
- [14] A 288 Gly (172.6, 81.0)
- [14] B 263 Arg (-20.1, 101.5)
- [14] B 288 Gly (145.6, 82.1)
- [14] B 317 Gly (-164.2, -70.3)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 15








87.1% (202/232) of all residues were in favored (98%) regions.
98.7% (229/232) of all residues were in allowed (>99.8%) regions.

There were 3 outliers (phi, psi):
[15] A 267 Lys (-56.4, 0.3)
[15] A 318 Met (-59.9, 6.2)
[15] B 280 Pro (-79.4, -52.1)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 16









89.7% (208/232) of all residues were in favored (98%) regions.
98.3% (228/232) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

- [16] A 288 Gly (164.2, 87.0)
- [16] B 249 Lys (52.2, 5.6)
- [16] B 261 Gln (-163.3, -12.7)
- [16] B 342 Asp (-68.3, 42.9)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 17









87.5% (203/232) of all residues were in favored (98%) regions.
97.8% (227/232) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

[17] A 317 Gly (-168.7, -53.9)
[17] A 342 Asp (-63.8, 52.2)
[17] B 251 Ser (-62.1, 81.5)
[17] B 285 Gly (139.5, 74.1)
[17] B 317 Gly (-170.9, -66.0)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 18









87.9% (204/232) of all residues were in favored (98%) regions.
98.3% (228/232) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):
[18] A 288 Gly (176.8, 72.8)
[18] B 267 Lys (-52.7, -3.2)
[18] B 290 Gln (44.0, -87.0)
[18] B 319 Ser (-65.2, 14.3)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 19









88.8% (206/232) of all residues were in favored (98%) regions.
99.1% (230/232) of all residues were in allowed (>99.8%) regions.

There were 2 outliers (phi, psi):
[19] B 279 Gly (-6.0, 112.5)
[19] B 318 Met (-57.1, 7.2)

MolProbity Ramachandran analysis

2jw8.H.pdb, model 20



90.9% (211/232) of all residues were in favored (98%) regions.
98.3% (228/232) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[20] A 317 Gly (-178.5, -64.7)
[20] B 317 Gly (-166.9, -39.0)
[20] B 319 Ser (-67.0, 28.7)
[20] B 320 Arg (37.0, 99.0)