


MolProbity Ramachandran analysis

7fae.H.pdb, model 1



90.1% (3831/4250) of all residues were in favored (98%) regions.
99.3% (4220/4250) of all residues were in allowed (>99.8%) regions.

There were 30 outliers (phi, psi):

A 591 Ser (-21.6, 137.3)	B 325 Ser (-170.8, 42.6)	H 157 Glu (-82.6, 67.0)	L 94 Tyr (-77.9, -166.2)
A 810 Ser (-173.2, 39.4)	B 527 Pro (-94.3, -127.9)	H 158 Pro (-132.0, 70.4)	L 95 Pro (-29.4, 121.0)
A 811 Lys (-148.2, -162.2)	C 88 Asp (-162.9, 0.3)	d 142 Arg (8.1, -85.3)	
A 812 Pro (-51.6, 99.9)	C 387 Leu (-36.5, -32.1)	d 143 Glu (-30.4, 82.6)	
A 813 Ser (67.7, -85.1)	C 520 Ala (-155.8, -60.2)	e 155 Phe (-86.1, -89.6)	
A 814 Lys (172.6, 74.2)	C 521 Pro (-25.1, -74.0)	e 156 Pro (-80.9, -153.8)	
A 986 Pro (-5.7, -66.8)	C 528 Lys (156.2, -10.4)	e 159 Val (10.0, 74.6)	
B 324 Glu (77.6, -152.0)	H 155 Phe (-85.6, -93.9)	g 94 Tyr (-77.9, -166.2)	
		g 95 Pro (-29.5, 121.0)	
		h 155 Phe (-126.4, -86.4)	