


MolProbity Ramachandran analysis

8aou.H.pdb, all models



79.4% (1421/1790) of all residues were in favored (98%) regions.

95.8% (1714/1790) of all residues were in allowed (>99.8%) regions.

This list is truncated; use the MolProbity multi-chart.html for complete list.







There were 76 outliers (phi, psi):

[1] 4 Leu (71.2, 122.4)	[2] 95 Ile (-29.8, -47.2)	[3] 155 Glu (33.0, 77.8)
[1] 28 Val (70.8, 78.6)	[2] 113 Glu (170.5, 150.8)	[3] 176 Glu (64.8, 106.7)
[1] 98 Gly (138.0, 63.8)	[2] 155 Glu (36.3, 76.6)	[4] 4 Leu (172.6, 128.8)
[1] 113 Glu (169.1, 151.2)	[3] 4 Leu (174.5, 135.0)	[4] 28 Val (71.5, 82.1)
[1] 132 Gly (-163.8, -53.0)	[3] 7 Gly (158.7, 43.3)	[4] 82 Gly (-153.1, -60.2)
[2] 4 Leu (67.1, 110.1)	[3] 28 Val (70.4, 76.3)	[4] 139 Asp (-171.5, 48.9)
[2] 6 Pro (-56.6, 96.5)	[3] 98 Gly (-130.7, -52.6)	[4] 144 Asp (51.0, 86.5)
[2] 28 Val (68.0, 81.9)	[3] 113 Glu (166.6, 150.4)	[4] 157 Phe (47.2, -96.7)
		[4] 159 Glu (-179.2, 109.5)
		[4] 28 Val (68.8, 84.0)
		[4] 177 Leu (64.9, -75.8)

[5] 6 Pro (-59.4, 91.0)	[6] 98 Gly (-163.8, -4.5)	[7] 143 Phe (-78.3, -114.0)
[5] 28 Val (67.9, 74.1)	[6] 113 Glu (175.6, 131.8)	[7] 144 Asp (55.3, 97.1)
[5] 75 Asp (58.8, -76.3)	[6] 147 Asp (62.3, 99.4)	[7] 146 Gly (-139.7, -50.2)
[5] 78 Thr (34.5, 76.2)	[6] 150 Gly (171.7, -21.0)	[7] 159 Glu (-172.5, 23.1)
[5] 113 Glu (167.2, 135.9)	[6] 155 Gly (25.3, 72.3)	[7] 163 Thr (60.1, 126.3)
[5] 171 Arg (-179.3, -25.0)	[6] 157 Phe (-171.4, -11.8)	[7] 175 Arg (70.1, 137.7)
[5] 178 Asn (-172.8, -50.4)	[6] 169 Val (66.3, 89.3)	[8] 4 Leu (66.7, 112.6)
[6] 4 Leu (73.4, 133.5)	[6] 172 Glu (-62.3, -82.9)	[8] 6 Pro (-47.3, 105.3)
[6] 6 Pro (-50.9, 100.6)	[7] 4 Leu (69.5, 121.4)	[8] 28 Val (67.7, 77.4)
[6] 8 Phe (68.7, 127.7)	[7] 28 Val (70.1, 81.6)	[8] 81 His (175.8, -76.6)
[6] 28 Val (68.8, 84.0)	[7] 80 Pro (-58.9, -174.7)	[8] 146 Gly (176.2, -65.4)
[6] 81 His (59.1, -91.5)	[7] 132 Gly (-176.4, 78.5)	[8] 159 Glu (55.2, 94.3)

MolProbity Ramachandran analysis

8aou.H.pdb, model 1









78.8% (141/179) of all residues were in favored (98%) regions.
97.2% (174/179) of all residues were in allowed (>99.8%) regions.

There were 5 outliers (phi, psi):

- [1] 4 Leu (71.2, 122.4)
- [1] 28 Val (70.8, 78.6)
- [1] 98 Gly (138.0, 63.8)
- [1] 113 Glu (169.1, 151.2)
- [1] 132 Gly (-163.8, -53.0)

MolProbity Ramachandran analysis

8aou.H.pdb, model 2









80.4% (144/179) of all residues were in favored (98%) regions.
96.6% (173/179) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [2] 4 Leu (67.1, 110.1)
- [2] 6 Pro (-56.6, 96.5)
- [2] 28 Val (68.0, 81.9)
- [2] 95 Ile (-29.8, -47.2)
- [2] 113 Glu (170.5, 150.8)
- [2] 155 Glu (36.3, 76.6)

MolProbity Ramachandran analysis

8aou.H.pdb, model 3









81.0% (145/179) of all residues were in favored (98%) regions.
96.1% (172/179) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

[3] 4 Leu (174.5, 135.0)
[3] 7 Gly (158.7, 43.3)
[3] 28 Val (70.4, 76.3)
[3] 98 Gly (-130.7, -52.6)
[3] 113 Glu (166.6, 150.4)
[3] 155 Glu (33.0, 77.8)
[3] 176 Glu (64.8, 106.7)

MolProbity Ramachandran analysis

8aou.H.pdb, model 4









79.9% (143/179) of all residues were in favored (98%) regions.
94.4% (169/179) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[4] 4 Leu (172.6, 128.8)	[4] 164 Lys (-160.6, -43.8)
[4] 7 Gly (-174.7, -64.4)	[4] 177 Leu (64.9, -75.8)
[4] 28 Val (71.5, 82.1)	
[4] 82 Gly (-153.1, -60.2)	
[4] 139 Asp (-171.5, 48.9)	
[4] 144 Asp (51.0, 86.5)	
[4] 157 Phe (47.2, -96.7)	
[4] 159 Glu (-179.2, 109.5)	

MolProbity Ramachandran analysis

8aou.H.pdb, model 5









79.9% (143/179) of all residues were in favored (98%) regions.
96.1% (172/179) of all residues were in allowed (>99.8%) regions.

There were 7 outliers (phi, psi):

- [5] 6 Pro (-59.4, 91.0)
- [5] 28 Val (67.9, 74.1)
- [5] 75 Asp (58.8, -76.3)
- [5] 78 Thr (34.5, 76.2)
- [5] 113 Glu (167.2, 135.9)
- [5] 171 Arg (-179.3, -25.0)
- [5] 178 Asn (-172.8, -50.4)

MolProbity Ramachandran analysis

8aou.H.pdb, model 6









77.7% (139/179) of all residues were in favored (98%) regions.
92.7% (166/179) of all residues were in allowed (>99.8%) regions.

There were 13 outliers (phi, psi):

[6] 4 Leu (73.4, 133.5)	[6] 150 Gly (171.7, -21.0)
[6] 6 Pro (-50.9, 100.6)	[6] 155 Glu (25.3, 72.3)
[6] 8 Phe (68.7, 127.7)	[6] 157 Phe (-171.4, -11.8)
[6] 28 Val (68.8, 84.0)	[6] 169 Val (66.3, 89.3)
[6] 81 His (59.1, -91.5)	[6] 172 Glu (-62.3, -82.9)
[6] 98 Gly (-163.8, -4.5)	
[6] 113 Glu (175.6, 131.8)	
[6] 147 Asp (62.3, 99.4)	

MolProbity Ramachandran analysis

8aou.H.pdb, model 7









75.4% (135/179) of all residues were in favored (98%) regions.
94.4% (169/179) of all residues were in allowed (>99.8%) regions.

There were 10 outliers (phi, psi):

[7] 4 Leu (69.5, 121.4)	[7] 163 Thr (60.1, 126.3)
[7] 28 Val (70.1, 81.6)	[7] 175 Arg (70.1, 137.7)
[7] 80 Pro (-58.9, -174.7)	
[7] 132 Gly (-176.4, 78.5)	
[7] 143 Phe (-78.3, -114.0)	
[7] 144 Asp (55.3, 97.1)	
[7] 146 Gly (-139.7, -50.2)	
[7] 159 Glu (-172.5, 23.1)	

MolProbity Ramachandran analysis

8aou.H.pdb, model 8









81.6% (146/179) of all residues were in favored (98%) regions.
95.5% (171/179) of all residues were in allowed (>99.8%) regions.

There were 8 outliers (phi, psi):

- [8] 4 Leu (66.7, 112.6)
- [8] 6 Pro (-47.3, 105.3)
- [8] 28 Val (67.7, 77.4)
- [8] 81 His (175.8, -76.6)
- [8] 146 Gly (176.2, -65.4)
- [8] 159 Glu (55.2, 94.3)
- [8] 171 Arg (-169.4, 27.8)
- [8] 178 Asn (-163.7, -50.4)

MolProbity Ramachandran analysis

8aou.H.pdb, model 9




79.3% (142/179) of all residues were in favored (98%) regions.
96.6% (173/179) of all residues were in allowed (>99.8%) regions.

There were 6 outliers (phi, psi):

- [9] 2 Glu (61.0, -90.1)
- [9] 28 Val (70.7, 80.9)
- [9] 113 Glu (167.1, 156.6)
- [9] 144 Asp (-175.1, 55.5)
- [9] 147 Asp (23.3, 62.1)
- [9] 161 Trp (-81.3, -114.3)

MolProbity Ramachandran analysis

8aou.H.pdb, model 10



79.9% (143/179) of all residues were in favored (98%) regions.
97.8% (175/179) of all residues were in allowed (>99.8%) regions.

There were 4 outliers (phi, psi):

[10] 28 Val (72.3, 81.0)
[10] 81 His (174.0, -81.8)
[10] 144 Asp (177.6, 69.7)
[10] 178 Asn (64.4, -75.2)