

# Wilson–’t Hooft lines as transfer matrices

Junya Yagi

Yau Mathematical Sciences Center, Tsinghua University

December 3, 2020

NTU–Kyoto High Energy Physics Workshop

Based on joint work with Kazunobu Maruyoshi and Toshihiro Ota

Various connections between

supersymmetric QFTs  $\longleftrightarrow$  quantum integrable systems

discovered in the past 10 years or so:

- Bethe/gauge correspondence (2d & 4d) [Nekrasov–Shatashvili]
- Bazhanov–Sergeev model from 4d  $\mathcal{N} = 1$  quiver gauge theories [Spiridonov, Yamazaki]
- Surface defects as transfer matrices [Maruyoshi–Yagi]
- 4d Chern–Simons (=  $\Omega$ -deformed 6d SYM [Costello–Y])  
[Costello, Costello–Yamazaki–Witten]
- ...

Many of them are related by string dualities [Costello–Y].

## Quantization of Donagi–Witten integrable system

- $\mathcal{N} = 2$  theory on  $\mathbb{R}^3 \times S^1$  on Coulomb branch
- IR:  $\mathcal{N} = 4$  sigma model on  $\mathbb{R}^3$
- Target  $\mathcal{M}$  is the phase space of a classical **complex integrable system** [Donagi–Witten]
- **$\Omega$ -deformation** on  $\mathbb{R}^2 \subset \mathbb{R}^3$  quantizes  $\mathcal{M}$   
[Nekrasov–Shatashvili, Nekrasov–Witten, Y]
- For class- $\mathcal{S}$  theories,  $\mathcal{M}$  is a Hitchin system.

## Surface defects as transfer matrices [Maruyoshi–Y, Y]

- $\mathcal{N} = 1$  theory constructed by “brane tiling” or of class  $\mathcal{S}_k$
- Place it on  $S^3 \times S^1$
- Insert **surface defects** on  $S^1 \times S^1$
- Surface defects act on SUSY index as difference operators, shifting flavor fugacities [Gadde–Gukov, Gaiotto–Rastelli–Razamat]
- Coincide with **transfer matrices** of elliptic QIS [Maruyoshi–Y, Y]
- Simplest case: **elliptic** Ruijsenaars–Schneider system [GRR, Bullimore–Fluder–Hollands–Richmond]


We found a new correspondence:

Wilson–’t Hooft lines = transfer matrices

- $\mathcal{N} = 2$  circular quiver theory (**class- $\mathcal{S}$** )
- Place it on  $S^1 \times \mathbb{R}^3$
- Wind a **Wilson–’t Hooft line**  $T$  around  $S^1$
- $\langle T \rangle$  is a function of Coulomb branch parameters
- Quantization of  $\langle T \rangle$  coincides with transfer matrix of **trigonometric** QIS

Related to other correspondences

Consider a periodic spin chain



Spins  $a^1, \dots, a^n \in \mathfrak{h}^*$ ,  $\mathfrak{h} = \text{Cartan of } \mathfrak{sl}_N$ :

$$a^r = \text{diag}(a_1^r, \dots, a_N^r), \quad \sum_{i=1}^N a_i^r = 0$$


Local Hilbert space:

$$\mathcal{M}_{\mathfrak{h}^*} = \{\text{meromorphic functions on } \mathfrak{h}^*\}$$


Total Hilbert space

$$\mathcal{H} = \underbrace{\mathcal{M}_{\mathfrak{h}^*} \otimes \dots \otimes \mathcal{M}_{\mathfrak{h}^*}}_n$$

## Equivalent lattice model



Spins live between double lines:



$a^r$  are called **dynamical parameters**.

$$L(z) = z \quad \text{---} \quad \begin{array}{c} \uparrow \\ \downarrow \end{array}$$



Dynamical parameters jump across solid lines:

$$L(z; a^1, a^2)_i^j = z \begin{array}{c} \xrightarrow{a^1} \textcircled{i} \xrightarrow{a^2} \textcircled{j} \xrightarrow{\quad} \\ \xleftarrow{a^1 - \epsilon h_i} \quad \quad \quad \xleftarrow{a^2 - \epsilon h_j} \end{array} \quad .$$

$\epsilon \in \mathbb{C}$ : fixed parameter (Planck constant)

$h_i$  are the weights of the vector rep  $\mathbb{C}^N$ :

$$\begin{aligned} h_1 &= \text{diag}(1 - \frac{1}{N}, -\frac{1}{N}, -\frac{1}{N}, \dots, -\frac{1}{N}), \\ h_2 &= \text{diag}(-\frac{1}{N}, 1 - \frac{1}{N}, -\frac{1}{N}, \dots, -\frac{1}{N}), \\ &\vdots \\ h_N &= \text{diag}(-\frac{1}{N}, -\frac{1}{N}, -\frac{1}{N}, \dots, 1 - \frac{1}{N}). \end{aligned}$$

$$L(z; a^1, a^2)_i^j = z \begin{array}{c} a^1 \\ \text{---} \bigcirc i \text{---} \\ a^1 - \epsilon h_i \end{array} \begin{array}{c} \updownarrow \\ \text{---} \bigcirc j \text{---} \\ a^2 - \epsilon h_j \end{array} \begin{array}{c} a^2 \\ \end{array}$$

Matrix elements  $L(z)_i^j$  are **difference operators** on  $\mathcal{M}_{\mathfrak{h}^*} \otimes \mathcal{M}_{\mathfrak{h}^*}$ :

$$L(z) = \sum_{i,j} L(z; a^1, a^2)_i^j \Delta_i^1 \Delta_j^2,$$

$$\Delta_i^r : a^r \mapsto a^r - \epsilon h_i.$$

Transfer matrix

$$T(z) = \sum_{i^1, \dots, i^n} \prod_{r=1}^n L(z; a^r, a^{r+1})_{i^r}^{i^{r+1}} \prod_{s=1}^n \Delta_{i^s}^s, \quad i^{n+1} = i^1$$

is a difference operator on  $\mathcal{H} = \mathcal{M}_{\mathfrak{h}^*}^{\otimes n}$ .

Crossing solid lines give **R-matrix**

$$R(z - z'; a)^{kl}_{ij} = \begin{array}{c} \uparrow \\ \textcircled{l} \\ a \\ \textcircled{i} \text{---} \textcircled{k} \\ \downarrow \\ \textcircled{j} \\ z' \end{array} .$$

R-matrix satisfies **dynamical Yang-Baxter equation**


$$\begin{array}{c} 1 \\ \diagdown \\ \diagup \\ 2 \end{array} \begin{array}{c} \uparrow \\ \diagup \\ \diagdown \\ 3 \end{array} = \begin{array}{c} 1 \\ \diagup \\ \diagdown \\ 2 \end{array} \begin{array}{c} \uparrow \\ \diagdown \\ \diagup \\ 3 \end{array} .$$

Just like the ordinary Yang-Baxter equation

$$R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12}$$


but with shifts in the dynamical parameters.

L-operator and R-matrix satisfy **RLL relation**




It follows that transfer matrices commute:

$$T(z)T(z') = T(z')T(z)$$




*Proof:*

By RLL relation



Multiply both sides by  $R^{-1}$ :



Take the trace, making the horizontal direction periodic.

Since

$$[T(z), T(z')] = 0,$$

coefficients of Laurent expansion

$$T(z) = \sum_{m=-\infty}^{\infty} T_m z^m$$

are commuting difference operators on  $\mathcal{H}$ :

$$[T_m, T_n] = 0.$$

This is **integrability**.

## Trigonometric L-operator [Hasegawa]

$$\mathcal{L}_{w,m}(z)_i^j = \sum_{i,j} (\Delta_i^1 \Delta_j^2)^{\frac{1}{2}} \frac{\sin \pi(z - w + a_j^2 - a_i^1)}{\sin \pi(z - w)} \ell_m(a^1, a^2)_i^j (\Delta_i^1 \Delta_j^2)^{\frac{1}{2}}$$

satisfies RLL relation with a trigonometric dynamical R-matrix (a limit of the 8vSOS R-matrix).

$$\ell_m(a^1, a^2)_i^j = \left( \frac{\prod_{k(\neq i)} \sin \pi(a_k^1 - a_j^2 - m) \prod_{l(\neq j)} \sin \pi(a_i^1 - a_l^2 - m)}{\prod_{k(\neq i)} \sin \pi(a_{ki}^1 - \frac{1}{2}\epsilon) \sin \pi(a_{ik}^1 - \frac{1}{2}\epsilon)} \right)^{\frac{1}{2}}$$

$w, m \in \mathbb{C}$  are spectral parameters assigned to the double line:

$$\mathcal{L}_{w,m}(z) = z \begin{array}{c} \uparrow \\ \text{---} \text{---} \text{---} \\ \downarrow \\ w, m \end{array} .$$

## Introduce **fundamental L-operators**

$$\mathcal{L}_{\pm,m} = \lim_{w \rightarrow \pm i\infty} \mathcal{L}_{w,m}.$$

Then

$$(\mathcal{L}_{\pm,m})_i^j = \sum_{i,j} (\Delta_i^1 \Delta_j^2)^{\frac{1}{2}} e^{\pm \pi i (a_j^2 - a_i^1)} \ell_m(a^1, a^2)_i^j (\Delta_i^1 \Delta_j^2)^{\frac{1}{2}}$$

and

$$\mathcal{L}_{w,m}(z) = \frac{e^{\pi i(z-w)} \mathcal{L}_{+,m} - e^{-\pi i(z-w)} \mathcal{L}_{-,m}}{\sin \pi(z-w)}.$$

We may as well consider  $\mathcal{L}_{\pm,m}$  without loss of generality.



Pick  $n$ -tuple of signs

$$\sigma = (\sigma^1, \dots, \sigma^n) \in \{\pm\}^n$$

and  $n$ -tuple of complex numbers

$$m = (m^1, \dots, m^n) \in \mathbb{C}^n .$$

Let  $\mathcal{T}_{\sigma,m}$  be the transfer matrix constructed from  $n$  L-operators

$$\mathcal{L}_{\sigma^1, m^1} , \dots , \mathcal{L}_{\sigma^n, m^n} .$$

$$\mathcal{T}_{\sigma,m} = \sum_{i^1, \dots, i^n} \left( \prod_{s=1}^n \Delta_{i_s}^s \right)^{\frac{1}{2}} \prod_{r=1}^n e^{\pi i \sigma^r (a_{i^{r+1}}^{r+1} - a_{i^r}^r)} \ell_{m^r}(a^r, a^{r+1})_{i^r}^{i^{r+1}} \left( \prod_{s=1}^n \Delta_{i_s}^s \right)^{\frac{1}{2}} .$$

This is the main character from the integrable system.

$\mathcal{N} = 2$  gauge theories have half-BPS **Wilson–’t Hooft lines**.

Worldlines of very massive dyonic particles

Charge of WH line

$$(\mathbf{m}, \mathbf{e}) \in (\Lambda_{\text{coweight}} \times \Lambda_{\text{weight}}) / \text{Weyl}.$$


Wilson line has  $\mathbf{m} = 0$  and is labeled by representation of  $\mathfrak{g}$ .

’t Hooft line has  $\mathbf{e} = 0$  and is labeled by representation of  ${}^L\mathfrak{g}$ .

Wilson–’t Hooft

= (’t Hooft) + (Wilson for subgroup of  $G$  leaving  $\mathbf{m}$  invariant)


$\mathcal{N} = 2$  gauge theory described by  $n$ -node **circular quiver**




Each node is  $SU(N)$  (more precisely,  $PSU(N)$ ).

Edges are bifundamental hypers with masses  $m^1, \dots, m^n$ .

Compactification of 6d  $\mathcal{N} = (2, 0)$  SCFT on  $n$ -punctured torus



WH lines = surface defects wrapping 1-cycles of the torus



Consider Wilson-'t Hooft line  $T_{\square, \sigma}$  corresponding to

$$\gamma_{\sigma} = b + \sum_r \frac{1 - \sigma^r}{2} c^r.$$

If  $\sigma^r = +1$  ( $-1$ ), the cycle passes above (below)  $r$ th puncture.

$\mathbf{m} = \square \oplus \cdots \oplus \square$  under  $\mathfrak{su}_N \oplus \cdots \oplus \mathfrak{su}_N$

$\mathbf{e}$  specified by  $\sigma \in \{\pm\}^n$

Put the theory on twisted product

$$S^1 \times_{\epsilon} \mathbb{R}^2 \times \mathbb{R}.$$

Wrap  $T_{\square, \sigma}$  around  $S^1 \times \{0\} \times \{t\}$ .

Ito–Okuda–Taki tell us how to compute the vev by **localization**:

$$\langle T_{\square, \sigma} \rangle = \sum_{i^1, \dots, i^n} \prod_{r=1}^n e^{2\pi i b_{i^r}^r} e^{\pi i \sigma^r (a_{i^r+1}^{r+1} - a_{i^r}^r)} \ell_{m^r}(a^r, a^{r+1})_{i^r}^{i^r+1}$$

in complexified Fenchel–Nielsen coordinates on Seiberg–Witten moduli space:

$$a = \frac{\theta_e}{2\pi} + i\beta \operatorname{Re} \phi + \dots, \quad b = \frac{\theta_m}{2\pi} - \frac{4\pi i \beta}{g^2} \operatorname{Im} \phi + i \frac{\vartheta}{2\pi} \beta \operatorname{Re} \phi + \dots.$$

Alternatively, we can compute it from **Toda theory** by AGT.

## Compare

$$\langle T_{\square, \sigma} \rangle = \sum_{i^1, \dots, i^n} \prod_{r=1}^n e^{2\pi i b_{i^r}^r} e^{\pi i \sigma^r (a_{i^r+1}^{r+1} - a_{i^r}^r)} \ell_{m^r}(a^r, a^{r+1})_{i^r}^{i^{r+1}},$$

$$\mathcal{T}_{\sigma, m} = \sum_{i^1, \dots, i^n} \left( \prod_{s=1}^n \Delta_{i_s}^s \right)^{\frac{1}{2}} \prod_{r=1}^n e^{\pi i \sigma^r (a_{i^r+1}^{r+1} - a_{i^r}^r)} \ell_{m^r}(a^r, a^{r+1})_{i^r}^{i^{r+1}} \left( \prod_{s=1}^n \Delta_{i_s}^s \right)^{\frac{1}{2}}.$$

If we quantize  $a^r, b^r$  so that

$$[\hat{a}_i^r, \hat{b}_j^s] = -i \frac{\epsilon}{2\pi} \delta^{rs} \left( \delta_{ij} - \frac{1}{N} \right),$$

then

$$\mathcal{T}_{\sigma, m} = \text{Weyl quantization of } \langle T_{\square, \sigma} \rangle.$$

# M-theory setup

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\mathbb{R}_9$	$S_{10}^1$
$N$ M5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	$S_6^1$	—	—	—	$S_{10}^1$
$n$ M5'	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	—	—	$\mathbb{R}_8$	$\mathbb{R}_9$	—
M2	—	—	$S_3^1$	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	—	—

12345 directions: twisted product  $\mathbb{R}_{12}^2 \times_{\epsilon} S_3^1 \times_{-\epsilon} \mathbb{R}_{45}^2$

M5: 6d  $\mathcal{N} = (2, 0)$  SCFT on  $\mathbb{R}_0 \times \mathbb{R}_{12}^2 \times_{\epsilon} S_3^1 \times S_6^1 \times S_{10}^1$

M5':  $n$  punctures on  $S_6^1 \times S_{10}^1$

M2: surface defect

Reduction on  $S_6^1 \times S_{10}^1$  gives the 4d setup with  $\sigma = (+, \dots, +)$ .

Compactify  $\mathbb{R}_9 \rightarrow S^1_9$ :

Spacetime	$\mathbb{R}_0$	$\mathbb{R}^2_{12}$	$S^1_3$	$\mathbb{R}^2_{45}$	$S^1_6$	$\mathbb{R}_7$	$\mathbb{R}_8$	$S^1_9$	$S^1_{10}$
$N$ M5	$\mathbb{R}_0$	$\mathbb{R}^2_{12}$	$S^1_3$	—	$S^1_6$	—	—	—	$S^1_{10}$
$n$ M5'	$\mathbb{R}_0$	$\mathbb{R}^2_{12}$	$S^1_3$	—	—	—	$\mathbb{R}_8$	$S^1_9$	—
M2	—	—	$S^1_3$	—	$S^1_6$	—	$\mathbb{R}^{\geq 0}_8$	—	—

Reduce on  $S^1_3$ :

Spacetime	$\mathbb{R}_0$	$\mathbb{R}^2_{12}$	$\mathbb{R}^2_{45}$	$S^1_6$	$\mathbb{R}_7$	$\mathbb{R}_8$	$S^1_9$	$S^1_{10}$
$N$ D4	$\mathbb{R}_0$	$\mathbb{R}^2_{12}$	—	$S^1_6$	—	—	—	$S^1_{10}$
$n$ D4	$\mathbb{R}_0$	$\mathbb{R}^2_{12}$	—	—	—	$\mathbb{R}_8$	$S^1_9$	
F1	—	—	—	$S^1_6$	—	$\mathbb{R}^{\geq 0}_8$	—	—



Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$S_9^1$	$S_{10}^1$
$N$ D4	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	$S_6^1$	—	—	—	$S_{10}^1$
$n$ D4	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	—	—	$\mathbb{R}_8$	$S_9^1$	
F1	—	—	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	—	—

Apply T-duality  $S_9^1 \rightarrow \check{S}_9^1$ :

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\check{S}_9^1$	$S_{10}^1$
$N$ D5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	$S_6^1$	—	—	$\check{S}_9^1$	$S_{10}^1$
$n$ D3	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	—	—	$\mathbb{R}_8$	—	
F1	—	—	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	—	—

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\check{S}_9^1$	$S_{10}^1$
$N$ D5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	$S_6^1$	—	—	$\check{S}_9^1$	$S_{10}^1$
$n$ D3	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	—	—	$\mathbb{R}_8$	—	
F1	—	—	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	—	—

D5: 6d  $\mathcal{N} = (1, 1)$  SYM on  $\mathbb{R}_0 \times \mathbb{R}_{12}^2 \times S_6^1 \times \check{S}_9^1 \times S_{10}^1$

D3: codim-3 operator on  $\mathbb{R}_0 \times \mathbb{R}_{12}^2$

F1: Wilson line on  $S_6^1$

**$\Omega$ -deformation** on  $\mathbb{R}_{12}^2$  from nontrivial background, due to the initial twisted product in 12345 directions [Hellerman–Orland–Reffert].

$\Omega$ -deformed 6d  $\mathcal{N} = (1, 1)$  SYM on  $\mathbb{R}_0 \times \mathbb{R}_{12}^2 \times S_6^1 \times \check{S}_9^1 \times S_{10}^1$


$\rightsquigarrow$  Costello's **4d Chern-Simons** on  $\mathbb{R}_0 \times S_6^1 \times \check{S}_9^1 \times S_{10}^1$  [Costello-Y]

Codim-3 operators on  $\mathbb{R}_0 \times \mathbb{R}_{12}^2$

$\rightsquigarrow$  line operators on  $\mathbb{R}_0$


Wilson line on  $S_6^1$

$\rightsquigarrow$  Wilson line on  $S_6^1$




$\mathbb{R}_0 \times S_6^1$

$\times$




$\check{S}_9^1 \times S_{10}^1$



$$\mathbb{R}_0 \times S^1_6$$

×





$$\check{S}^1_9 \times S^1_{10}$$

Topological on  $\mathbb{R}_0 \times S^1_6$ , holomorphic on  $\check{S}^1_9 \times S^1_{10}$

2d TQFT + line defects  $\implies$  lattice model

TQFT + extra dimensions  $\implies$  integrability [Costello]





Wilson line gives transfer matrix of elliptic QIS with

$$\tau = iR_{10}/\check{R}_9 .$$

Now, decompactify  $S^1_9 \rightarrow \mathbb{R}_9$ . Take  $R_9 \rightarrow \infty$ , or  $\check{R}_9 \rightarrow 0$ .

This is the **trigonometric limit**  $\tau \rightarrow i\infty$ .

Dependence on the position on the torus (spectral parameter) is gone in this limit.

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\check{S}_9^1$	$S_{10}^1$
$N$ D5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	$S_6^1$	—	—	$\check{S}_9^1$	$S_{10}^1$
$n$ D3	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	—	—	$\mathbb{R}_8$	—	—
F1	—	—	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	—	—

For **Nekrasov–Shatashvili**, apply S-duality:

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\check{S}_9^1$	$S_{10}^1$
$N$ NS5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	$S_6^1$	—	—	$\check{S}_9^1$	$S_{10}^1$
$n$ D3	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	—	—	$\mathbb{R}_8$	—	—
D1	—	—	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	—	—


Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\check{S}_9^1$	$S_{10}^1$
$N$ NS5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	$S_6^1$	—	—	$\check{S}_9^1$	$S_{10}^1$
$n$ D3	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	—	—	$\mathbb{R}_8$	—	—
D1	—	—	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	—	—

Then T-duality on  $S_6^1$ :

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$\mathbb{R}_{45}^2$	$\check{S}_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\check{S}_9^1$	$S_{10}^1$
$N$ NS5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	$\check{S}_6^1$	—	—	$\check{S}_9^1$	$S_{10}^1$
$n$ D4	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	$\check{S}_6^1$	—	$\mathbb{R}_8$	—	—
D0	—	—	—	—	—	$\mathbb{R}_8^{\geq 0}$	—	—

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$\mathbb{R}_{45}^2$	$\check{S}_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\check{S}_9^1$	$S_{10}^1$
$N$ NS5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	$\check{S}_6^1$	—	—	$\check{S}_9^1$	$S_{10}^1$
$n$ D4	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	—	$\check{S}_6^1$	—	$\mathbb{R}_8$	—	—
D0	—	—	—	—	—	$\mathbb{R}_8^{\geq 0}$	—	—

D4–NS5: 4d  $\mathcal{N} = 2$  theory for  $(N + 1)$ -node linear quiver



placed on  $\mathbb{R}_0 \times \mathbb{R}_{12}^2 \times \check{S}_6^1$ .

$\Omega$ -deformation quantizes DW system (trigonometric Gaudin)  
 $\implies$  noncompact XXX spin chain

D0 is a local operator, acting as a transfer matrix.

Actuality, 9 & 10 directions are compact, so it's a 6d lift. We get the elliptic version of the integrable system.



## Go back to M-theory

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$S_9^1$	$S_{10}^1$
$N$ M5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	$S_6^1$	—	—	—	$S_{10}^1$
$n$ M5'	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	—	—	$\mathbb{R}_8$	$S_9^1$	—
M2	—	—	$S_3^1$	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	—	—

Reduce on  $S_{10}^1$ :

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\check{S}_9^1$
$N$ D4	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	$S_6^1$	—	—	—
$n$ NS5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	—	—	$\mathbb{R}_8$	$\check{S}_9^1$
D2	—	—	$S_3^1$	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	—

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\check{S}_9^1$
$N$ D4	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	$S_6^1$	—	—	—
$n$ NS5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	—	—	$\mathbb{R}_8$	$\check{S}_9^1$
D2	—	—	$S_3^1$	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	—

Apply T-duality on  $S_9^1$ :

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\check{S}_9^1$
$N$ D5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	$S_6^1$	—	—	$\check{S}_9^1$
$n$ NS5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	—	—	$\mathbb{R}_8$	$\check{S}_9^1$
D3	—	—	$S_3^1$	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	$\check{S}_9^1$

Spacetime	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	$\mathbb{R}_{45}^2$	$S_6^1$	$\mathbb{R}_7$	$\mathbb{R}_8$	$\check{S}_9^1$
$N$ D5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	$S_6^1$	—	—	$\check{S}_9^1$
$n$ NS5	$\mathbb{R}_0$	$\mathbb{R}_{12}^2$	$S_3^1$	—	—	—	$\mathbb{R}_8$	$\check{S}_9^1$
D3	—	—	$S_3^1$	—	$S_6^1$	—	$\mathbb{R}_8^{\geq 0}$	$\check{S}_9^1$

D5–NS5: 5d circular quiver theory on  $\mathbb{R}_0 \times \mathbb{R}_{12}^2 \times_{\epsilon} S_3^1 \times \check{S}_9^1$

D3: surface defect on  $S_3^1 \times \check{S}_9^1$

We can add more NS5s, preserving 4d  $\mathcal{N} = 1$  SUSY on  $\mathbb{R}_{12}^2 \times_{\epsilon} S_3^1 \times \check{S}_9^1$ . This leads to the brane tiling story [Maruyoshi–Y].

## Summary

- We considered a class of Wilson–’t Hooft lines in 4d  $\mathcal{N} = 2$  circular quiver theories.
- We found that they can be identified with transfer matrices of trigonometric QIS.
- By embedding into string theory, this correspondence can be related to other known correspondences via dualities.

## Further directions

- Surface defects in 5d circular quiver theory correspond to transfer matrices of elliptic QIS.
- Variations of the present setup
- Circular quiver theories deconstruct 6d  $\mathcal{N} = (2, 0)$  SCFT. Integrability is behind surface operators in 6d theory.