

TikhonovFenichelReductions.jl: A systematic approach to geometric singular perturbation theory

Johannes Apelt¹

¹ Institute of Mathematics and Computer Science, University of Greifswald, Germany

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Natural systems often evolve on different characteristic time scales, with process rates differing substantially by orders of magnitude ([Hek, 2010](#)). To study such systems, singular perturbation theory provides a mathematical toolbox that translates this feature to ODE models and enables dimensionality reduction by focusing on a single characteristic time scale, (typically the slow one governing the long-term behaviour) ([Wechselberger, 2020](#)).

This separation of time scales is reflected by the presence of a small parameter $\varepsilon > 0$, which can be interpreted as the ratio between the two time scales. Roughly speaking, the dynamics in slow time $\tau = \varepsilon t$ can be approximated by replacing the fast part of the system with its steady state, so that the fast dynamics effectively appear instantaneous on the slow time scale.

More precisely, we obtain the *reduced system* (or *reduction* in short) as the limit

$$\begin{aligned} \dot{u} &= g(u, v) \\ \varepsilon \dot{v} &= h(u, v) \end{aligned} \xrightarrow{\varepsilon \rightarrow 0} \begin{aligned} \dot{u} &= g(u, v), & u &\in \mathbb{R}^s \\ 0 &= h(u, v), & v &\in \mathbb{R}^r \end{aligned} \quad (1)$$

with $\dot{} = d/d\tau$ as stated in Tikhonov's theorem ([1952](#); Theorem 1.1 in [Verhulst, 2007](#)). The reduced system is defined on the so-called slow manifold $M_0 = \{(u, v) \mid h(u, v) = 0\}$.

For autonomous systems, the results of Fenichel ([1979](#)) allow a clear geometric interpretation — hence the name geometric singular perturbation theory (GSPT) — and guarantee that for sufficiently small $\varepsilon > 0$, the reduction captures the behaviour of the full system.

In practice, GSPT can mitigate the realism-complexity trade-off, as modellers can work with the reduction instead of the full system, which is often more tractable. Because the reduction remains embedded in the original system through the time scale separation, the original interpretability is retained. This allows one to follow the modelling paradigm: Start with a complex but realistic model, reduce later.

The algebraic approach recently developed by Goeke & Walcher (and colleagues) allows one to work with GSPT systematically and is coordinate-free, i.e. we do not rely on the standard form (1) with a time scale separation of components.

Instead, we consider systems of the form

$$x' = f^{(0)}(x) + \varepsilon f^{(1)}(x) + \mathcal{O}(\varepsilon^2), \quad x \in \mathbb{R}^n, \quad (2)$$

where $f^{(0)}$ contains the fast processes and $f^{(1)}$ contains the slow ones. Note that system (1) can also be written in fast time t as

$$\begin{aligned} u' &= \varepsilon g(u, v) \\ v' &= h(u, v) \end{aligned} \quad (3)$$

where $' = d/dt$. Thus, we can translate (2) into (3) by setting $x = (u, v)^T$, $f^{(0)} = (0, h)^T$ and $f^{(1)} = (g, 0)^T$. However, the reverse direction requires a suitable coordinate transformation, which is not always feasible. Fortunately, this transformation is not explicitly needed to obtain the reduced system, which can be computed using (4).

The main benefit of this framework is however that it allows to find all critical parameters admitting a reduction for polynomial or rational ODE systems of the form (2). Thus, modellers do not have to rely on prior knowledge to find suitable time scale separations as they can systematically consider all of them. This can be achieved by evaluating necessary conditions for the existence of a reduction (Goeke, 2013; Goeke et al., 2015; Goeke & Walcher, 2013, 2014). TikhonovFenichelReductions.jl is a Julia (Bezanson et al., 2017) package implementing this approach for polynomial ODE systems. Apelt & Liebscher (2025) provide a showcasing example and more detailed explanations.

Statement of need

The ad-hoc approach to singular perturbation theory requires prior knowledge of a suitable time scale separation and substantial mathematical effort to compute the reduction. In contrast, the algebraic framework developed by Goeke & Walcher systematically yields *all* possible reductions for a given polynomial ODE system, enabling modellers to consider several scenarios defined by different time scale separations (Goeke et al., 2015). TikhonovFenichelReductions.jl makes the required computations easily accessible, even for non-expert users.

To the author's knowledge, no publicly available implementation of this theory currently exists. However, there is an implementation of a related approach for computing invariant manifolds by Roberts (n.d., 1997).

Software design

TikhonovFenichelReductions.jl is implemented in Julia due to its flexibility, the use of multiple dispatch and the availability of the feature-rich computer algebra system Oscar.jl (Decker et al., 2025; The OSCAR Team, 2025).

Core features are the search for critical parameters admitting a reduction, so-called *Tikhonov-Fenichel Parameter Values (TFPVs)*, and the computation of the corresponding reduced systems. Crucially, this requires various computations with multiple symbolic representations (i.e. different polynomial rings, rational function fields and matrix spaces) and therefore parsing of data between the corresponding types in Oscar.jl, which TikhonovFenichelReductions.jl performs hidden away from the user. Thus, the user mostly works in an object-oriented manner with the types and methods provided.

The package is essentially designed around two types: `ReductionProblem`, which constructs all symbolic data types needed for the search of TFPVs, and `Reduction`, which holds all relevant information for the reduced system and the steps required to compute it. The latter also contains the reduced system and other information parsed to the appropriate types from Oscar.jl, that can be further used, e.g. for a symbolic analysis.

Detailed explanations and examples are provided by Apelt & Liebscher (2025) and in the [documentation](#). The core features are summarized in the following.

Finding TFPVs

The package provides a method to obtain all possible TFPVs implicitly by computing a Gröbner Basis and one to find *slow-fast separations of rates*, i.e. TFPVs with some parameters set to zero. Although the former method is an exhaustive search, the latter is usually better suited in practice as it yields the TFPVs one is typically interested in explicitly, is more efficient, and

77 directly outputs the corresponding slow manifolds (implicitly as affine varieties in phase space).
 78 In both cases, one utilizes the correspondence between algebraic and geometric properties of
 79 the slow manifold, which renders this an algebraic approach to GSPT.

80 Computing reductions

81 Computing a reduction for a slow-fast separation of rates as in Theorem 1 in Goeke & Walcher
 82 (2014) requires essentially two steps. First, we need to provide a parametric representation of
 83 the slow manifold, which is given as an irreducible component of the affine variety $\mathcal{V}(f^{(0)})$.
 84 Then, we need to find a product decomposition $f^{(0)} = P\psi$, where P is a $n \times r$ matrix of
 85 rational functions and ψ is a vector of polynomials locally satisfying $\mathcal{V}(\psi) = \mathcal{V}(f^{(0)})$ and
 86 $\text{rank } P = \text{rank } D\psi = r$.

87 With this, the reduced system in the sense of Tikhonov is given by

$$x' = [1_n - P(x)(D\psi(x)P(x))^{-1}D\psi(x)] f^{(1)}(x). \quad (4)$$

88 For convenience, the packages provides multiple heuristics, which automate the steps required
 89 to compute a reduction and allows bulk computation of multiple reductions at once.

90 Integration with the Julia ecosystem

91 The input system can be given as a reaction network defined with `Catalyst.jl` (Loman et al.,
 92 2023). Because the reduced systems are represented using types from `Oscar.jl`, the latter's
 93 functions can be used to aid the symbolic analysis. Julia's support for metaprogramming
 94 allows to perform further tasks such as a numerical analysis without having to copy or parse
 95 code (see e.g. `TFRSimulations.jl`). For convenience, there are several methods for displaying
 96 the output, including printing as $\text{\LaTeX}$ source code via `Latexify.jl`.

97 Research impact statement

98 Time scale separations are widely used in various areas of mathematical modelling (Wechsel-
 99 berger, 2020). However, as far as the author is aware, the systematic approach due to Goeke &
 100 Walcher seems to be scarcely adopted, even though it comes with many advantages compared
 101 to the ad-hoc approach. This package aims to make the theory more accessible and convenient
 102 to use.

103 In the field of mathematical ecology (which the author is most familiar with), the theory was
 104 successfully used by Kruff et al. (2019) and more recently by Apelt & Liebscher (2025). For
 105 the model introduced in the former, the application was relatively straightforward, but more
 106 complex models as in the latter require some more work. In particular, the method for finding
 107 TFPVs that relies on the computation of a Gröbner Basis may fail for complex systems. In
 108 this case, `TikhonovFenichelReductions.jl` may enable and simplify the search for the most
 109 common TFPVs. Given the ubiquity of time scale separation techniques in this field alone
 110 (see e.g. Abbott et al., 2020; Poggiale & Auger, 2004; Revilla, 2015), the package can be a
 111 potentially useful tool for modellers.

112 AI usage disclosure

113 No generative AI tools were used in the development of this software, the writing of this
 114 manuscript, or the preparation of supporting materials.

Acknowledgements

This work was supported by a scholarship awarded by the University of Greifswald according to the “Landesgraduiertenförderungsgesetz (LGFG) MV”. I like to thank Volkmar Liebscher for his supervision and support, Sebastian Walcher and Alexandra Goeke for developing the Tikhonov-Fenichel reduction theory, Leonard Schmitz for discussing aspects of the computational algebra, and the OSCAR team for their great work and helpful replies to my questions.

References

- Abbott, K. C., Ji, F., Stieha, C. R., & Moore, C. M. (2020). Fast and slow advances toward a deeper integration of theory and empiricism. *Theoretical Ecology*, 13(1), 7–15. <https://doi.org/10.1007/s12080-019-00441-x>
- Apelt, J., & Liebscher, V. (2025). Tikhonov-Fenichel reductions and their application to a novel modelling approach for mutualism. *Theoretical Population Biology*, 16–35. <https://doi.org/10.1016/j.tpb.2025.08.004>
- Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. B. (2017). Julia: A Fresh Approach to Numerical Computing. *SIAM Review*, 59(1), 65–98. <https://doi.org/10.1137/141000671>
- Decker, W., Eder, C., Fieker, C., Horn, M., & Joswig, M. (Eds.). (2025). *The Computer Algebra System OSCAR: Algorithms and Examples*. Springer Nature Switzerland. <https://doi.org/10.1007/978-3-031-62127-7>
- Fenichel, N. (1979). Geometric singular perturbation theory for ordinary differential equations. *Journal of Differential Equations*, 31(1), 53–98. [https://doi.org/10.1016/0022-0396\(79\)90152-9](https://doi.org/10.1016/0022-0396(79)90152-9)
- Goeke, A. (2013). *Reduktion und asymptotische Reduktion von Reaktionsgleichungen* [RWTH Aachen]. <https://publications.rwth-aachen.de/record/229008>
- Goeke, A., & Walcher, S. (2013). Quasi-Steady State: Searching for and Utilizing Small Parameters. In A. Johann, H.-P. Kruse, F. Rupp, & S. Schmitz (Eds.), *Recent Trends in Dynamical Systems* (Vol. 35, pp. 153–178). Springer Basel. https://doi.org/10.1007/978-3-0348-0451-6_8
- Goeke, A., & Walcher, S. (2014). A constructive approach to quasi-steady state reductions. *Journal of Mathematical Chemistry*, 52(10), 2596–2626. <https://doi.org/10.1007/s10910-014-0402-5>
- Goeke, A., Walcher, S., & Zerz, E. (2015). Determining “small parameters” for quasi-steady state. *Journal of Differential Equations*, 259(3), 1149–1180. <https://doi.org/10.1016/j.jde.2015.02.038>
- Hek, G. (2010). Geometric singular perturbation theory in biological practice. *Journal of Mathematical Biology*, 60(3), 347–386. <https://doi.org/10.1007/s00285-009-0266-7>
- Kruff, N., Lax, C., Liebscher, V., & Walcher, S. (2019). The Rosenzweig–MacArthur system via reduction of an individual based model. *Journal of Mathematical Biology*, 78(1–2), 413–439. <https://doi.org/10.1007/s00285-018-1278-y>
- Loman, T. E., Ma, Y., Ilin, V., Gowda, S., Korsbo, N., Yewale, N., Rackauckas, C., & Isaacson, S. A. (2023). Catalyst: Fast and flexible modeling of reaction networks. *PLOS Computational Biology*, 19(10), e1011530. <https://doi.org/10.1371/journal.pcbi.1011530>
- Poggiale, J.-C., & Auger, P. (2004). Impact of spatial heterogeneity on a predator–prey system dynamics. *Comptes Rendus. Biologies*, 327(11), 1058–1063. <https://doi.org/10.1016/j.crv.2004.06.006>

- 159 Revilla, T. A. (2015). Numerical responses in resource-based mutualisms: A time scale approach.
160 *Journal of Theoretical Biology*, 378, 39–46. <https://doi.org/10.1016/j.jtbi.2015.04.012>
- 161 Roberts, A. J. (n.d.). *Construct invariant manifolds*. <https://tuck.adelaide.edu.au/gencm.php>
- 162 Roberts, A. J. (1997). Low-dimensional modelling of dynamics via computer algebra. *Computer*
163 *Physics Communications*, 100(3), 215–230. [https://doi.org/10.1016/s0010-4655\(96\)](https://doi.org/10.1016/s0010-4655(96)00162-2)
164 [00162-2](https://doi.org/10.1016/s0010-4655(96)00162-2)
- 165 The OSCAR Team. (2025). *OSCAR – Open Source Computer Algebra Research system*.
166 <https://www.oscar-system.org>
- 167 Tikhonov, A. N. (1952). Systems of differential equations containing small parameters in the
168 derivatives. *Matematicheskii sbornik*, 73(3), 575–586.
- 169 Verhulst, F. (2007). Singular perturbation methods for slow–fast dynamics. *Nonlinear*
170 *Dynamics*, 50(4), 747–753. <https://doi.org/10.1007/s11071-007-9236-z>
- 171 Wechselberger, M. (2020). *Geometric Singular Perturbation Theory Beyond the Standard*
172 *Form*. Springer International Publishing. <https://doi.org/10.1007/978-3-030-36399-4>

DRAFT