

Model fitting

Joachim Vandekerckhove

Winter 2025

Item Response Models

- The two-parameter logistic (2PL) model describes the probability of a correct response:

$$P(y|\theta, a, b_i) = \frac{1}{1 + e^{-a(\theta - b_i)}} \quad (1)$$

Item Response Models

- The two-parameter logistic (2PL) model describes the probability of a correct response:

$$P(y|\theta, a, b_i) = \frac{1}{1 + e^{-a(\theta - b_i)}} \quad (1)$$

- Parameters:

Item Response Models

- The two-parameter logistic (2PL) model describes the probability of a correct response:

$$P(y|\theta, a, b_i) = \frac{1}{1 + e^{-a(\theta - b_i)}} \quad (1)$$

- Parameters:
 - a : Discrimination (slope)

Item Response Models

- The two-parameter logistic (2PL) model describes the probability of a correct response:

$$P(y|\theta, a, b_i) = \frac{1}{1 + e^{-a(\theta - b_i)}} \quad (1)$$

- Parameters:
 - a : Discrimination (slope)
 - b_i : Difficulty (location) for item or condition i

Item Response Models

- The two-parameter logistic (2PL) model describes the probability of a correct response:

$$P(y|\theta, a, b_i) = \frac{1}{1 + e^{-a(\theta - b_i)}} \quad (1)$$

- Parameters:
 - a : Discrimination (slope)
 - b_i : Difficulty (location) for item or condition i
 - θ : Subject ability

Item Response Models


- The two-parameter logistic (2PL) model describes the probability of a correct response:

$$P(y|\theta, a, b_i) = \frac{1}{1 + e^{-a(\theta - b_i)}} \quad (1)$$

- Parameters:
 - a : Discrimination (slope)
 - b_i : Difficulty (location) for item or condition i
 - θ : Subject ability
- The three-parameter logistic (3PL) adds a guessing parameter c :

$$P(y|\theta, a, b_i, c) = c + (1 - c) \frac{1}{1 + e^{-a(\theta - b_i)}} \quad (2)$$

Visualizing the 2PL Model



Ordinary Least Squares

- Ordinary Least Squares (OLS) is a common estimation method:

$$\hat{\beta} = \arg \min_{\beta} \sum_{i=1}^N (y_i - f(x_i, \beta))^2 \quad (3)$$

Ordinary Least Squares

- Ordinary Least Squares (OLS) is a common estimation method:

$$\hat{\beta} = \arg \min_{\beta} \sum_{i=1}^N (y_i - f(x_i, \beta))^2 \quad (3)$$

- Properties:

Ordinary Least Squares

- Ordinary Least Squares (OLS) is a common estimation method:

$$\hat{\beta} = \arg \min_{\beta} \sum_{i=1}^N (y_i - f(x_i, \beta))^2 \quad (3)$$

- Properties:
 - Minimizes squared differences between observed and predicted values (“loss”)

Ordinary Least Squares

- Ordinary Least Squares (OLS) is a common estimation method:

$$\hat{\beta} = \arg \min_{\beta} \sum_{i=1}^N (y_i - f(x_i, \beta))^2 \quad (3)$$

- Properties:
 - Minimizes squared differences between observed and predicted values (“loss”)
 - Optimal when errors are normally distributed

Ordinary Least Squares

- Ordinary Least Squares (OLS) is a common estimation method:

$$\hat{\beta} = \arg \min_{\beta} \sum_{i=1}^N (y_i - f(x_i, \beta))^2 \quad (3)$$

- Properties:
 - Minimizes squared differences between observed and predicted values (“loss”)
 - Optimal when errors are normally distributed
 - Has closed-form solution for linear models

Ordinary Least Squares

- Ordinary Least Squares (OLS) is a common estimation method:

$$\hat{\beta} = \arg \min_{\beta} \sum_{i=1}^N (y_i - f(x_i, \beta))^2 \quad (3)$$

- Properties:
 - Minimizes squared differences between observed and predicted values (“loss”)
 - Optimal when errors are normally distributed
 - Has closed-form solution for linear models
 - Simple to implement and interpret

Squared Error Estimation

- Implementing OLS for the 2PL model, parameters are estimated by minimizing squared error:

$$\text{SSE}(\theta, a, \mathbf{b}, \mathbf{y}) = \sum_{i=1}^N (y_i - P(y_i | \theta, a, b_i))^2 \quad (4)$$

Squared Error Estimation

- Implementing OLS for the 2PL model, parameters are estimated by minimizing squared error:

$$\text{SSE}(\theta, a, \mathbf{b}, \mathbf{y}) = \sum_{i=1}^N (y_i - P(y_i | \theta, a, b_i))^2 \quad (4)$$

- Where:

Squared Error Estimation

- Implementing OLS for the 2PL model, parameters are estimated by minimizing squared error:

$$\text{SSE}(\theta, a, \mathbf{b}, \mathbf{y}) = \sum_{i=1}^N (y_i - P(y_i | \theta, a, b_i))^2 \quad (4)$$

- Where:
 - y_i is the participant's observed response (0 or 1) to the i^{th} item of N items of varying difficulty b_i

Squared Error Estimation

- Implementing OLS for the 2PL model, parameters are estimated by minimizing squared error:

$$\text{SSE}(\theta, a, \mathbf{b}, \mathbf{y}) = \sum_{i=1}^N (y_i - P(y_i|\theta, a, b_i))^2 \quad (4)$$

- Where:
 - y_i is the participant's observed response (0 or 1) to the i^{th} item of N items of varying difficulty b_i
 - $P(y_i|\theta, a, b_i)$ is the model prediction of y_i

Squared Error Estimation

- Implementing OLS for the 2PL model, parameters are estimated by minimizing squared error:

$$\text{SSE}(\theta, a, \mathbf{b}, \mathbf{y}) = \sum_{i=1}^N (y_i - P(y_i|\theta, a, b_i))^2 \quad (4)$$

- Where:
 - y_i is the participant's observed response (0 or 1) to the i^{th} item of N items of varying difficulty b_i
 - $P(y_i|\theta, a, b_i)$ is the model prediction of y_i
- This approach:

Squared Error Estimation

- Implementing OLS for the 2PL model, parameters are estimated by minimizing squared error:

$$\text{SSE}(\theta, a, \mathbf{b}, \mathbf{y}) = \sum_{i=1}^N (y_i - P(y_i|\theta, a, b_i))^2 \quad (4)$$

- Where:
 - y_i is the participant's observed response (0 or 1) to the i^{th} item of N items of varying difficulty b_i
 - $P(y_i|\theta, a, b_i)$ is the model prediction of y_i
- This approach:
 - Is intuitive and widely used in simple cases


Squared Error Estimation

- Implementing OLS for the 2PL model, parameters are estimated by minimizing squared error:

$$\text{SSE}(\theta, a, \mathbf{b}, \mathbf{y}) = \sum_{i=1}^N (y_i - P(y_i|\theta, a, b_i))^2 \quad (4)$$


- Where:
 - y_i is the participant's observed response (0 or 1) to the i^{th} item of N items of varying difficulty b_i
 - $P(y_i|\theta, a, b_i)$ is the model prediction of y_i
- This approach:
 - Is intuitive and widely used in simple cases
 - Assumes normally distributed errors

Normality Assumption




- May not be appropriate for non-normal errors

Normality Assumption



- May not be appropriate for non-normal errors
- Can give biased estimates for binary outcomes

Normality Assumption



- May not be appropriate for non-normal errors
- Can give biased estimates for binary outcomes
- Fine for a first pass, but not optimal for many applications

Maximum Likelihood Estimation

- For binary outcomes, likelihood is more appropriate:

$$L(\theta, a, \mathbf{b}, \mathbf{y}) = \prod_{i=1}^N P(y_i|\theta, a, b_i)^{y_i} (1 - P(y_i|\theta, a, b_i))^{1-y_i} \quad (5)$$

Maximum Likelihood Estimation

- For binary outcomes, likelihood is more appropriate:

$$L(\theta, a, \mathbf{b}, \mathbf{y}) = \prod_{i=1}^N P(y_i|\theta, a, b_i)^{y_i} (1 - P(y_i|\theta, a, b_i))^{1-y_i} \quad (5)$$

- Taking the log gives the log-likelihood function:

$$\ell(\theta, a, \mathbf{b}, \mathbf{y}) = \sum_{i=1}^N [y_i \log P(y_i|\theta, a, b_i) + (1 - y_i) \log(1 - P(y_i|\theta, a, b_i))] \quad (6)$$

Maximum Likelihood Estimation

- For binary outcomes, likelihood is more appropriate:

$$L(\theta, a, \mathbf{b}, \mathbf{y}) = \prod_{i=1}^N P(y_i|\theta, a, b_i)^{y_i} (1 - P(y_i|\theta, a, b_i))^{1-y_i} \quad (5)$$

- Taking the log gives the log-likelihood function:

$$\ell(\theta, a, \mathbf{b}, \mathbf{y}) = \sum_{i=1}^N [y_i \log P(y_i|\theta, a, b_i) + (1 - y_i) \log(1 - P(y_i|\theta, a, b_i))] \quad (6)$$

- Using the likelihood function instead of squared error appropriately accounts for the fact that **some prediction errors are more severe than others**

Regularization

- Regularization is a technique used to prevent overfitting in models

Regularization

- Regularization is a technique used to prevent overfitting in models
- It adds a penalty term to the loss function to discourage large parameter values

Regularization

- Regularization is a technique used to prevent overfitting in models
- It adds a penalty term to the loss function to discourage large parameter values
- This helps in cases where there is a lot of data and many parameters

Regularization

- Regularization is a technique used to prevent overfitting in models
- It adds a penalty term to the loss function to discourage large parameter values
- This helps in cases where there is a lot of data and many parameters
- Let's consider a 2PL model with predictors on item difficulty:

$$P(y|\theta, a, x, \beta) = \frac{1}{1 + e^{-a(\theta - f(x, \beta))}} \quad (7)$$

with $f(x, \beta) = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_k x_k$

Regularization

- Regularization is a technique used to prevent overfitting in models
- It adds a penalty term to the loss function to discourage large parameter values
- This helps in cases where there is a lot of data and many parameters
- Let's consider a 2PL model with predictors on item difficulty:

$$P(y|\theta, a, x, \beta) = \frac{1}{1 + e^{-a(\theta - f(x, \beta))}} \quad (7)$$

with $f(x, \beta) = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_k x_k$

- Here, β is a vector of parameters for the predictors that replaces the item difficulty parameters **b**

Regularization

- Regularization is a technique used to prevent overfitting in models
- It adds a penalty term to the loss function to discourage large parameter values
- This helps in cases where there is a lot of data and many parameters
- Let's consider a 2PL model with predictors on item difficulty:

$$P(y|\theta, a, x, \beta) = \frac{1}{1 + e^{-a(\theta - f(x, \beta))}} \quad (7)$$

with $f(x, \beta) = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_k x_k$

- Here, β is a vector of parameters for the predictors that replaces the item difficulty parameters \mathbf{b}
- Fewer predictors is better for model parsimony and interpretability

Regularization

- Regularization is a technique used to prevent overfitting in models
- It adds a penalty term to the loss function to discourage large parameter values
- This helps in cases where there is a lot of data and many parameters
- Let's consider a 2PL model with predictors on item difficulty:

$$P(y|\theta, a, x, \beta) = \frac{1}{1 + e^{-a(\theta - f(x, \beta))}} \quad (7)$$

with $f(x, \beta) = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_k x_k$

- Here, β is a vector of parameters for the predictors that replaces the item difficulty parameters \mathbf{b}
- Fewer predictors is better for model parsimony and interpretability
- Certainly, we need there to be fewer predictors than conditions, so $k < N$

LASSO and Ridge Regularization

$$\ell(\theta, a, x, \beta) = \sum_{i=1}^N [y_i \log P(y|\theta, a, x, \beta) + (1 - y_i) \log(1 - P(y|\theta, a, x, \beta))]$$

- LASSO (Least Absolute Shrinkage and Selection Operator) equation:

$$\ell_{\text{LASSO}}(\theta, a, x, \beta) = \ell(\theta, a, x, \beta) - \lambda \sum_{j=1}^k |\beta_j| \quad (8)$$

LASSO and Ridge Regularization

$$\ell(\theta, a, x, \beta) = \sum_{i=1}^N [y_i \log P(y|\theta, a, x, \beta) + (1 - y_i) \log(1 - P(y|\theta, a, x, \beta))]$$

- LASSO (Least Absolute Shrinkage and Selection Operator) equation:

$$\ell_{\text{LASSO}}(\theta, a, x, \beta) = \ell(\theta, a, x, \beta) - \lambda \sum_{j=1}^k |\beta_j| \quad (8)$$

- Ridge regularization equation:

$$\ell_{\text{Ridge}}(\theta, a, x, \beta) = \ell(\theta, a, x, \beta) - \lambda \sum_{j=1}^k \beta_j^2 \quad (9)$$

Interpretation of LASSO and Ridge Regularization

- LASSO:

Interpretation of LASSO and Ridge Regularization

- LASSO:
 - Shrinks the coefficients towards zero

Interpretation of LASSO and Ridge Regularization

- LASSO:
 - Shrinks the coefficients towards zero
 - Can set some coefficients to zero, performing “feature selection”

Interpretation of LASSO and Ridge Regularization

- LASSO:
 - Shrinks the coefficients towards zero
 - Can set some coefficients to zero, performing “feature selection”
 - Is equivalent to a Bayesian model with a Laplace prior on the coefficients

Interpretation of LASSO and Ridge Regularization

- LASSO:
 - Shrinks the coefficients towards zero
 - Can set some coefficients to zero, performing “feature selection”
 - Is equivalent to a Bayesian model with a Laplace prior on the coefficients
- Ridge:

Interpretation of LASSO and Ridge Regularization

- LASSO:
 - Shrinks the coefficients towards zero
 - Can set some coefficients to zero, performing “feature selection”
 - Is equivalent to a Bayesian model with a Laplace prior on the coefficients
- Ridge:
 - Shrinks the coefficients towards zero

Interpretation of LASSO and Ridge Regularization

- LASSO:
 - Shrinks the coefficients towards zero
 - Can set some coefficients to zero, performing “feature selection”
 - Is equivalent to a Bayesian model with a Laplace prior on the coefficients
- Ridge:
 - Shrinks the coefficients towards zero
 - Does not set any coefficients to zero

Interpretation of LASSO and Ridge Regularization

- LASSO:
 - Shrinks the coefficients towards zero
 - Can set some coefficients to zero, performing “feature selection”
 - Is equivalent to a Bayesian model with a Laplace prior on the coefficients
- Ridge:
 - Shrinks the coefficients towards zero
 - Does not set any coefficients to zero
 - Is equivalent to a Bayesian model with a Gaussian prior on the coefficients

Interpretation of LASSO and Ridge Regularization

- LASSO:
 - Shrinks the coefficients towards zero
 - Can set some coefficients to zero, performing “feature selection”
 - Is equivalent to a Bayesian model with a Laplace prior on the coefficients
- Ridge:
 - Shrinks the coefficients towards zero
 - Does not set any coefficients to zero
 - Is equivalent to a Bayesian model with a Gaussian prior on the coefficients
- Regularization is a form of Bayesian inference

Interpretation of LASSO and Ridge Regularization

- LASSO:
 - Shrinks the coefficients towards zero
 - Can set some coefficients to zero, performing “feature selection”
 - Is equivalent to a Bayesian model with a Laplace prior on the coefficients
- Ridge:
 - Shrinks the coefficients towards zero
 - Does not set any coefficients to zero
 - Is equivalent to a Bayesian model with a Gaussian prior on the coefficients
- Regularization is a form of Bayesian inference
- Can be combined with likelihoods or more basic loss functions (like squared error)

- The 2PL model is a close relative to Signal Detection Theory

Conclusion

- The 2PL model is a close relative to Signal Detection Theory
- You can fit the 2PL model using maximum likelihood estimation

Conclusion

- The 2PL model is a close relative to Signal Detection Theory
- You can fit the 2PL model using maximum likelihood estimation
- You can regularize the model using LASSO and Ridge regularization

Conclusion

- The 2PL model is a close relative to Signal Detection Theory
- You can fit the 2PL model using maximum likelihood estimation
- You can regularize the model using LASSO and Ridge regularization
- This is very similar to Bayesian parameter estimation

Conclusion

- The 2PL model is a close relative to Signal Detection Theory
- You can fit the 2PL model using maximum likelihood estimation
- You can regularize the model using LASSO and Ridge regularization
- This is very similar to Bayesian parameter estimation
- Soon, we will create python classes for:

Conclusion

- The 2PL model is a close relative to Signal Detection Theory
- You can fit the 2PL model using maximum likelihood estimation
- You can regularize the model using LASSO and Ridge regularization
- This is very similar to Bayesian parameter estimation
- Soon, we will create python classes for:
 - the 2PL model

Conclusion

- The 2PL model is a close relative to Signal Detection Theory
- You can fit the 2PL model using maximum likelihood estimation
- You can regularize the model using LASSO and Ridge regularization
- This is very similar to Bayesian parameter estimation
- Soon, we will create python classes for:
 - the 2PL model
 - the regularized 2PL model

Conclusion

- The 2PL model is a close relative to Signal Detection Theory
- You can fit the 2PL model using maximum likelihood estimation
- You can regularize the model using LASSO and Ridge regularization
- This is very similar to Bayesian parameter estimation
- Soon, we will create python classes for:
 - the 2PL model
 - the regularized 2PL model
- These classes will allow us to fit the model to data and perform inference

Conclusion

- The 2PL model is a close relative to Signal Detection Theory
- You can fit the 2PL model using maximum likelihood estimation
- You can regularize the model using LASSO and Ridge regularization
- This is very similar to Bayesian parameter estimation
- Soon, we will create python classes for:
 - the 2PL model
 - the regularized 2PL model
- These classes will allow us to fit the model to data and perform inference
- I will add a twist that is likely to confuse an AI assistant

Model fitting

Joachim Vandekerckhove

Winter 2025