Continuous random variable

Probability density functio

Cumulative distribution functions

Common distributions

Continuous Random Variables

James Heald¹

¹Gatsby Computational Neuroscience Unit University College London

Gatsby Bridging Programme 2023



Objectives

Continuous

Probability density function

Cumulativo distribution functions

Common distributio

- Introduce the concept and formal definition of a continuous random variable *X* and a probability density function.
- Learn how to find the probability that a continuous random variable falls in some interval [a, b].
- Learn that if X is continuous, the probability that X takes on any specific value is 0.
- Introduce the concept and formal definition of a cumulative distribution function of a continuous random variable.
- Learn how to find the cumulative distribution function of a continuous random variable X from the probability density function of X.

Discrete vs. continuous random variables

Continuous random variables

Probability density function

Cumulativo distribution functions

Common distributior Unlike discrete random variables, which can take on a countable number of possible values (e.g. faces of a die or cards in a deck), continuous random variables can take on an uncountable number of possible values (e.g. all the real numbers in an interval).

Discrete vs. continuous random variables

Continuous random variables

Probability density function

Cumulativo distribution functions

Common distributio Unlike discrete random variables, which can take on a countable number of possible values (e.g. faces of a die or cards in a deck), continuous random variables can take on an uncountable number of possible values (e.g. all the real numbers in an interval).

Examples

- the voltage membrane potential of a cell
- the interspike interval of a neuron
- the force generated by a muscle
- the velocity of an eye movement

Continuous random variables

Continuous random variables

Probability density function

Cumulativo distribution functions

Common distribution

Definition

A random variable X is continuous if:

- possible values comprise either a single interval on the number line (i.e. for some a < b, any number x between a and b is a possible value) or a union of disjoint intervals, and
- P(X = c) = 0 for any number c that is a possible value of X.


Discrete probability distributions in the limit

Continuous

Probability density functions

Cumulative distribution functions

Common distributions Continuous random variables can be discretised into bins to form a discrete distribution that can be viewed as a probability histogram. As the bins become narrower, the histogram approaches a smooth curve.



Probabilities as integrals


Continuous

Probability density functions

Cumulative distributior functions

Common distribution: The probability that a continuous random variable X takes on a value in the interval [a, b] is given by the area under the probability density function f(x).

$$P(a \le X \le b) = \int_a^b f(x) dx$$



The probability density function

Continuous random variable

Probability density functions

Cumulative distributior functions

Common distributions

Definition

A random variable X is continuous if there exists a nonnegative function f(x) defined on the interval $(-\infty, \infty)$, such that for any interval [a, b] we have

$$P(a \le X \le b) = \int_a^b f(x) dx.$$

The probability density function

Continuous

Probability density functions

Cumulative distribution functions

Common distributions

Definition

A random variable X is continuous if there exists a nonnegative function f(x) defined on the interval $(-\infty, \infty)$, such that for any interval [a, b] we have

$$P(a \le X \le b) = \int_a^b f(x) dx.$$

A valid probability density function (PDF) f(x) has the following properties:

$$f(x) \ge 0 \text{ for all } x$$
 (1)

$$\int_{-\infty}^{\infty} f(x)dx = 1. \tag{2}$$

Density as probability per unit length

Continuous

Probability density functions

Cumulativ distributio functions

Common distribution

The probability of a small interval δ is approximately the density \times δ :

$$P(x \le X \le x + \delta) = \int_{x}^{x+\delta} f(t)dt$$
$$\approx f(x) \times \delta$$

Density as probability per unit length

Continuous random variable

Probability density functions

Cumulativ distributio functions

Common distributions The probability of a small interval δ is approximately the density \times δ :

$$P(x \le X \le x + \delta) = \int_{x}^{x+\delta} f(t)dt$$
$$\approx f(x) \times \delta$$

Thus density is probability per unit length (probability accumulation rate):

$$\frac{P(x \le X \le x + \delta)}{\delta} \approx f(x)$$

Each possible value has zero probability

Continuous random variable

Probability density functions

Cumulativ distributio functions

Common distribution

The probability that X takes on a particular value a is 0, as

$$P(X = a) = \int_{a}^{a} f(x)dx$$
$$= \lim_{\epsilon \to 0} \int_{a-\epsilon}^{a+\epsilon} f(x)dx$$
$$= 0.$$

Each possible value has zero probability

Continuous

Probability density functions

Cumulativ distributio functions

Common distribution The probability that X takes on a particular value a is 0, as

$$P(X = a) = \int_{a}^{a} f(x)dx$$
$$= \lim_{\epsilon \to 0} \int_{a-\epsilon}^{a+\epsilon} f(x)dx$$
$$= 0.$$

This implies that probabilities don't depend on interval end points:

$$P(a \le X \le b) = P(a < X < b) = P(a < X \le b) = P(a \le X < b),$$

as $P(X = a) = P(X = b) = 0.$

Mean and variance

Continuous random variable

Probability density functions

Cumulativ distributio functions

Common distribution

The expected value (mean) of a continuous random variable X is:

$$\mathbb{E}[X] = \int_{-\infty}^{\infty} x f(x) \mathrm{d}x.$$

Mean and variance

Continuous random variable

Probability density functions

Cumulativo distribution functions

Common distribution The expected value (mean) of a continuous random variable X is:

$$\mathbb{E}[X] = \int_{-\infty}^{\infty} x f(x) \mathrm{d}x.$$

The expected value of a function g(x) of X is:

$$\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)f(x)dx.$$

Mean and variance

Continuous random variable

Probability density functions

Cumulative distribution functions

Common distribution The expected value (mean) of a continuous random variable X is:

$$\mathbb{E}[X] = \int_{-\infty}^{\infty} x f(x) \mathrm{d}x.$$

The expected value of a function g(x) of X is:

$$\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)f(x)dx.$$

The variance of X is:

$$Var[X] = \mathbb{E}[(x - \mathbb{E}[X])^2]$$
$$= \int_{-\infty}^{\infty} (x - \mathbb{E}[X])^2 f(x) dx.$$

Example: the uniform distribution

Continuous random variable

Probability density functions

Cumulativ distributio functions

Common distributions When X has a uniform distribution:

$$f(x) = \begin{cases} \frac{1}{b-a} & \text{if } a \le x \le b \\ 0 & \text{otherwise.} \end{cases}$$

Example: the uniform distribution

Continuous random variable

Probability density functions

Cumulativ distributio functions

Common distributions When X has a uniform distribution:

$$f(x) = \begin{cases} \frac{1}{b-a} & \text{if } a \le x \le b \\ 0 & \text{otherwise.} \end{cases}$$

The expected value of X is:

$$\mathbb{E}[X] = \int_a^b x \frac{1}{b-a} dx = \frac{a+b}{2}.$$

Example: the uniform distribution

Continuous random variable

Probability density functions

Cumulativ distributio functions

Common distribution: When X has a uniform distribution:

$$f(x) = \begin{cases} \frac{1}{b-a} & \text{if } a \le x \le b \\ 0 & \text{otherwise.} \end{cases}$$

The expected value of X is:

$$\mathbb{E}[X] = \int_a^b x \frac{1}{b-a} \mathrm{d}x = \frac{a+b}{2}.$$

The variance of X is:

$$Var[X] = \int_{a}^{b} \left(x - \frac{a+b}{2}\right)^{2} \frac{1}{b-a} dx = \frac{(b-a)^{2}}{12}.$$

The cumulative distribution function


Continuous random variables

Probability density function

Cumulative distribution functions

Common distribution

The cumulative distribution function (CDF) F(x) is the area under the probability density function f(x) to the left of x.



The cumulative distribution function

Continuous random variables

Probability density function

Cumulative distribution functions

Common distribution:

Definition

Let X be a continuous random variable with probability density function f(x), then the cumulative distribution function is defined as

$$F(x) = P(X \le x)$$
$$= \int_{-\infty}^{x} f(t)dt.$$

The cumulative distribution function

Continuous random variables

Probability density functio

Cumulative distribution functions

Common distribution:

Definition

Let X be a continuous random variable with probability density function f(x), then the cumulative distribution function is defined as

$$F(x) = P(X \le x)$$
$$= \int_{-\infty}^{x} f(t)dt.$$

The CDF is a monotonically-increasing continuous function $F: \mathbb{R} \mapsto [0,1]$ satisfying $\lim_{x \to -\infty} F(x) = 0$ and $\lim_{x \to \infty} F(x) = 1$.


Computing probabilities using the CDF

Continuous random variables

Probability density function


Cumulative distribution functions


Common distribution:



Computing probabilities using the CDF

Cumulative distribution functions





Obtaining the PDF from the CDF

Continuous random variables

Probability density function

Cumulative distribution functions

Common distribution At every x at which the derivative F'(x) exists, F'(x) = f(x).

Obtaining the PDF from the CDF

Continuous random variable

Probability density functio

Cumulative distribution functions

distribution

At every x at which the derivative F'(x) exists, F'(x) = f(x).

Examples

When X has a uniform distribution, for a < x < b:

$$F'(x) = \frac{d}{dx} \left(\frac{x - a}{b - a} \right) = \frac{1}{b - a} = f(x)$$

Sampling using the CDF

Continuous random variables

Probability density function

Cumulative distribution functions

Common distribution

The inverse transform sampling algorithm can be used to sample a continuous random variable using the inverse of its cumulative distribution function.

Sampling using the CDF

Continuous random variables

Probability density function

Cumulative distribution functions

Common distribution

The inverse transform sampling algorithm can be used to sample a continuous random variable using the inverse of its cumulative distribution function.

Recall that $F: \mathbb{R} \mapsto [0,1]$.

Sampling using the CDF

Continuous random variable

Probability density function

Cumulative distribution functions

Common distribution The inverse transform sampling algorithm can be used to sample a continuous random variable using the inverse of its cumulative distribution function.

Recall that $F: \mathbb{R} \mapsto [0,1]$.

To draw a sample $x \sim f(x)$:

- **③** Sample $u \sim \mathrm{U}\left(0,1\right)$ (using a pseudo-random number generator)
- **2** Let $x = F^{-1}(u)$

Continuous random variable

Probability density function

Cumulative distribution functions

Common distribution: The PDF of the exponential distribution is:

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & \text{if } x \ge 0 \\ 0 & \text{otherwise,} \end{cases}$$

Continuous random variable

Probability density function

Cumulative distribution functions

Common distribution: The PDF of the exponential distribution is:

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & \text{if } x \ge 0\\ 0 & \text{otherwise,} \end{cases}$$

which implies that the CDF is:

$$F(x) = 1 - e^{-\lambda x} = u,$$

Continuous random variable

Probability density function

Cumulative distribution functions

Common distribution: The PDF of the exponential distribution is:

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & \text{if } x \ge 0\\ 0 & \text{otherwise,} \end{cases}$$

which implies that the CDF is:

$$F(x) = 1 - e^{-\lambda x} = u,$$

and the inverse of the CDF is:

$$F^{-1}(u) = -\frac{\log(1-u)}{\lambda} = x.$$

Continuous random variable

Probability density function

Cumulative distribution functions

Common distributions The PDF of the exponential distribution is:

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & \text{if } x \ge 0\\ 0 & \text{otherwise,} \end{cases}$$

which implies that the CDF is:

$$F(x) = 1 - e^{-\lambda x} = u,$$

and the inverse of the CDF is:

$$F^{-1}(u) = -\frac{\log(1-u)}{\lambda} = x.$$

Hence, to sample $x \sim f(x)$:

- Sample $u \sim U(0,1)$
- 2 Let $x = -\frac{\log(1-u)}{1}$

The normal (Gaussian) distribution

Continuous random variables

Probability density function

Cumulative distribution functions

Common distributions