

Exercises: unsupervised inference with iid observations and the linear Gaussian model

Joaquín Rapela*

August 6, 2023

1 Inferring location of a static submarine from its sonar measurements

This exercise is an extension of Example 3.3.4 from another great book in machine learning, [Murphy \(2022\)](#)¹. A static submarine is located in a 2D planar surface deep in the sea. A priori, we model its unknown location with a 2D random variable \mathbf{z} with

$$p(\mathbf{z}) = \mathcal{N}(\mathbf{z}|\mu_z, \Sigma_z) \quad (1)$$

We obtain noisy measurements of the location of the submarine with a sonar. We represent a 2D sonar measurement with random variable \mathbf{y}_n , with

$$p(\mathbf{y}_n|\mathbf{z}) = \mathcal{N}(\mathbf{y}_n|\mathbf{z}, \Sigma_y) \quad (2)$$

- (a) Sample 100 a priory locations of the submarine (i.e., $\mathbf{z}_1, \dots, \mathbf{z}_{100}$) using a mean $\mu_z = [2, 3]^\top$, a standard deviation along the horizontal direction $\sigma_{zx} = 1.0$, a standard deviation along the vertical direction $\sigma_{zy} = 1.0$, and a correlation coefficient between the vertical and horizontal directions $\rho_z = -0.8$.

*j.rapela@ucl.ac.uk

¹See this related [discussion](#) and its [follow-up](#).

Plot μ_z , the samples of the a-priori submarine location. and the ellipse containing 95% of samples of a-priori submarine locations. We call this ellipse the 95% prediction ellipse for samples of the a-priori submarine location.

You may want to complete the script `doExSubmarine.a.py` to address this item.

- (b) Select the submarine location \mathbf{z}_1 generated in the previous item. Sample $N = 5$ sonar measurements, assuming the submarine is at location \mathbf{z}_1 (i.e., sample from $p(\mathbf{y}|\mathbf{z}_1)$ to obtain $\mathbf{y}_1, \dots, \mathbf{y}_N$). Use a standard deviation of 1.0 for the measurement noise along the horizontal and vertical directions, and a correlation coefficient between the vertical and horizontal directions $\rho_{\mathbf{y}} = 0.8$.

Plot \mathbf{z}_1 , the sonar measurements samples. and the 95% prediction ellipse for sonar measurement samples.

You may want to complete the script `doExSubmarine.b.py` to address this item.

- (c) derive a mathematical expression for the posterior of the submarine location, given sonar measurements; i.e., $p(\mathbf{z}|\mathbf{y}_1, \dots, \mathbf{y}_N)$.

Hints:

- The posterior of the submarine location given sonar measurements is proportional to the joint distribution of the submarine location and sonar measurements; i.e., $p(\mathbf{z}|\mathbf{y}_1, \dots, \mathbf{y}_N) = \frac{p(\mathbf{y}_1, \dots, \mathbf{y}_N, \mathbf{z})}{p(\mathbf{y}_1, \dots, \mathbf{y}_N)} = K p(\mathbf{y}_1, \dots, \mathbf{y}_N, \mathbf{z})$, where K is a value that does not depend on \mathbf{z} . Thus, to obtain the posterior we can just keep the terms of the joint that depend on \mathbf{z} and normalise the resulting expression to integrate to one.
- The joint is the product of the likelihood and the prior; i.e., $p(\mathbf{y}_1, \dots, \mathbf{y}_N, \mathbf{z}) = p(\mathbf{y}_1, \dots, \mathbf{y}_N|\mathbf{z})p(\mathbf{z})$. Thus, to keep the terms of the joint that depend on \mathbf{z} , we can just keep the term of the likelihood that depend on \mathbf{z} and combine the result with the prior.
- As shown in Claim. 1, the terms of the likelihood that depend on \mathbf{z} are proportional to a Gaussian distribution with mean \mathbf{z} and covariance $\frac{1}{N}\Sigma$; i.e., $p(\mathbf{y}_1, \dots, \mathbf{y}_N|\mathbf{z}) = K \mathcal{N}(\bar{\mathbf{y}}|\mathbf{z}, \frac{1}{N}\Sigma_y)$, where K is a value that does not depend on \mathbf{z} .

- From the previous arguments, to obtain the posterior of \mathbf{z} we can multiply $\mathcal{N}(\bar{\mathbf{y}}|\mathbf{z}, \frac{1}{N}\Sigma_y)$ with the prior $p(\mathbf{z}) = \mathcal{N}(\mathbf{z}|\mu_z, \Sigma_z)$ and normalise the result. To do this we can use the expression for the posterior of the linear Gaussian model described in class.
- (d) plot the sample mean of the measurements, the 95% prediction ellipse for sample means of the measurements, the mean of the posterior distribution and the 95% prediction ellipse for samples from the posterior. You may want to complete the script `doExSubmarine.d.py` to address this item.
- (e) repeat (b) and (d) with $N \in \{1, 10, 100, 1000\}$ measurements, and show the plots generated in (d). How do the posterior and sample mean estimates change as N increases?
- (f) write expressions of the posterior mean and covariances to show that:
- as the number of measurements increases, the relative contribution of the prior to estimates of the posterior mean and covariance decreases,
 - in the limit when the number of measurements approaches infinity, the posterior covariance approaches zero and the posterior mean approaches the measurements sample mean. That is, for an infinite number of measurements, the posterior estimate becomes deterministic and the contribution of the prior to this estimate disappears.

Can you see these points in the previous simulations?

Claim 1. If $P(\mathbf{y}_i|\mathbf{z}) = \mathcal{N}(\mathbf{y}_i|\mathbf{z}, \Sigma)$, $i = 1, \dots, N$,
and $P(\mathbf{y}_1, \dots, \mathbf{y}_N|\mathbf{z}) = \prod_{i=1}^N P(\mathbf{y}_i|\mathbf{z})$,
then $P(\mathbf{y}_1, \dots, \mathbf{y}_N|\mathbf{z}) = K\mathcal{N}(\bar{\mathbf{y}}_N|\mathbf{z}, \frac{1}{N}\Sigma)$
where K is a value unrelated to \mathbf{z} .

Proof. By induction: $P_n = P(\mathbf{y}_1, \dots, \mathbf{y}_n|\mathbf{z}) = K\mathcal{N}(\bar{\mathbf{y}}_n|\mathbf{z}, \frac{1}{n}\Sigma)$
 P_1 :

$$P(\mathbf{y}_1|\mathbf{z}) = \mathcal{N}(\mathbf{y}_1|\mathbf{z}, \Sigma) = \mathcal{N}(\bar{\mathbf{y}}_1|\mathbf{z}, \frac{1}{1}\Sigma)$$

$$P_n \rightarrow P_{n+1}:$$

$$\begin{aligned} P(\mathbf{y}_1, \dots, \mathbf{y}_n, \mathbf{y}_{n+1}|\mathbf{z}) &= \prod_{i=1}^{n+1} P(\mathbf{y}_i|\mathbf{z}) \\ &= P(\mathbf{y}_1, \dots, \mathbf{y}_n|\mathbf{z})P(\mathbf{y}_{n+1}|\mathbf{z}) \\ &= \mathcal{N}(\bar{\mathbf{y}}_n|\mathbf{z}, \frac{1}{n}\Sigma)\mathcal{N}(\mathbf{y}_{n+1}|\mathbf{z}, \Sigma) \end{aligned}$$

then

$$\begin{aligned} \log P(\mathbf{y}_1, \dots, \mathbf{y}_n, \mathbf{y}_{n+1}|\mathbf{z}) &= K - \frac{1}{2}(\bar{\mathbf{y}}_n - \mathbf{z})^\top n\Sigma^{-1}(\bar{\mathbf{y}}_n - \mathbf{z}) - \\ &\quad \frac{1}{2}(\mathbf{y}_{n+1} - \mathbf{z})^\top \Sigma^{-1}(\mathbf{y}_{n+1} - \mathbf{z}) \\ &= K_1 - \frac{1}{2}(\mathbf{z}^\top(n+1)\Sigma^{-1}\mathbf{z} - 2\mathbf{z}^\top n\Sigma^{-1}\bar{\mathbf{y}}_n - 2\mathbf{z}^\top \Sigma^{-1}\mathbf{y}_{n+1}) \\ &= K_1 - \frac{1}{2}\left(\mathbf{z}^\top(n+1)\Sigma^{-1}\mathbf{z} - 2\mathbf{z}^\top \Sigma^{-1}\sum_{i=1}^n \mathbf{y}_i - 2\mathbf{z}^\top \Sigma^{-1}\mathbf{y}_{n+1}\right) \\ &= K_1 - \frac{1}{2}\left(\mathbf{z}^\top(n+1)\Sigma^{-1}\mathbf{z} - 2\mathbf{z}^\top \Sigma^{-1}\sum_{i=1}^{n+1} \mathbf{y}_i\right) \\ &= K_1 - \frac{1}{2}(\mathbf{z}^\top(n+1)\Sigma^{-1}\mathbf{z} - 2\mathbf{z}^\top(n+1)\Sigma^{-1}\bar{\mathbf{y}}_{n+1}) \end{aligned}$$

Therefore

$$P(\mathbf{y}_1, \dots, \mathbf{y}_n, \mathbf{y}_{n+1} | \mathbf{z}) = K_2 \mathcal{N} \left(\mathbf{z} \left| \bar{\mathbf{y}}_{n+1}, \frac{1}{n+1} \Sigma \right. \right) = K_2 \mathcal{N} \left(\bar{\mathbf{y}}_{n+1} \left| \mathbf{z}, \frac{1}{n+1} \Sigma \right. \right)$$

□

References

Murphy, K. P. (2022). *Probabilistic machine learning: an introduction*. MIT press.