

Foundations of probability theory

Historical notes

Axiomatic definition
of probability

Some basic rules

Conditional probability and Bayes

Conditional
probability

Assigning
probabilities by
conditioning

Independent events

Law of conditional
probability

Baye's rule in odds
form

Bayesian inference –
discrete case

References

Foundations of probability theory

Joaquín Rapela

Gatsby Computational Neuroscience Unit
University College London

June 14, 2023

Contents

Foundations of probability theory

Historical notes
Axiomatic definition of probability
Some basic rules

Conditional probability and Bayes

Conditional probability
Assigning probabilities by conditioning
Independent events
Law of conditional probability
Baye's rule in odds form
Bayesian inference – discrete case

References

- 1 Foundations of probability theory
 - Historical notes
 - Axiomatic definition of probability
 - Some basic rules
- 2 Conditional probability and Bayes
 - Conditional probability
 - Assigning probabilities by conditioning
 - Independent events
 - Law of conditional probability
 - Baye's rule in odds form
 - Bayesian inference – discrete case

Main reference

Foundations of probability theory

- Historical notes
- Axiomatic definition of probability
- Some basic rules

Conditional probability and Bayes

- Conditional probability
- Assigning probabilities by conditioning
- Independent events
- Law of conditional probability
- Baye's rule in odds form
- Bayesian inference – discrete case

References

I will mainly follow chapters seven *Foundations of probability theory* and eight *Conditional probability and Bayes* from Tijms (2012).

Contents

Foundations of probability theory

- Historical notes
- Axiomatic definition of probability
- Some basic rules

Conditional probability and Bayes

- Conditional probability
- Assigning probabilities by conditioning
- Independent events
- Law of conditional probability
- Baye's rule in odds form
- Bayesian inference – discrete case

References

1 Foundations of probability theory

2 Conditional probability and Bayes

Historical notes

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- explain the frequency-based interpretation of probability.
- constructing the mathematical foundations of probability theory has proven to be a long-lasting process of trial and error.
- the approach of defining probability as relative frequencies of repeatable experiments lead to unsatisfactory theory (why?)
<https://www.jstor.org/stable/pdf/20115155.pdf>
- the frequency view of probability has a long history that goes back to Aristotle.
- in 1933 the Russian mathematician Andrej Kolmogorov (1903-1987) laid a satisfactory mathematical foundation of probability theory.

He created a set of axioms. Axioms state a number of minimal requirements that the probability objects should satisfy. From these few algorithms all claims of probability can be derived, as we will see.

Probability model

Foundations
of probability
theory

Historical notes

Axiomatic definition
of probability

Some basic rules

Conditional
probability
and Bayes

Conditional
probability

Assigning
probabilities by
conditioning

Independent events

Law of conditional
probability

Baye's rule in odds
form

Bayesian inference –
discrete case

References

Definition 1 (Probability model)

A probability model *is a mathematical representation of a random experiment.*

Probability model

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Definition 1 (Probability model)

A **probability model** is a mathematical representation of a random experiment. It consists of a description of all possible outcomes of the experiment (i.e., **sample space**), a set of subsets of the sample space (i.e., **events**), and an assignment of probability to events (i.e., **probability measure**).

Sample space

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Definition 2 (Sample space)

*The set of all samples in an experiment is called the **sample space**. It is denoted by Ω .*

Event

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Definition 3 (Event)

*An **event** is a subset of the sample space. We denote the collection of all events by \mathcal{F} .*

Event

Foundations
of probability
theory

Historical notes

Axiomatic definition
of probability

Some basic rules

Conditional
probability
and Bayes

Conditional
probability

Assigning
probabilities by
conditioning

Independent events

Law of conditional
probability

Baye's rule in odds
form

Bayesian inference –
discrete case

References

Definition 3 (Event)

*An **event** is a subset of the sample space. We denote the collection of all events by \mathcal{F} .*

Notes:

- ① We will only assign probabilities to events (i.e., to sets $A \in \mathcal{F}$).
- ② For finite or countable sample spaces, we can assign probabilities to any subset of the sample space. Thus, any subset of a finite or countable sample space can be an event.
- ③ For uncountable sample spaces, we can only assign probabilities to well behaved subsets of the sample space (i.e., to elements in a σ algebra of subsets of the sample space). Only well-behaved subsets of an uncountable sample space can be events.

Probability measure

Foundations
of probability
theory

Historical notes

Axiomatic definition
of probability

Some basic rules

Conditional
probability
and Bayes

Conditional
probability

Assigning
probabilities by
conditioning

Independent events

Law of conditional
probability

Baye's rule in odds
form

Bayesian inference –
discrete case

References

Definition 4 (Probability measure)

A probability measure is a function that assigns numbers between zero and one to events (i.e., $P : \mathcal{F} \rightarrow [0, 1]$).

Probability model: definitions

Foundations
of probability
theory

Historical notes
Axiomatic definition
of probability
Some basic rules

Conditional
probability
and Bayes

Conditional
probability
Assigning
probabilities by
conditioning
Independent events

Law of conditional
probability

Baye's rule in odds
form

Bayesian inference –
discrete case

References

Definition (Sample space)

*The set of all samples in an experiment is called the **sample space**. It is denoted by Ω .*

Definition (Event)

*An **event** is a subset of the sample space. We denote the collection of all events by \mathcal{F} .*

Definition (Probability measure)

*A **probability measure** is a function that assigns numbers between zero and one to events (i.e., $P : \mathcal{F} \rightarrow [0, 1]$).*

Definition (Probability model)

*A **probability model**, \mathcal{M} , is a mathematical representation of a random experiment consisting of a sample space, Ω , a set of events, \mathcal{F} , and a probability measure, P (i.e., $\mathcal{M} = \{\Omega, \mathcal{F}, P\}$).*

Examples of probability models

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

For each of the following examples, let's find the sample space and propose a probability measure.

- 1 The experiment is to toss a fair coin once.

Examples of probability models

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

For each of the following examples, let's find the sample space and propose a probability measure.

- 1 The experiment is to toss a fair coin once. The sample space is the set $[H, T]$.

Examples of probability models

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

For each of the following examples, let's find the sample space and propose a probability measure.

- 1 The experiment is to toss a fair coin once. The sample space is the set $[H, T]$. We assign a probability of 0.5 to each element of the sample space.

Examples of probability models

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

For each of the following examples, let's find the sample space and propose a probability measure.

- 1 The experiment is to toss a fair coin once. The sample space is the set $[H, T]$. We assign a probability of 0.5 to each element of the sample space.
- 2 The experiment is to repeatedly roll a fair die and count the number of rolls until the first six shows up.

Examples of probability models

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

For each of the following examples, let's find the sample space and propose a probability measure.

- 1 The experiment is to toss a fair coin once. The sample space is the set $[H, T]$. We assign a probability of 0.5 to each element of the sample space.
- 2 The experiment is to repeatedly roll a fair die and count the number of rolls until the first six shows up. The sample space is the set of positive integers.

Examples of probability models

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

For each of the following examples, let's find the sample space and propose a probability measure.

- 1 The experiment is to toss a fair coin once. The sample space is the set $[H, T]$. We assign a probability of 0.5 to each element of the sample space.
- 2 The experiment is to repeatedly roll a fair die and count the number of rolls until the first six shows up. The sample space is the set of positive integers. The probabilities $\frac{1}{6}, \frac{5}{6} \times \frac{1}{6}, \left(\frac{5}{6}\right)^2 \times \frac{1}{6}, \dots$ can be assigned to the outcomes $1, 2, 3, \dots$

Examples of probability models

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

For each of the following examples, let's find the sample space and propose a probability measure.

- 1 The experiment is to toss a fair coin once. The sample space is the set $[H, T]$. We assign a probability of 0.5 to each element of the sample space.
- 2 The experiment is to repeatedly roll a fair die and count the number of rolls until the first six shows up. The sample space is the set of positive integers. The probabilities $\frac{1}{6}, \frac{5}{6} \times \frac{1}{6}, \left(\frac{5}{6}\right)^2 \times \frac{1}{6}, \dots$ can be assigned to the outcomes $1, 2, 3, \dots$.
- 3 The experiment is to measure the time between the first and second spikes in an experimental trial.

Examples of probability models

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

For each of the following examples, let's find the sample space and propose a probability measure.

- 1 The experiment is to toss a fair coin once. The sample space is the set $[H, T]$. We assign a probability of 0.5 to each element of the sample space.
- 2 The experiment is to repeatedly roll a fair die and count the number of rolls until the first six shows up. The sample space is the set of positive integers. The probabilities $\frac{1}{6}, \frac{5}{6} \times \frac{1}{6}, \left(\frac{5}{6}\right)^2 \times \frac{1}{6}, \dots$ can be assigned to the outcomes $1, 2, 3, \dots$.
- 3 The experiment is to measure the time between the first and second spikes in an experimental trial.

Examples of probability models

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

For each of the following examples, let's find the sample space and propose a probability measure.

- 1 The experiment is to toss a fair coin once. The sample space is the set $[H, T]$. We assign a probability of 0.5 to each element of the sample space.
- 2 The experiment is to repeatedly roll a fair die and count the number of rolls until the first six shows up. The sample space is the set of positive integers. The probabilities $\frac{1}{6}, \frac{5}{6} \times \frac{1}{6}, \left(\frac{5}{6}\right)^2 \times \frac{1}{6}, \dots$ can be assigned to the outcomes $1, 2, 3, \dots$
- 3 The experiment is to measure the time between the first and second spikes in an experimental trial. The sample space is the set $(0, \infty)$ of positive real numbers. We can assign a probability of $1 - \exp(-\lambda t)$ to the event that the second spike is fired less than t seconds after the first spike.

Axioms of probability theory

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Axiom 1

$$P(A) \geq 0, \quad \forall A \in \mathcal{F}$$

Axiom 2

$$P(\Omega) = 1$$

Axiom 3

$P(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i)$ for every collection of pairwise disjoint events A_1, A_2, \dots

Experiment with equally likely outcomes

Foundations of probability theory

Historical notes

Axiomatic definition
of probability

Some basic rules

Conditional probability and Bayes

Conditional
probability

Assigning
probabilities by
conditioning

Independent events

Law of conditional
probability

Bayes's rule in odds
form

Bayesian inference –
discrete case

References

An experiment with equally likely outcomes is one with a finite number of outcomes $\omega_1, \dots, \omega_N$, where all outcomes have the same probability (i.e., $P(\omega_i) = \frac{1}{N}$).

Experiment with equally likely outcomes

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

An experiment with equally likely outcomes is one with a finite number of outcomes $\omega_1, \dots, \omega_N$, where all outcomes have the same probability (i.e., $P(\omega_i) = \frac{1}{N}$).

Claim 1

For any event A , $P(A) = \frac{N(A)}{N}$, where $N(A)$ is the number of outcomes in the set A .

Example: equally likely outcomes

Foundations
of probability
theory

Historical notes

Axiomatic definition
of probability

Some basic rules

Conditional
probability
and Bayes

Conditional
probability

Assigning
probabilities by
conditioning

Independent events

Law of conditional
probability

Baye's rule in odds
form

Bayesian inference –
discrete case

References

Example 1

John, Pedro and Rosita each roll on fair die. How do we calculate the probability that the score of Rosita is equal to the sum of the scores of John and Pedro?

Some basic rules

Foundations of probability theory

Historical notes
Axiomatic definition of probability
Some basic rules

Conditional probability and Bayes

Conditional probability
Assigning probabilities by conditioning
Independent events
Law of conditional probability
Bayes's rule in odds form
Bayesian inference – discrete case

References

Rule 1

For any finite number of mutually exclusive events A_1, \dots, A_N ,

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + \dots + P(A_N)$$

Rule 2

For any event A ,

$$P(A) = 1 - P(A^c)$$

where the event A^c consists of all outcomes that are not in A .

Rule 3

Let A, B be two events such that $A \subseteq B$. Then $P(A) \leq P(B)$.

Rule 4

For any two events A and B ,

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

Rule 1

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Proof.

Denote by \emptyset the empty set. We will first prove that $P(\emptyset) = 0$. Take $A_i = \emptyset$ for $i = 1, 2, \dots$. Then $\emptyset = \bigcup_{i=1}^{\infty} A_i$. Next, by Axiom 3, $P(\emptyset) = \sum_{i=1}^{\infty} P(A_i) = \sum_{i=1}^{\infty} P(\emptyset)$. This implies that $P(\emptyset) = 0$. Define $A_{N+j} = \emptyset$ for $j = 1, 2, \dots$. Then

$$\begin{aligned} P(A_1 \cup A_2 \cup \dots \cup A_N) &= P(A_1 \cup A_2 \cup \dots \cup A_N \cup A_{N+1} \cup A_{N+2} \cup \dots) \\ &= P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i) \\ &= \sum_{i=1}^N P(A_i) + \sum_{j=1}^{\infty} P(A_{N+j}) = \sum_{i=1}^N P(A_i) \end{aligned}$$



Notes:

- 1 the last equality in the second line holds by Axiom 1
- 2 the last equality in the third line holds because $P(A_{N+j}) = P(\emptyset) = 0$.

Proof of rule 2

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Proof.

$\Omega = A \cup A^c$. A and A^c are disjoint. Then, by Rule 1, $P(\Omega) = P(A) + P(A^c)$. From Axiom 2, $P(\Omega) = 1$. Thus, $1 = P(A) + P(A^c)$.



Exercise

Foundations of probability theory

Historical notes

Axiomatic definition
of probability

Some basic rules

Conditional probability and Bayes

Conditional
probability

Assigning
probabilities by
conditioning

Independent events

Law of conditional
probability

Baye's rule in odds
form

Bayesian inference –
discrete case

References

Prove rule 3.

Proof of rule 4

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Proof.

$$A \cup B = (A \setminus B) \cup (B \setminus A) \cup (A \cap B)$$

$$A = (A \setminus B) \cup (A \cap B)$$

$$B = (B \setminus A) \cup (A \cap B)$$

Since the sets in the right-hand-side of the above equations are pairwise disjoint, by rule 1, we obtain

$$P(A \cup B) = P(A \setminus B) + P(B \setminus A) + P(A \cap B)$$

$$P(A) = P(A \setminus B) + P(A \cap B) \rightarrow P(A \setminus B) = P(A) - P(A \cap B)$$

$$P(B) = P(B \setminus A) + P(A \cap B) \rightarrow P(B \setminus A) = P(B) - P(A \cap B)$$

Replacing the equations on the right of the second and third line in the equation on the first line the rule is proved.

$$\begin{aligned} P(A \cup B) &= P(A) - P(A \cap B) + P(B) - P(A \cap B) + P(A \cap B) \\ &= P(A) + P(B) - P(A \cap B) \end{aligned}$$



Example 7.7: Chevalier de Mere to Blaise Pascal

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

The gambler Chevalier de Mere posed the following problem to the famous French mathematician Blaise Pascal in 1654. This problem marks the beginning of probability theory.

Example 7.7

- a *How many rolls of a fair die are required to have at least a 50% chance of rolling at least one six?*
- b *How many rolls of two fair dice are required to have at least a 50% chance of rolling at least one double six?*

Analytical solution to example 7.7a

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

(a)

- Let's fix the number of rolls r . The sample space is $\Omega = \{(i_1, \dots, i_r) : 1 \leq i_k \leq 6\}$, where i_k is the up face of the die on the k th roll. The outcomes in Ω are equiprobable.

Analytical solution to example 7.7a

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

(a)

- Let's fix the number of rolls r . The sample space is $\Omega = \{(i_1, \dots, i_r) : 1 \leq i_k \leq 6\}$, where i_k is the up face of the die on the k th roll. The outcomes in Ω are equiprobable.
- We want to calculate the probability of the event A = "at least one six shows up in the r rolls.". When you see the keyword "at least one" in an event, it is easier to calculate the probability of the complement A^c = "no six shows up in the r rolls."

Analytical solution to example 7.7a

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

(a)

- Let's fix the number of rolls r . The sample space is $\Omega = \{(i_1, \dots, i_r) : 1 \leq i_k \leq 6\}$, where i_k is the up face of the die on the k th roll. The outcomes in Ω are equiprobable.
- We want to calculate the probability of the event A = "at least one six shows up in the r rolls." When you see the keyword "at least one" in an event, it is easier to calculate the probability of the complement A^c = "no six shows up in the r rolls."
- $$P(A^c) = \frac{N(A^c)}{N} = \frac{5^r}{6^r} = \left(\frac{5}{6}\right)^r.$$

Analytical solution to example 7.7a

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

(a)

- Let's fix the number of rolls r . The sample space is $\Omega = \{(i_1, \dots, i_r) : 1 \leq i_k \leq 6\}$, where i_k is the up face of the die on the k th roll. The outcomes in Ω are equiprobable.
- We want to calculate the probability of the event A = "at least one six shows up in the r rolls." When you see the keyword "at least one" in an event, it is easier to calculate the probability of the complement A^c = "no six shows up in the r rolls."
- $P(A^c) = \frac{N(A^c)}{N} = \frac{5^r}{6^r} = \left(\frac{5}{6}\right)^r$.
- $\frac{1}{2} < P(A) = 1 - P(A^c) = 1 - \left(\frac{5}{6}\right)^r$ iff $\left(\frac{5}{6}\right)^r < \frac{1}{2}$ iff $\log \left(\frac{5}{6}\right)^r < \log \frac{1}{2}$ iff $r \log \frac{5}{6} < \log \frac{1}{2}$ iff $r > \frac{\log \frac{1}{2}}{\log \frac{5}{6}} = 3.8$

Simulated solution to example 7.7a

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Please see code [here](#).

Exercise

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Solve example 7.7b analytically and by simulation. Answer: you need at least 25 draws.

Example 7.9: soccer teams in quarterfinal

Foundations
of probability
theory

Historical notes

Axiomatic definition
of probability

Some basic rules

Conditional
probability
and Bayes

Conditional
probability

Assigning
probabilities by
conditioning

Independent events

Law of conditional
probability

Baye's rule in odds
form

Bayesian inference –
discrete case

References

Example 7.9

The eight soccer teams which have reached the quarterfinals of the Championship League are formed by two teams from each of the countries England, Germany, Italy and Spain. The four matches to be played in the quarterfinal are determined by drawing lots.

- a *What is the probability that the two teams from the same country play against each other in each of the four matches?*
- b *What is the probability that there is a match between the two teams from England or between the two teams from Germany?*

Analytical solution to example 7.9: setup

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- Let T be the set of eight teams
$$T = \{E_1, E_2, G_1, G_2, I_1, I_2, S_1, S_2\}$$

Analytical solution to example 7.9: setup

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- Let T be the set of eight teams
$$T = \{E_1, E_2, G_1, G_2, I_1, I_2, S_1, S_2\}$$
- A match is a set of two different elements from T (i.e., $\text{match} = \{T_i, T_j\}, \text{match} \subset T, T_i \neq T_j$).

Analytical solution to example 7.9: setup

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- Let T be the set of eight teams
$$T = \{E_1, E_2, G_1, G_2, I_1, I_2, S_1, S_2\}$$
- A match is a set of two different elements from T (i.e., $\text{match} = \{T_i, T_j\}, \text{match} \subset T, T_i \neq T_j$).
- A quarterfinal is a set of four different matches.

Analytical solution to example 7.9: setup

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- Let T be the set of eight teams
$$T = \{E_1, E_2, G_1, G_2, I_1, I_2, S_1, S_2\}$$
- A match is a set of two different elements from T (i.e., $\text{match} = \{T_i, T_j\}, \text{match} \subset T, T_i \neq T_j$).
- A quarterfinal is a set of four different matches.
- Ω is the set of all possible quarterfinals (e.g., $\{\{E_1, G_1\}, \{I_1, G_2\}, \{S_2, G_2\}, \{I_2, G_1\}\} \in \Omega$).

Analytical solution to example 7.9: setup

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- Let T be the set of eight teams
$$T = \{E_1, E_2, G_1, G_2, I_1, I_2, S_1, S_2\}$$
- A match is a set of two different elements from T (i.e., $\text{match} = \{T_i, T_j\}, \text{match} \subset T, T_i \neq T_j$).
- A quarterfinal is a set of four different matches.
- Ω is the set of all possible quarterfinals (e.g., $\{\{E_1, G_1\}, \{I_1, G_2\}, \{S_2, G_2\}, \{I_2, G_1\}\} \in \Omega$).
- Ω contains equally-likely outcomes. Thus, for any event A ,
$$P(A) = \frac{N(A)}{N}.$$

Analytical solution to example 7.9: setup

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- Let T be the set of eight teams
$$T = \{E_1, E_2, G_1, G_2, I_1, I_2, S_1, S_2\}$$
- A match is a set of two different elements from T (i.e., $\text{match} = \{T_i, T_j\}, \text{match} \subset T, T_i \neq T_j$).
- A quarterfinal is a set of four different matches.
- Ω is the set of all possible quarterfinals (e.g., $\{\{E_1, G_1\}, \{I_1, G_2\}, \{S_2, G_2\}, \{I_2, G_1\}\} \in \Omega$).
- Ω contains equally-likely outcomes. Thus, for any event A ,
$$P(A) = \frac{N(A)}{N}.$$
- $N = \frac{\binom{8}{2}\binom{6}{2}\binom{4}{2}}{4!}$. There are $\binom{8}{2}$, $\binom{6}{2}$, $\binom{4}{2}$ and 1 ways of selecting the first, second, third and fourth matches, respectively. We divide by $4!$ because the order between matches does not matter.

Analytical solution to example 7.9a

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- The event $A =$ “*two teams from the same country play against each other in each of the four matches*” contains only one outcome (i.e.,
$$A = \{\{E_1, E_2\}, \{G_1, G_2\}, \{I_1, I_2\}, \{S_1, S_2\}\}).$$

Analytical solution to example 7.9a

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- The event $A = \text{"two teams from the same country play against each other in each of the four matches"}$ contains only one outcome (i.e.,
 $A = \{\{E_1, E_2\}, \{G_1, G_2\}, \{I_1, I_2\}, \{S_1, S_2\}\}$).
- $$P(A) = \frac{N(A)}{N} = \frac{1}{\frac{\binom{8}{2}\binom{6}{2}\binom{4}{2}}{4!}} = 0.009524$$

Analytical solution to example 7.9b

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- The event $B = \text{"there is a match between the two teams from England or between the two teams from Germany"}$ is the union of the events $B_E = \text{"there is a match between the two teams from England"}$ and $B_G = \text{"there is a match between the two teams from Germany"}$ (i.e., $B = B_E \cup B_G$).

Analytical solution to example 7.9b

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- The event $B = \text{"there is a match between the two teams from England or between the two teams from Germany"}$ is the union of the events $B_E = \text{"there is a match between the two teams from England"}$ and $B_G = \text{"there is a match between the two teams from Germany"}$ (i.e., $B = B_E \cup B_G$).
- Since B_E and B_G are not disjoint, we should use Rule 3 to compute $P(B_E \cup B_G)$ (i.e., $P(B_E \cup B_G) = P(B_E) + P(B_G) - P(B_E \cap B_G)$).

Analytical solution to example 7.9b

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Bayes's rule in odds form

Bayesian inference – discrete case

References

- The event $B =$ “there is a match between the two teams from England or between the two teams from Germany” is the union of the events $B_E =$ “there is a match between the two teams from England” and $B_G =$ “there is a match between the two teams from Germany” (i.e., $B = B_E \cup B_G$).
- Since B_E and B_G are not disjoint, we should use Rule 3 to compute $P(B_E \cup B_G)$ (i.e., $P(B_E \cup B_G) = P(B_E) + P(B_G) - P(B_E \cap B_G)$).
- $P(B_E) = P(B_G)$. Since the match $\{E_1, E_2\}$ is in all quarterfinals in B_E , to calculate $N(B_E)$ we need to compute the number of matches between six teams of three countries, as done in part (a). This gives $N(B_E) = \frac{\binom{6}{2}\binom{4}{2}}{3!}$.

$$\text{Then } P(B_E) = P(B_G) = \frac{\frac{\binom{6}{2}\binom{4}{2}}{3!}}{\frac{\binom{8}{2}\binom{6}{2}\binom{4}{2}}{4!}} = 0.142857.$$

Analytical solution to example 7.9b

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Bayes's rule in odds form

Bayesian inference – discrete case

References

- The event $B =$ “there is a match between the two teams from England or between the two teams from Germany” is the union of the events $B_E =$ “there is a match between the two teams from England” and $B_G =$ “there is a match between the two teams from Germany” (i.e., $B = B_E \cup B_G$).
- Since B_E and B_G are not disjoint, we should use Rule 3 to compute $P(B_E \cup B_G)$ (i.e., $P(B_E \cup B_G) = P(B_E) + P(B_G) - P(B_E \cap B_G)$).
- $P(B_E) = P(B_G)$. Since the match $\{E_1, E_2\}$ is in all quarterfinals in B_E , to calculate $N(B_E)$ we need to compute the number of matches between six teams of three countries, as done in part (a). This gives $N(B_E) = \frac{\binom{6}{2}\binom{4}{2}}{3!}$.
Then $P(B_E) = P(B_G) = \frac{\frac{\binom{6}{2}\binom{4}{2}}{3!}}{\frac{\binom{8}{2}\binom{6}{2}\binom{4}{2}}{4!}} = 0.142857$.
- Since the match $\{E_1, E_2\}$ and $\{G_1, G_2\}$ are in all quarterfinals in $B_E \cap B_G$, as in part (a), $N(B_E \cap B_G) = \frac{\binom{4}{2}}{2!}$. Then
$$P(B_E \cap B_G) = \frac{\frac{\binom{4}{2}}{2!}}{\frac{\binom{8}{2}\binom{6}{2}\binom{4}{2}}{4!}} = 0.028571.$$

Analytical solution to example 7.9b

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Bayes's rule in odds form

Bayesian inference – discrete case

References

- The event $B =$ “there is a match between the two teams from England or between the two teams from Germany” is the union of the events $B_E =$ “there is a match between the two teams from England” and $B_G =$ “there is a match between the two teams from Germany” (i.e., $B = B_E \cup B_G$).
- Since B_E and B_G are not disjoint, we should use Rule 3 to compute $P(B_E \cup B_G)$ (i.e., $P(B_E \cup B_G) = P(B_E) + P(B_G) - P(B_E \cap B_G)$).
- $P(B_E) = P(B_G)$. Since the match $\{E_1, E_2\}$ is in all quarterfinals in B_E , to calculate $N(B_E)$ we need to compute the number of matches between six teams of three countries, as done in part (a). This gives $N(B_E) = \frac{\binom{6}{2}\binom{4}{2}}{3!}$.
Then $P(B_E) = P(B_G) = \frac{\frac{\binom{6}{2}\binom{4}{2}}{3!}}{\frac{\binom{8}{2}\binom{6}{2}\binom{4}{2}}{4!}} = 0.142857$.
- Since the match $\{E_1, E_2\}$ and $\{G_1, G_2\}$ are in all quarterfinals in $B_E \cap B_G$, as in part (a), $N(B_E \cap B_G) = \frac{\binom{4}{2}}{2!}$. Then
$$P(B_E \cap B_G) = \frac{\frac{\binom{4}{2}}{2!}}{\frac{\binom{8}{2}\binom{6}{2}\binom{4}{2}}{4!}} = 0.028571.$$
- Thus $P(B_E \cup B_G) = P(B_E) + P(B_G) - P(B_E \cap B_G) = 2 \times 0.142857 - 0.028571 = 0.257143$.

Simulated solution to example 7.9a

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Please see code [here](#).

Exercise

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Solve example 7.7b by simulation. Answer: you should obtain a solution close to the analytical one.

Contents

Foundations of probability theory

- Historical notes
- Axiomatic definition of probability
- Some basic rules

Conditional probability and Bayes

- Conditional probability
- Assigning probabilities by conditioning
- Independent events
- Law of conditional probability
- Baye's rule in odds form
- Bayesian inference – discrete case

References

1 Foundations of probability theory

2 Conditional probability and Bayes

Definition of conditional probability: motivation

- We are given a probability model (Ω, \mathcal{F}, P) .

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Definition of conditional probability: motivation

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- We are given a probability model (Ω, \mathcal{F}, P) .
- We are interested in the probability of event $A \in \mathcal{F}$. This model provides us the unconditioned probability of A , $P(A)$.

Definition of conditional probability: motivation

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- We are given a probability model (Ω, \mathcal{F}, P) .
- We are interested in the probability of event $A \in \mathcal{F}$. This model provides us the unconditioned probability of A , $P(A)$.
- Our colleague performs an experiment and tells us that event B occurred. How does the fact that B occurred changes the probability of A ?

$$P(A) \xrightarrow{\text{B occurred}} P(A|B)$$

Definition of conditional probability: motivation

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- We are given a probability model (Ω, \mathcal{F}, P) .
- We are interested in the probability of event $A \in \mathcal{F}$. This model provides us the unconditioned probability of A , $P(A)$.
- Our colleague performs an experiment and tells us that event B occurred. How does the fact that B occurred changes the probability of A ?

$$P(A) \xrightarrow{\text{B occurred}} P(A|B)$$

- Definition 1: $P(\cdot|B) = P(\cdot \cap B)$, where \cdot can be any event.

Definition of conditional probability: motivation

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- We are given a probability model (Ω, \mathcal{F}, P) .
- We are interested in the probability of event $A \in \mathcal{F}$. This model provides us the unconditioned probability of A , $P(A)$.
- Our colleague performs an experiment and tells us that event B occurred. How does the fact that B occurred changes the probability of A ?

$$P(A) \xrightarrow{\text{B occurred}} P(A|B)$$

- Definition 1: $P(\cdot|B) = P(\cdot \cap B)$, where \cdot can be any event.
- Problem with Definition 1: we want $P(B|B) = 1$, but from Definition 1, $P(B|B) = P(B \cap B) = P(B) \leq 1$.

Definition of conditional probability: motivation

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- We are given a probability model (Ω, \mathcal{F}, P) .
- We are interested in the probability of event $A \in \mathcal{F}$. This model provides us the unconditioned probability of A , $P(A)$.
- Our colleague performs an experiment and tells us that event B occurred. How does the fact that B occurred changes the probability of A ?

$$P(A) \xrightarrow{\text{B occurred}} P(A|B)$$

- Definition 1: $P(\cdot|B) = P(\cdot \cap B)$, where \cdot can be any event.
- Problem with Definition 1: we want $P(B|B) = 1$, but from Definition 1, $P(B|B) = P(B \cap B) = P(B) \leq 1$.
- Definition 2: $P(\cdot|B) = \frac{P(\cdot \cap B)}{P(B)}$.

Definition of conditional probability: motivation

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- We are given a probability model (Ω, \mathcal{F}, P) .
- We are interested in the probability of event $A \in \mathcal{F}$. This model provides us the unconditioned probability of A , $P(A)$.
- Our colleague performs an experiment and tells us that event B occurred. How does the fact that B occurred changes the probability of A ?

$$P(A) \xrightarrow{\text{B occurred}} P(A|B)$$

- Definition 1: $P(\cdot|B) = P(\cdot \cap B)$, where \cdot can be any event.
- Problem with Definition 1: we want $P(B|B) = 1$, but from Definition 1, $P(B|B) = P(B \cap B) = P(B) \leq 1$.
- Definition 2: $P(\cdot|B) = \frac{P(\cdot \cap B)}{P(B)}$.
 - $P(B|B) = 1 \checkmark$

Definition of conditional probability: motivation

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- We are given a probability model (Ω, \mathcal{F}, P) .
- We are interested in the probability of event $A \in \mathcal{F}$. This model provides us the unconditioned probability of A , $P(A)$.
- Our colleague performs an experiment and tells us that event B occurred. How does the fact that B occurred changes the probability of A ?

$$P(A) \xrightarrow{\text{B occurred}} P(A|B)$$

- Definition 1: $P(\cdot|B) = P(\cdot \cap B)$, where \cdot can be any event.
- Problem with Definition 1: we want $P(B|B) = 1$, but from Definition 1, $P(B|B) = P(B \cap B) = P(B) \leq 1$.
- Definition 2: $P(\cdot|B) = \frac{P(\cdot \cap B)}{P(B)}$.
 - $P(B|B) = 1 \checkmark$
 - $P(A) = P(A|\Omega) = \frac{P(A \cap \Omega)}{P(\Omega)} = \frac{P(A)}{1} = P(A) \checkmark$

Definition of conditional probability

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Definition 5 (Conditional probability)

For any two events A and B , with $P(B) > 0$, the conditional probability of A given B , $P(A|B)$, is defined as

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Example 8.1: conditional probability for two dice

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Example 8.1

Someone has rolled two dice. You know that one of the dice turned up a face value of six. What is the probability that the other die turned up a six as well?

Analytical solution to example 8.1

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- $\Omega = \{(i_1, i_2) : 1 \leq i_1, i_2 \leq 6\}$. $N = 36$. Equally probable outcomes.

Analytical solution to example 8.1

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Bayes's rule in odds form

Bayesian inference – discrete case

References

- $\Omega = \{(i_1, i_2) : 1 \leq i_1, i_2 \leq 6\}$. $N = 36$. Equally probable outcomes.
- $B = \text{"one die turned up a face value of six"} = \{(6, i), (j, 6), (6, 6) : 1 \leq i, j \leq 5\}$. $N(B) = 11$.
$$P(B) = \frac{N(B)}{N} = \frac{11}{36}.$$

Analytical solution to example 8.1

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Bayes's rule in odds form

Bayesian inference – discrete case

References

- $\Omega = \{(i_1, i_2) : 1 \leq i_1, i_2 \leq 6\}$. $N = 36$. Equally probable outcomes.
- $B =$ “one die turned up a face value of six” = $\{(6, i), (j, 6), (6, 6) : 1 \leq i, j \leq 5\}$. $N(B) = 11$.
$$P(B) = \frac{N(B)}{N} = \frac{11}{36}.$$
- $A =$ “the other die turned up a face value of six”

Analytical solution to example 8.1

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- $\Omega = \{(i_1, i_2) : 1 \leq i_1, i_2 \leq 6\}$. $N = 36$. Equally probable outcomes.
- $B =$ “one die turned up a face value of six” = $\{(6, i), (j, 6), (6, 6) : 1 \leq i, j \leq 5\}$. $N(B) = 11$.
 $P(B) = \frac{N(B)}{N} = \frac{11}{36}$.
- $A =$ “the other die turned up a face value of six”
- $A \cap B =$ “the two dice turned up a face value of six” = $\{(6, 6)\}$. $N(A \cap B) = 1$. $P(A \cap B) = \frac{N(A \cap B)}{N} = \frac{1}{36}$.

Analytical solution to example 8.1

Foundations of probability theory

Historical notes

Axiomatic definition
of probability

Some basic rules

Conditional probability and Bayes

Conditional
probability

Assigning
probabilities by
conditioning

Independent events

Law of conditional
probability

Bayes's rule in odds
form

Bayesian inference –
discrete case

References

- $\Omega = \{(i_1, i_2) : 1 \leq i_1, i_2 \leq 6\}$. $N = 36$. Equally probable outcomes.
- $B =$ “one die turned up a face value of six” = $\{(6, i), (j, 6), (6, 6) : 1 \leq i, j \leq 5\}$. $N(B) = 11$.
$$P(B) = \frac{N(B)}{N} = \frac{11}{36}.$$
- $A =$ “the other die turned up a face value of six”
- $A \cap B =$ “the two dice turned up a face value of six” = $\{(6, 6)\}$. $N(A \cap B) = 1$. $P(A \cap B) = \frac{N(A \cap B)}{N} = \frac{1}{36}.$
- Approach 1 [definition]: $P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{1/36}{11/36} = \frac{1}{11}.$

Analytical solution to example 8.1

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- $\Omega = \{(i_1, i_2) : 1 \leq i_1, i_2 \leq 6\}$. $N = 36$. Equally probable outcomes.
- $B =$ “one die turned up a face value of six” = $\{(6, i), (j, 6), (6, 6) : 1 \leq i, j \leq 5\}$. $N(B) = 11$.
 $P(B) = \frac{N(B)}{N} = \frac{11}{36}$.
- $A =$ “the other die turned up a face value of six”
- $A \cap B =$ “the two dice turned up a face value of six” = $\{(6, 6)\}$. $N(A \cap B) = 1$. $P(A \cap B) = \frac{N(A \cap B)}{N} = \frac{1}{36}$.
- Approach 1 [definition]: $P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{1/36}{11/36} = \frac{1}{11}$.
- Approach 2 [B as Ω]: $P(A|B) = \frac{N(A \cap B)}{N(B)} = \frac{1}{11}$.

Simulated solution to example 8.1

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Please see code [here](#).

Exercise

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Someone has rolled two dice. You know that the first die turned up a face value of six. What is the probability that the second die turned up a six as well?

Assigning probabilities by conditioning

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Rule 5

For any $n \in \mathbb{N}$, $n \geq 2$, and any sequence of events A_1, \dots, A_n ,

$$P(A_1 \cap \dots \cap A_n) = P(A_n | A_{n-1} \cap \dots \cap A_1) P(A_{n-1} | A_{n-2} \cap \dots \cap A_1) \dots P(A_1)$$

Proof of rule 5

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Proof.

By induction.

$$P_2: P(A_1 \cap A_2) = P(A_2|A_1)P(A_1)$$

$$P_n \rightarrow P_{n+1}:$$

$$\begin{aligned} P(A_1 \cap A_2 \cap \dots \cap A_n \cap A_{n+1}) &= P(B \cap A_{n+1}) \\ &= P(A_{n+1}|B)P(B) \\ &= P(A_{n+1}|A_n \cap A_{n-1} \cap \dots \cap A_1) \\ &\quad P(A_n \cap A_{n-1} \cap \dots \cap A_1) \\ &= P(A_{n+1}|A_n \cap A_{n-1} \cap \dots \cap A_1) \\ &\quad P(A_n|A_{n-1} \cap \dots \cap A_1) \\ &\quad P(A_{n-1}|A_{n-2} \cap \dots \cap A_1) \dots P(A_1) \end{aligned}$$



Notes:

- 1 P_2 follows from the definition of conditional probability.
- 2 in the first equality in $P_n \rightarrow P_{n+1}$ we defined the event $B = A_1 \cap \dots \cap A_n$.
- 3 in the second equality in $P_n \rightarrow P_{n+1}$ we used the definition of conditional probability.
- 4 in the third equality in $P_n \rightarrow P_{n+1}$ we replaced B by its definition.
- 5 in the fourth equality in $P_n \rightarrow P_{n+1}$ we used the inductive hypothesis $P(A_1 \cap \dots \cap A_n) = P(A_n|A_{n-1} \cap \dots \cap A_1)P(A_{n-1}|A_{n-2} \cap \dots \cap A_1) \dots P(A_1)$.

Example 8.3: allocating tourists to hotels

Foundations
of probability
theory

Historical notes

Axiomatic definition
of probability

Some basic rules

Conditional
probability
and Bayes

Conditional
probability

Assigning
probabilities by
conditioning

Independent events

Law of conditional
probability

Baye's rule in odds
form

Bayesian inference –
discrete case

References

Example 8.3

A group of 15 tourists is stranded in a city with four hotels of the same class. Each of the hotels has enough room available to accomodate the 15 tourists. The group's guide, who has a good working relationship with each of the four hotels, assigns the tourists to the hotels as follows. First, he randomly determines how many tourists will go to hotel A, then how many of the remaining tourists will go to hotel B, and next how many of the still reamining tourists will go to hotel C. All remaining tourists are sent to hotel D. At each stage of the assignment the guide draws a random number between zero and the number of tourists left.

- a** Calculate the probability of any assignment of tourists to hotels.
- b** Check that the probability of all possible assignments equals one.
- c** What is the probability that all four hotels receive guests?
- d** Is the selected rule fair to the four hotels?

Analytical solution to example 8.3

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Let i, j and k be the number of tourists assigned to hotels A, B and C respectively. Then

$$\begin{aligned}\Omega = \{(i, j, k, 15 - (i + j + k)) : 0 \leq i \leq 15, \\ 0 \leq j \leq 15 - i; \\ 0 \leq k \leq 15 - (i + j)\}\end{aligned}$$

- a Define the events $E_j(k)$ = "hotel W receives k tourists," with $W \in \{A, B, C, D\}$. The assignment $(i, j, k, 15 - (i + j + k))$ is the only member of the event $E_A(i) \cap E_B(j) \cap E_C(k) \cap E_D(15 - (i + j + k))$. Then

$$\begin{aligned}P(\{(i, j, k, 15 - (i + j + k))\}) &= P(E_A(i) \cap E_B(j) \cap E_C(k)) \\ &= P(E_C(k) | E_A(i) \cap E_B(j)) P(E_B(j) | E_A(i)) P(E_A(i))\end{aligned}$$

$$P(E_A(i)) = \frac{1}{16}$$

$$P(E_B(j) | E_A(i)) = \frac{1}{16 - i}$$

$$P(E_C(k) | E_B(j) \cap E_A(i)) = \frac{1}{16 - (i + j)}$$

Therefore

$$P(\{(i, j, k, 15 - (i + j + k))\}) = \frac{1}{16 - (i + j)} \frac{1}{16 - i} \frac{1}{16}$$

Analytical solution to example 8.3

Foundations of probability theory

Historical notes
Axiomatic definition of probability
Some basic rules

Conditional probability and Bayes

Conditional probability
Assigning probabilities by conditioning
Independent events
Law of conditional probability
Bayes's rule in odds form
Bayesian inference – discrete case

References

Let i, j and k be the number of tourists assigned to hotels A, B and C respectively. Then

$$\Omega = \{(i, j, k, 15 - (i + j + k)) : 0 \leq i \leq 15, \\ 0 \leq j \leq 15 - i; \\ 0 \leq k \leq 15 - (i + j)\}$$

b

$$\begin{aligned} P(\Omega) &= \sum_{\omega \in \Omega} P(\omega) = \sum_{i=0}^{15} \sum_{j=0}^{15-i} \sum_{k=0}^{15-(i+j)} P(\{(i, j, k, 15 - (i + j + k))\}) \\ &= \sum_{i=0}^{15} \sum_{j=0}^{15-i} \sum_{k=0}^{15-(i+j)} \frac{1}{16 - (i + j)} \frac{1}{16 - i} \frac{1}{16} \\ &= \frac{1}{16} \sum_{i=0}^{15} \frac{1}{16 - i} \sum_{j=0}^{15-i} \frac{1}{16 - (i + j)} \sum_{k=0}^{15-(i+j)} 1 \\ &= \frac{1}{16} \sum_{i=0}^{15} \frac{1}{16 - i} \sum_{j=0}^{15-i} \frac{1}{16 - (i + j)} (16 - (i + j)) \\ &= \frac{1}{16} \sum_{i=0}^{15} \frac{1}{16 - i} \sum_{j=0}^{15-i} 1 = \frac{1}{16} \sum_{i=0}^{15} \frac{1}{16 - i} (16 - i) = \frac{1}{16} \sum_{i=0}^{15} 1 = \frac{1}{16} 16 = 1 \end{aligned}$$

Analytical solution to example 8.3

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References



Define the event $E = \text{"all hotels receive guests."}$ Then

$$E = \{(i, j, k, 15 - (i + j + k)) : 1 \leq i \leq 12, \\ 1 \leq j \leq 13 - i; \\ 1 \leq k \leq 14 - (i + j)\}$$

$$\begin{aligned} P(E) &= \sum_{i=1}^{12} \sum_{j=1}^{13-i} \sum_{k=1}^{14-(i+j)} P(\{(i, j, k, 15 - (i + j + k))\}) \\ &= \sum_{i=1}^{12} \sum_{j=1}^{13-i} \sum_{k=1}^{14-(i+j)} \frac{1}{16 - (i + j)} \frac{1}{16 - i} \frac{1}{16} \\ &= 0.2856 \end{aligned}$$



$$\begin{aligned} P(\{(15, 0, 0, 0)\}) &= \frac{1}{16} \\ P(\{(0, 15, 0, 0)\}) &= \frac{1}{16^2} \end{aligned}$$

Thus, the selected rule is unfair to the four hotels.

Simulated solution to example 8.3

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Please see code [here](#).

Independent events

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- motivation of independence definition with conditional probabilities
- Definition 8.2

Example

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- Example 8.5

Example

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- Example 8.6 (uses birthday problem, example 7.10)

Law of conditional probability

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- example of die followed by coin tosses

Rule 5 law of conditional probability. Let A be an event that can only occur if one of the mutually exclusive events B_1, \dots, B_n occurs. Then

$$P(A) = P(A|B_1)P(B_1) + \dots + P(A|B_n)P(B_n)$$

Example

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- example 8.6: tour the France (difficult!)

Bayes rule in odds form

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- true/false hypothesis

Rule 6 The posterior probability $P(H|E)$ satisfies

$$\frac{P(H|E)}{P(\bar{H}|E)} = \frac{P(H) P(E|H)}{P(\bar{H}) P(E|\bar{H})}$$

- interpretation of rule 6
- avoid need of $P(E)$
- prior odds + likelihood ratio or Bayes factor
- prior odds update with new evidence
- sequential update (mention Bayesian linear regression)

Example

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- example 8.8

Example

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- example 8.11
- add to the problem statement:
- in 1992, 4936 women were murdered in the US, of which roughly 1430 were murdered by their (ex)husbands or boyfriends
- 5% of the married women in the US have at some point been physically abused by their husbands.
- assume that a woman who has been murdered by some other than her husband had the same chance of being abused by her husband as a randomly selected woman
- Alan Dershowitz admitted that a substantial percentage of the husbands who murder their wives, previous to the murder, also physically abuse their wives. Given this statement, we assume that the probability that a husband physically abused his wife, given that he killed her, is 50 percent.

Bayesian inference – discrete case

Foundations of probability theory

- Historical notes
- Axiomatic definition of probability
- Some basic rules

Conditional probability and Bayes

- Conditional probability
- Assigning probabilities by conditioning
- Independent events
- Law of conditional probability
- Baye's rule in odds form
- Bayesian inference – discrete case**

References

- explain posterior sequential update

Example

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

- example 8.13 (solve it analytically and by sampling)

References

Foundations of probability theory

Historical notes

Axiomatic definition of probability

Some basic rules

Conditional probability and Bayes

Conditional probability

Assigning probabilities by conditioning

Independent events

Law of conditional probability

Baye's rule in odds form

Bayesian inference – discrete case

References

Tijms, H. (2012). *Understanding probability*. Cambridge University Press.