

## CHAPTER 6. PANEL DATA

Joan Llull

Quantitative Statistical Methods II  
Barcelona School of Economics

# INTRODUCTION

## *Panel data*

The term **panel data** refers to data sets with **repeated observations** over time (or other dimensions) for a given cross-section of units.

**Units** can be persons, households, firms, countries,...

It is different from **repeated cross-sections**.

Main **advantages** of panel data:

- Permanent unobserved heterogeneity.
- Dynamic responses and error components.

# *Cigarette Consumption*

We will use the same **example** throughout the chapter.

Consider the estimation of the demand for cigarettes:

$$\ln C_{it} = \beta_0 + \beta_1 \ln P_{it} + \beta_2 \ln Y_{it} + \eta_i + v_{it},$$

where:

- $\ln C_{it}$  is log consumption of cigarettes,
- $\ln P_{it}$  is log price of the cigarettes,
- $\ln Y_{it}$  is log income of the individual,
- $\eta_i + v_{it}$  is unobserved.

# STATIC MODELS

# General notation

We consider the following **model**:

$$y_{it} = \mathbf{x}_{it}'\boldsymbol{\beta} + (\eta_i + v_{it}).$$

where  $y_{it}$  and  $\mathbf{x}_{it}$  are **observed**, and  $\eta_i + v_{it}$  is **unobserved**.

Let  $\{y_{it}, \mathbf{x}_{it}\}_{i=1, \dots, N}^{t=1, \dots, T}$  be our **sample**. We define:

$$\mathbf{y}_i \equiv \begin{pmatrix} y_{i1} \\ \vdots \\ y_{iT} \end{pmatrix}, X_i = \begin{pmatrix} \mathbf{x}_{i1}' \\ \vdots \\ \mathbf{x}_{iT}' \end{pmatrix}, \boldsymbol{\eta}_i = \eta_i \boldsymbol{\iota}_T, \text{ and } \mathbf{v}_i = \begin{pmatrix} v_{i1} \\ \vdots \\ v_{iT} \end{pmatrix},$$
$$\mathbf{y} \equiv \begin{pmatrix} \mathbf{y}_1 \\ \vdots \\ \mathbf{y}_N \end{pmatrix}, X = \begin{pmatrix} X_1 \\ \vdots \\ X_N \end{pmatrix}, \boldsymbol{\eta} = \begin{pmatrix} \eta_1 \\ \vdots \\ \eta_N \end{pmatrix}, \text{ and } \mathbf{v} = \begin{pmatrix} \mathbf{v}_1 \\ \vdots \\ \mathbf{v}_N \end{pmatrix},$$

where  $\boldsymbol{\iota}_T$  is a size  $T$  **vector of ones**.

Hence, we can use **compact notation**:  $\mathbf{y}_i = X_i\boldsymbol{\beta} + (\boldsymbol{\eta}_i + \mathbf{v}_i)$ , and  $\mathbf{y} = X\boldsymbol{\beta} + (\boldsymbol{\eta} + \mathbf{v})$ .

## *General assumptions for static models*

For static models, we assume:

- **Fixed effects:**  $\mathbb{E}[\mathbf{x}_{it}\eta_i] \neq 0$  or **random effects:**  $\mathbb{E}[\mathbf{x}_{it}\eta_i] = 0$ .
- **Strict exogeneity:**  $\mathbb{E}[\mathbf{x}_{it}v_{is}] = 0 \ \forall s, t$ . This assumption rules out effects of past  $v_{is}$  on current  $\mathbf{x}_{it}$  (e.g.  $\mathbf{x}_{it}$  cannot include lagged dependent variables).
- **Error components:**  $\mathbb{E}[\eta_i] = \mathbb{E}[v_{it}] = \mathbb{E}[\eta_i v_{it}] = 0$ .
- **Serially uncorrelated shocks:**  $\mathbb{E}[v_{it}v_{is}] = 0 \ \forall s \neq t$ .
- **Homoskedasticity and i.i.d. errors:**  $\eta_i \sim iid(0, \sigma_\eta^2)$  and  $v_{it} \sim iid(0, \sigma_v^2)$ , which does not affect any crucial result, but simplifies some derivations.

## Pooled OLS

A simple approach: define:  $\mathbf{u} \equiv \boldsymbol{\eta} + \mathbf{v}$  and estimate  $\boldsymbol{\beta}$  by OLS:

$$\hat{\boldsymbol{\beta}}_{OLS} = (X'X)^{-1}X'\mathbf{y}.$$

The **properties** of  $\hat{\boldsymbol{\beta}}_{OLS}$  depend on  $\mathbb{E}[\mathbf{x}_{it}\eta_i]$ , as  $\mathbb{E}[\mathbf{x}_{it}v_{it}] = 0$ :

- If  $\mathbb{E}[\mathbf{x}_{it}\eta_i] = 0 \quad \Rightarrow \quad \mathbb{E}[\mathbf{x}_j u_j] = 0$  (**random effects**):  
 $\mathbb{E}[\mathbf{x}_{it}v_{it}] = 0$ 
  - $\hat{\boldsymbol{\beta}}_{OLS}$  is **consistent** as  $N \rightarrow \infty$ , or  $T \rightarrow \infty$ , or both.
  - it is **efficient** only if  $\sigma_{\eta}^2 = 0$ .
- If  $\mathbb{E}[\mathbf{x}_{it}\eta_i] \neq 0 \Rightarrow \mathbb{E}[\mathbf{x}_j u_j] \neq 0$  (**fixed effects**):
  - $\hat{\boldsymbol{\beta}}_{OLS}$  is **inconsistent** as  $N \rightarrow \infty$ , or  $T \rightarrow \infty$ , or both.
  - **cross-section** results are also inconsistent (but panel helps in constructing a consistent alternative).



## *Pooled OLS in Cigarette Demand*

In our previous example, we would **redefine** our regression as:

$$\ln C_j = \beta_0 + \beta_1 \ln P_j + \beta_2 \ln Y_j + u_j,$$

where I used subindex  $j$  to emphasize that each observation *it* is considered as one **independent** observation.

| OLS                             |        |         |
|---------------------------------|--------|---------|
| $\ln \text{ Prices } (\beta_1)$ | -0.083 | (0.015) |
| $\ln \text{ Income } (\beta_2)$ | -0.032 | (0.006) |

Potential problems:

- preferences,
- education,
- parental smoking behavior,...

**The fixed effects model.  
Within groups estimation.**

## *Within groups estimator*

Write the model in **deviations from individual means**,  $\tilde{y}_{it} \equiv y_{it} - \bar{y}_i$ , where  $\bar{y}_i \equiv T^{-1} \sum_{t=1}^T y_{it}$ :

$$\tilde{y}_{it} = (\mathbf{x}_{it} - \bar{\mathbf{x}}_i)' \boldsymbol{\beta} + (\eta_i - \bar{\eta}_i) + (v_{it} - \bar{v}_i) = \tilde{\mathbf{x}}_{it}' \boldsymbol{\beta} + \tilde{v}_{it}.$$

Given the previous **assumptions**:

$$\mathbb{E}[\tilde{\mathbf{x}}_{it} \tilde{v}_{it}] = 0.$$

Therefore, **OLS on the transformed model**:

$$\hat{\boldsymbol{\beta}}_{WG} = \left( \tilde{X}' \tilde{X} \right)^{-1} \tilde{X}' \tilde{y},$$

is a **consistent** estimator either if  $\mathbb{E}[\mathbf{x}_{it} \eta_i] \neq 0$  or  $\mathbb{E}[\mathbf{x}_{it} \eta_i] = 0$ .

**Strict exogeneity** is a crucial assumption (see next slide).

## *The role of strict exogeneity*

In the case where  $N \rightarrow \infty$  and  $T$  is fixed, consistency depends on **strict exogeneity**.

To see it, recall that:

$$\tilde{\mathbf{x}}_{it} = \mathbf{x}_{it} - \frac{1}{T}(\mathbf{x}_{i1} + \dots + \mathbf{x}_{iT}) \text{ and } \tilde{v}_{it} = v_{it} - \frac{1}{T}(v_{i1} + \dots + v_{iT}).$$

Therefore  $\mathbb{E}[\tilde{\mathbf{x}}_{it}\tilde{v}_{it}] = 0$  requires  $\mathbb{E}[\mathbf{x}_{it}v_{is}] = 0 \ \forall \ s, t$  unless  $T \rightarrow \infty$ .

This has motivated the development of **dynamic panel data models**, to relax this assumption.

## *Pros and cons of within groups estimator*

**Advantage:** consistent either if  $\mathbb{E}[\mathbf{x}_{it}\eta_i] \neq 0$  or  $\mathbb{E}[\mathbf{x}_{it}\eta_i] = 0$ .

Limitations:

- **Not efficient:**

- When  $N \rightarrow \infty$  but  $T$  is fixed, less efficient than e.g.  $\hat{\beta}_{GLS}$  if  $\mathbb{E}[\mathbf{x}_{it}\eta_i] = 0$ .
- Even if  $\mathbb{E}[\mathbf{x}_{it}\eta_i] \neq 0$ , differencing introduces correlation in the errors.

- It does not allow to identify coefficients for **time-invariant regressors**, and identification is through **switchers**.

## *Within Groups in Cigarette Demand*

In our example:

|                         | OLS    |         | WG     |         |
|-------------------------|--------|---------|--------|---------|
| ln Prices ( $\beta_1$ ) | -0.083 | (0.015) | -0.292 | (0.023) |
| ln Income ( $\beta_2$ ) | -0.032 | (0.006) | 0.107  | (0.019) |

## *Least Squares Dummy Variables*

The Within Groups estimator can also be obtained by including a set of  $N$  individual **dummy variables**:

$$y_{it} = \mathbf{x}'_{it}\boldsymbol{\beta} + \eta_1 D_{1i} + \dots + \eta_N D_{Ni} + v_{it},$$

where  $D_{hi} = \mathbb{1}\{h = i\}$  (e.g.  $D_{1i}$  takes the value of 1 for the observations on individual 1 and 0 for all other observations).

OLS estimation of this model gives **numerically equivalent** estimates to WG (that's why  $\hat{\boldsymbol{\beta}}_{WG}$  is a.k.a.  $\hat{\boldsymbol{\beta}}_{LSDV}$ ).

## *LSDV in Cigarette Demand*

We can generate individual (firm) dummies and estimate by OLS, to check that it delivers the same results:

|                                     | OLS               | WG                | LSDV              |
|-------------------------------------|-------------------|-------------------|-------------------|
| ln Prices ( $\beta_1$ )             | -0.083<br>(0.015) | -0.292<br>(0.023) | -0.292<br>(0.023) |
| ln Income ( $\beta_2$ )             | -0.032<br>(0.006) | 0.107<br>(0.019)  | 0.107<br>(0.019)  |
| Individual 1 ( $\beta_0 + \eta_1$ ) |                   |                   | 2.804<br>(0.288)  |
| Individual 2 ( $\beta_0 + \eta_2$ ) |                   |                   | 3.455<br>(0.398)  |
| Individual 3 ( $\beta_0 + \eta_3$ ) |                   |                   | 2.891<br>(0.416)  |
| Individual 4 ( $\beta_0 + \eta_4$ ) |                   |                   | 2.908<br>(0.384)  |
| Individual 5 ( $\beta_0 + \eta_5$ ) |                   |                   | 3.490<br>(0.433)  |
| Individual 6 ( $\beta_0 + \eta_6$ ) |                   |                   | 2.092             |



## *First-Differenced Least Squares*

Another transformation we can consider is **first differences**:

$$\Delta y_{it} = \Delta \mathbf{x}_{it}' \boldsymbol{\beta} + \Delta v_{it}, \text{ for } i = 1, \dots, N; t = 2, \dots, T$$

where  $\Delta y_{it} = y_{it} - y_{it-1}$ .

**Takes out time-invariant** individual effects ( $\Delta \eta_i = \eta_i - \eta_i = 0$ ), so OLS on the differenced model is **consistent**.

**Consistency** requires  $\mathbb{E}[\Delta \mathbf{x}_{it} \Delta v_{it}] = 0$  which is implied by but weaker than strict exogeneity.

**WG more efficient** than FDLS under **classical assumptions**.

**FDLS more efficient** if  $v_{it}$  random walk ( $\Delta v_{it} = \varepsilon_{it} \sim iid(0, \sigma_\varepsilon^2)$ ).

## *FDLS in Cigarette Demand*

To get FDLS we generate first differences and estimate by OLS:

|                                     | OLS               | WG                | LSDV              | FDLS              |
|-------------------------------------|-------------------|-------------------|-------------------|-------------------|
| ln Prices ( $\beta_1$ )             | -0.083<br>(0.015) | -0.292<br>(0.023) | -0.292<br>(0.023) | -0.413<br>(0.035) |
| ln Income ( $\beta_2$ )             | -0.032<br>(0.006) | 0.107<br>(0.019)  | 0.107<br>(0.019)  | 0.178<br>(0.055)  |
| Individual 1 ( $\beta_0 + \eta_1$ ) |                   |                   | 2.804<br>(0.288)  |                   |
| Individual 2 ( $\beta_0 + \eta_2$ ) |                   |                   | 3.455<br>(0.398)  |                   |
| Individual 3 ( $\beta_0 + \eta_3$ ) |                   |                   | 2.891<br>(0.416)  |                   |
| Individual 4 ( $\beta_0 + \eta_4$ ) |                   |                   | 2.908<br>(0.384)  |                   |
| ...                                 |                   |                   | ...               |                   |

**The random effects model.  
Error components.**

## *Uncorrelated effects*

Now we assume uncorrelated or **random effects**:  $\mathbb{E}[\mathbf{x}_{it}\eta_i] = 0$ .

In this case, OLS is **consistent**, but **not efficient**.

The inefficiency is provided by the **serial correlation** induced by  $\eta_i$ :

$$\mathbb{E}[u_{it}u_{is}] = \mathbb{E}[(\eta_i + v_{it})(\eta_i + v_{is})] = \mathbb{E}[\eta_i^2] = \sigma_\eta^2.$$

The **variance** of the unobservables (under classical assumptions) is:

$$\mathbb{E}[u_{it}^2] = \mathbb{E}[\eta_i^2] + \mathbb{E}[v_{it}^2] = \sigma_\eta^2 + \sigma_v^2.$$

## *Error structure*

Therefore, the variance-covariance matrix of the unobservables is:

$$\mathbb{E}[\mathbf{u}_i \mathbf{u}_i'] = \begin{pmatrix} \sigma_\eta^2 + \sigma_v^2 & \sigma_\eta^2 & \dots & \sigma_\eta^2 \\ \sigma_\eta^2 & \sigma_\eta^2 + \sigma_v^2 & \dots & \sigma_\eta^2 \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_\eta^2 & \sigma_\eta^2 & \dots & \sigma_\eta^2 + \sigma_v^2 \end{pmatrix} = \Omega_i,$$

whose dimensions are  $T \times T$ , and  $\mathbb{E}[\mathbf{u}_i \mathbf{u}_h'] = 0 \ \forall i \neq h$ , or:

$$\mathbb{E}[\mathbf{u} \mathbf{u}'] = \begin{pmatrix} \Omega_1 & 0 & \dots & 0 \\ 0 & \Omega_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \Omega_N \end{pmatrix} = \Omega,$$

which is block-diagonal with dimension  $NT \times NT$ .

## *Generalized Least Squares*

Under the classical assumptions, GLS (Balestra-Nerlove) estimator is **consistent and efficient** if  $\mathbb{E}[\mathbf{x}_{it}\eta_i] = 0$ :

$$\hat{\beta}_{GLS} = (X'\Omega^{-1}X)^{-1} X'\Omega^{-1}\mathbf{y}.$$

If  $\mathbb{E}[\mathbf{x}_{it}\eta_i] \neq 0$  GLS is **inconsistent** as  $N \rightarrow \infty$  and  $T$  is fixed.

This estimator is **unfeasible** because we do not know  $\sigma_\eta^2$  and  $\sigma_v^2$ .

## *Theta-differencing*

$\hat{\beta}_{GLS}$  is **equivalent** to OLS on the theta-differenced model:

$$y_{it}^* = \mathbf{x}_{it}^{*'} \boldsymbol{\beta} + u_{it}^*,$$

where:

$$y_{it}^* = y_{it} - (1 - \theta)\bar{y}_i,$$

and:

$$\theta^2 = \frac{\sigma_v^2}{\sigma_v^2 + T\sigma_\eta^2}.$$

**Consistency** relies on  $\mathbb{E}[\mathbf{x}_{it}\eta_i] = 0$  (as  $\eta_i$  not eliminated).

**Two special cases:**

- If  $\sigma_\eta^2 = 0$  (i.e. no individual effect), OLS is efficient.
- If  $T \rightarrow \infty$ , then  $\theta \rightarrow 0$ , and  $y_{it}^* \rightarrow \tilde{y}_{it} = y_{it} - \bar{y}_i$ : WG is efficient.

## *Feasible GLS*

$\hat{\beta}_{GLS}$  is **unfeasible** because we do not know  $\sigma_\eta^2$  and  $\sigma_v^2$ .

A consistent estimator of  $\sigma_v^2$  is provided by the **WG residuals**:

$$\hat{v}_{it} \equiv \tilde{y}_{it} - \tilde{\mathbf{x}}'_{it} \hat{\beta}_{WG}$$

$$\hat{\sigma}_v^2 = \frac{\hat{\mathbf{v}}' \hat{\mathbf{v}}}{N(T-1) - K}$$

Then, a consistent estimator of  $\sigma_\eta^2$  is given by the **BG residuals**:

$$\bar{y}_i = \bar{\mathbf{x}}'_i \boldsymbol{\beta} + \bar{\eta}_i + \bar{v}_i, \quad i = 1, \dots, N \Rightarrow \hat{\beta}_{BG}$$

$$\hat{u}_i \equiv \bar{y}_i - \bar{\mathbf{x}}'_i \hat{\beta}_{BG}$$

$$\hat{\sigma}_u^2 = \left( \widehat{\sigma_\eta^2 + \frac{1}{T} \sigma_v^2} \right) = \frac{\hat{\mathbf{u}}' \hat{\mathbf{u}}}{N - K} \quad \Rightarrow \quad \hat{\sigma}_\eta^2 = \hat{\sigma}_u^2 - \frac{1}{T} \hat{\sigma}_v^2.$$



## *Feasible GLS in Cigarette Demand*

In our example, if we now estimate  $\hat{\beta}_{FGLS}$ , we get:

|                         | OLS               | WG                | FGLS              |
|-------------------------|-------------------|-------------------|-------------------|
| ln Prices ( $\beta_1$ ) | -0.083<br>(0.015) | -0.292<br>(0.023) | -0.122<br>(0.014) |
| ln Income ( $\beta_2$ ) | -0.032<br>(0.006) | 0.107<br>(0.019)  | -0.012<br>(0.004) |

**Testing for correlated individual effects.**

## *Testing for correlated effects (Hausman test)*

$\hat{\beta}_{WG}$  is **consistent** regardless of  $\mathbb{E}[\mathbf{x}_{it}\eta_i]$  being equal to zero or not.

$\hat{\beta}_{FGLS}$  is **consistent only** if  $\mathbb{E}[\mathbf{x}_{it}\eta_i] = 0$ .

$\Rightarrow$  we can test whether both **estimates are similar!**

The **Hausman test** does exactly this comparison:

$$h = \hat{\mathbf{q}}' [\text{avar}(\hat{\mathbf{q}})]^{-1} \hat{\mathbf{q}} \stackrel{a}{\sim} \chi^2(K),$$

under the **null hypothesis**  $\mathbb{E}[\mathbf{x}_{it}\eta_i] = 0$ , where:

$$\hat{\mathbf{q}} = \hat{\beta}_{WG} - \hat{\beta}_{FGLS},$$

and:

$$\text{avar}(\hat{\mathbf{q}}) = \text{avar}(\hat{\beta}_{WG}) - \text{avar}(\hat{\beta}_{FGLS}).$$

Requires **classical assumptions** (FGLS to be more efficient than WG).

## *Hausman test in Cigarette Demand*

Output from software packages often includes the Hausman test. In our example:

|              | Statistic | P-value |
|--------------|-----------|---------|
| Hausman test | 24.661    | 0.000   |

# DYNAMIC MODELS

# Autoregressive models with individual effects

# *Autoregressive panel data model*

We consider the following model:

$$y_{it} = \alpha y_{it-1} + \eta_i + v_{it} \quad |\alpha| < 1.$$

**Other regressors** can be included, but main results unaffected.

We assume:

- **Error components:**  $\mathbb{E}[\eta_i] = \mathbb{E}[v_{it}] = \mathbb{E}[\eta_i v_{it}] = 0.$
- **Serially uncorrelated shocks:**  $\mathbb{E}[v_{it} v_{is}] = 0 \ \forall \ s \neq t.$
- **Predetermined initial cond.:**  $\mathbb{E}[y_{i0} v_{it}] = 0 \ \forall \ t = 1, \dots, T.$

# *Properties of pooled OLS and WG estimators*

Even assuming  $\mathbb{E}[y_{it-1}v_{it}] = 0$ , still **OLS** delivers:

$$\text{plim}_{N \rightarrow \infty} \hat{\alpha}_{OLS} > \alpha,$$

$$\text{because } \mathbb{E}[y_{it-1}\eta_i] = \sigma_\eta^2 \left( \frac{1-\alpha^{t-1}}{1-\alpha} \right) + \alpha^{t-1} \mathbb{E}[y_{i0}\eta_i] > 0.$$

Doing the **within groups** transformation we see that:

$$\text{plim}_{N \rightarrow \infty} \hat{\alpha}_{WG} < \alpha,$$

$$\text{because } \mathbb{E}[\tilde{y}_{it-1}\tilde{v}_{it}] = -A\sigma_v^2 < 0. \quad \left( A = \frac{(1-\alpha)(1+T(1-\alpha^{t-1}-\alpha^{T-1-t})) + \alpha^T(1-\alpha^{T-1})}{T^2(1-\alpha)^2} \right)$$

**WG bias vanishes** as  $T \rightarrow \infty$  (bias not small even with  $T = 15$ ).

Supposedly consistent estimators with  $\hat{\alpha} \gg \alpha_{OLS}$  or  $\hat{\alpha} \ll \hat{\alpha}_{WG}$  should be **seen with suspicion**.



# Anderson and Hsiao

Consider the model in **first differences**:

$$\Delta y_{it} = \alpha \Delta y_{it-1} + \Delta v_{it}.$$

OLS in first differences is **inconsistent**:  $\mathbb{E}[\Delta y_{it-1} \Delta v_{it}] = -\sigma_v^2 < 0$ .

However,  $y_{it-2}$  or  $\Delta y_{it-2}$  are valid **instruments** for  $\Delta y_{it-1}$ :

- **Relevance**:  $\mathbb{E}[\Delta y_{it-2} \Delta y_{it-1}] \neq 0$ ,  $\mathbb{E}[y_{it-2} \Delta y_{it-1}] \neq 0$ .
- **Orthogonality**:  $\mathbb{E}[\Delta y_{it-2} \Delta v_{it}] = \mathbb{E}[y_{it-2} \Delta v_{it}] = 0$ .

Anderson and Hsiao (1981) proposed this **2SLS** estimators:

$$\hat{\alpha}_{AH} = \left( \widehat{\Delta \mathbf{y}'_{-1} \Delta \mathbf{y}_{-1}} \right)^{-1} \widehat{\Delta \mathbf{y}'_{-1} \Delta \mathbf{y}},$$

where:

$$\widehat{\Delta \mathbf{y}_{-1}} = Z (Z' Z)^{-1} Z' \Delta \mathbf{y}_{-1},$$

where  $Z$  can be  $\mathbf{y}_{-2}$  or  $\Delta \mathbf{y}_{-2}$ .

Requires min. **three periods** ( $T = 2$  and  $y_{i0}$ ). Only **efficient** if  $T = 2$ .

## *AR(1) Cigarette Demand (No Covariates)*

In our example, we redefine the model to be an AR(1) process (for now without regressors):

$$\ln C_{it} = \alpha \ln C_{it-1} + \eta_i + v_{it}.$$

The Anderson-Hsiao results (together with OLS and WG) are:

|                                 | OLS              | WG               | Anderson-Hsiao   |
|---------------------------------|------------------|------------------|------------------|
| Lagged cons. ( $\ln C_{it-1}$ ) | 0.982<br>(0.003) | 0.884<br>(0.061) | 1.395<br>(0.090) |

## Differenced GMM estimation.

## *GMM in 3 slides (I): the setup*

GMM finds parameter estimates that come as close as possible to satisfy **orthogonality conditions** (or moment conditions) in the sample.

E.g., consider  $K$  regressors  $\mathbf{x}_i$  and  $L$  “instruments”  $\mathbf{z}_i$ :

$$u_i = y_i - f(\mathbf{x}_i, \boldsymbol{\beta}) \quad \mathbf{z}_i = g(\mathbf{x}_i).$$

The model specifies  $L$  **moment conditions**:  $\mathbb{E}[\mathbf{z}_i u_i] = \mathbf{0}$ .

**Sample analogue:**

$$\mathbf{b}_N(\boldsymbol{\beta}) = \frac{1}{N} \sum_{i=1}^N \mathbf{z}_i u_i(\boldsymbol{\beta}).$$

## *GMM in 3 slides (II): the estimation*

Two possible cases **cases**:

- $L = K$  (**just identified**):  $\mathbf{b}_N(\hat{\beta}_{GMM}) = \mathbf{0}$ .
- $L > K$  (**overidentified**):  $\hat{\beta}_{GMM} = \arg \min_{\beta} \mathbf{b}_N(\beta)' W_N \mathbf{b}_N(\beta)$ .

$W_N$  is a **positive definite** weighting matrix.

**Optimal GMM** (efficient) uses  $\left( \frac{1}{N} \sum_{i=1}^N \mathbb{E}[z_i u_i u_i' z_i'] \right)^{-1}$ , the inverse of the variance-covariance matrix, as weighting matrix.

## *GMM in 3 slides (III): asymptotic properties*

$\hat{\beta}_{GMM}$  is a **consistent** estimator of  $\beta$ .

It is **asymptotically normal**, with the following variance:

$$avar(\hat{\beta}_{GMM}) = (D'WD)^{-1}D'WS_0WD(D'WD)^{-1},$$

where:

$$D \equiv \text{plim}_{N \rightarrow \infty} \frac{\partial \mathbf{b}_N(\beta)}{\partial \beta'},$$

$$W \equiv \text{plim}_{N \rightarrow \infty} W_N,$$

$$S_0 \equiv \frac{1}{N} \sum_{i=1}^N \mathbb{E}[z_i u_i u_i' z_i'].$$

# Moment conditions

Given previous assumptions, several **moment conditions**:

| Equation   | Instruments                               | Orthogonality cond.   |
|--|---|---|
| $\Delta y_{i2} = \alpha \Delta y_{i1} + \Delta v_{i2}$   | $y_{i0}$                                  | $\mathbb{E}[\Delta v_{i2} y_{i0}] = 0$  |
| $\Delta y_{i3} = \alpha \Delta y_{i2} + \Delta v_{i3}$   | $y_{i0}, y_{i1}$                          | $\mathbb{E} \left[ \Delta v_{i3} \begin{pmatrix} y_{i0} \\ y_{i1} \end{pmatrix} \right] = \mathbf{0}$                                 |
| $\Delta y_{i4} = \alpha \Delta y_{i3} + \Delta v_{i4}$   | $y_{i0}, y_{i1}, y_{i2}$                  | $\mathbb{E} \left[ \Delta v_{i4} \begin{pmatrix} y_{i0} \\ y_{i1} \\ y_{i2} \end{pmatrix} \right] = \mathbf{0}$                       |
| $\vdots$   | $\vdots$                                  | $\vdots$  |
| $\Delta y_{iT} = \alpha \Delta y_{iT-1} + \Delta v_{iT}$ | $y_{i0}, y_{i1}, y_{i2}, \dots, y_{iT-2}$ | $\mathbb{E} \left[ \Delta v_{iT} \begin{pmatrix} y_{i0} \\ y_{i1} \\ y_{i2} \\ \vdots \\ y_{iT-2} \end{pmatrix} \right] = \mathbf{0}$ |

We end up with  $(T - 1)T/2$  **moment conditions**.

## Moment conditions in matrix form

We can write these **moment conditions** as  $\mathbb{E}[Z_i' \Delta \mathbf{v}_i] = 0$ , where:

$$Z_i = \begin{pmatrix} y_{i0} & 0 & 0 & 0 & \dots & 0 & 0 & \dots & 0 \\ 0 & y_{i0} & y_{i1} & 0 & \dots & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & 0 & \dots & y_{i0} & y_{i1} & \dots & y_{iT-2} \end{pmatrix} \text{ and } \Delta \mathbf{v}_i = \begin{pmatrix} \Delta v_{i2} \\ \Delta v_{i3} \\ \vdots \\ \Delta v_{iT} \end{pmatrix},$$

and the **sample analogue** is:

$$\mathbf{b}_N(\alpha) = \frac{1}{N} \sum_{i=1}^N Z_i' \Delta \mathbf{v}_i(\alpha).$$

Hence, the **GMM estimator** (proposed by Arellano and Bond, 1991) is:

$$\begin{aligned} \hat{\alpha}_{GMM} &= \arg \min_{\alpha} \left( \frac{1}{N} \sum_{i=1}^N \Delta \mathbf{v}_i'(\alpha) Z_i \right) W_N \left( \frac{1}{N} \sum_{i=1}^N Z_i' \Delta \mathbf{v}_i(\alpha) \right) = \\ &= (\Delta \mathbf{y}_{-1}' Z W_N Z' \Delta \mathbf{y}_{-1})^{-1} \Delta \mathbf{y}_{-1}' Z W_N Z' \Delta \mathbf{y}. \end{aligned}$$



## Optimal weighting matrix

The **optimal weighting matrix** (efficient GMM) is:

$$W_N = \left( \frac{1}{N} \sum_{i=1}^N \mathbb{E}[Z_i' \Delta \mathbf{v}_i \Delta \mathbf{v}_i' Z_i] \right)^{-1}.$$

The **sample analogue** is obtained in a **two-step** procedure:

$$W_N = \left( \frac{1}{N} \sum_{i=1}^N [Z_i' \widehat{\Delta \mathbf{v}_i(\hat{\alpha})} \widehat{\Delta \mathbf{v}_i'(\hat{\alpha})} Z_i] \right)^{-1}.$$

Windeijer (2005) proposes a **finite sample correction** of the variance that accounts for  $\alpha$  being estimated.

The most common **one-step** (and first-step) matrix uses the structure of  $\mathbb{E}[\Delta \mathbf{v}_i \Delta \mathbf{v}_i']$ :

$$\mathbb{E}[\Delta \mathbf{v}_i \Delta \mathbf{v}_i'] = \sigma_v^2 \begin{pmatrix} 2 & -1 & 0 & 0 & \dots & 0 \\ -1 & 2 & -1 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \\ 0 & 0 & 0 & 0 & \dots & 2 \end{pmatrix}.$$

## *GMM in Cigarette Demand*

GMM results in our example are:

|                           | Coefficient | Standard Error |
|---------------------------|-------------|----------------|
| Least Squares (OLS)       | 0.982       | (0.003)        |
| Within Groups (WG)        | 0.884       | (0.061)        |
| Anderson-Hsiao            | 1.395       | (0.090)        |
| One-step GMM              | 1.023       | (0.104)        |
| Two-step GMM              | 0.994       | (0.040)        |
| Two-step GMM small sample | 0.994       | (0.121)        |

# *Potential limitations of Arellano-Bond*

## Weak instruments:

- When  $\alpha \rightarrow 1$ , **relevance** of the instrument decreases.
- Instruments are still valid, but have **poor small sample** properties.
- **Monte Carlo evidence** shows that with  $\alpha > 0.8$ , estimator behaves poorly unless huge samples available.
- There are **alternatives** in the literature.

## Overfitting:

- “**Too many**” instruments if  $T$  relative to  $N$  is relatively large.
- We might want to **restrict** the number of instruments used.
- It is always good practice to check **robustness** to different combinations of instruments.

## *Dynamic linear model*

Once we include **regressors**, the model is:

$$y_{it} = \alpha y_{it-1} + \mathbf{x}_{it}'\boldsymbol{\beta} + \eta_i + v_{it} \quad |\alpha| < 1.$$

We maintain the **previous assumptions**: error components, serially uncorrelated shocks, and predetermined initial conditions.

Therefore, moment conditions of the kind:

$$\mathbb{E}[y_{it-s}\Delta v_{it}] = 0 \quad t = 2, \dots, T, s \geq 2,$$

are **still valid**.

## *Assumptions on regressors*

Different assumptions regarding  $\mathbf{x}_{it}$  will enable different additional **orthogonality conditions**:

- $\mathbf{x}_{it}$  can be correlated or uncorrelated with  $\eta_i$ .
- $\mathbf{x}_{it}$  can be endogenous, predetermined, or strictly exogenous with respect to  $v_{it}$ .

For instance, if assumptions are analogous to those for  $y_{it-1}$ , we may use  $\mathbf{x}_{it-1}$  (and longer lags) as instruments:

$$Z_i = \begin{pmatrix} y_{i0} & \mathbf{x}'_{i0} & \mathbf{x}'_{i1} & \dots & 0 & \dots & 0 & \mathbf{0}' & \dots & \mathbf{0}' \\ \vdots & \vdots & \vdots & \ddots & \vdots & \dots & \vdots & \vdots & \vdots & \vdots \\ 0 & \mathbf{0}' & \mathbf{0}' & \dots & y_{i0} & \dots & y_{iT-2} & \mathbf{x}'_{i0} & \dots & \mathbf{x}'_{iT-1} \end{pmatrix}.$$

## *Dynamic Cigarette Demand with Regressors*

In our example, we rewrite the employment equations as follows:

$$\ln C_{it} = \alpha \ln C_{it-1} + \beta_1 \ln P_{it} + \beta_2 \ln Y_{it} + \eta_i + v_{it}.$$

Results are:

|                         | OLS              | WG                | GMM               |
|-------------------------|------------------|-------------------|-------------------|
| Lagged dep ( $\alpha$ ) | 0.947<br>(0.011) | 0.528<br>(0.064)  | 0.495<br>(0.127)  |
| ln Prices ( $\beta_1$ ) | 0.010<br>(0.006) | -0.501<br>(0.098) | -0.607<br>(0.143) |
| ln Income ( $\beta_2$ ) | 0.049<br>(0.011) | 0.369<br>(0.044)  | 0.338<br>(0.051)  |

# System GMM estimation

## *Additional orthogonality conditions*

Recall our  $(T - 1)T/2$  **moment conditions**:

$$\mathbb{E}[y_{it-s}\Delta v_{it}] = 0 \quad t = 2, \dots, T; s \geq 2.$$

**System GMM** (Arellano and Bover, 1995) uses the assumption  $\mathbb{E}[y_{i0}|\eta_i] = \frac{\eta_i}{1-\alpha}$ :

$$\mathbb{E}[\Delta y_{it}\eta_i] = 0, \quad \forall t$$

or, alternatively:

$$\mathbb{E}[\Delta y_{iT-s}u_{iT}] = 0, \quad u_{iT} = \eta_i + v_{iT}, \quad s = 1, \dots, T - 1.$$



## *The System GMM estimator*

**Analogously** to the first-differenced GMM, the estimator is given by  $\mathbb{E}[(Z^*)'u_i^*] = 0$ :

$$\hat{\alpha}_{Sys-GMM} = ((X^*)'Z^*W_N(Z^*)'X^*)^{-1} X^*Z^*W_N(Z^*)'\mathbf{y}^*,$$

where:

$$Z_i^* = \begin{pmatrix} Z_i & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0}' & \Delta y_{i1} & \dots & \Delta y_{iT-1} \end{pmatrix}, \mathbf{u}_i^* = \begin{pmatrix} \Delta \mathbf{v}_i \\ \eta_i + v_{iT} \end{pmatrix}, X_i^* = \begin{pmatrix} \Delta \mathbf{y}_{-1i} \\ y_{iT-1} \end{pmatrix} \text{ and } \mathbf{y}_i^* = \begin{pmatrix} \Delta \mathbf{y}_i \\ y_{iT} \end{pmatrix},$$

This estimator is **more efficient**, as it uses additional moment conditions.

It **reduces small sample bias**, especially when  $\alpha \rightarrow 1$ .

## *System-GMM in Cigarette Demand*

GMM results in our example are:

|                           | Coefficient | Standard Error |
|---------------------------|-------------|----------------|
| Least Squares (OLS)       | 0.982       | (0.003)        |
| Within Groups (WG)        | 0.884       | (0.061)        |
| Anderson-Hsiao            | 1.395       | (0.090)        |
| One-step GMM              | 1.023       | (0.104)        |
| Two-step GMM              | 0.994       | (0.040)        |
| Two-step GMM small sample | 0.994       | (0.121)        |
| One-step System-GMM       | 0.926       | (0.023)        |
| Two-step System-GMM small | 0.911       | (0.032)        |

# Specification tests

## *Specification tests*

There are several relevant aspects for the validity of the estimation that can be **tested formally**.

- **Orthogonality conditions:** are they small enough to not reject that they are zero (overidentifying restrictions).
- **Validity** of a subset of more restrictive assumptions (difference Sargan test, Hausman test).
- **Serial correlation in the data:** vital for the validity of the instruments (Arellano-Bond test).

## *Sargan/Hansen overidentifying restrictions test*

The null hypothesis is whether the **orthogonality** conditions are **satisfied** (i.e. moments are equal to zero).

The test can only be implemented if the problem is **overidentified** (otherwise the sample moments are exactly zero by construction).

The **test** is:

$$S = NJ_N(\beta) = N \left( \frac{1}{N} \sum_{i=1}^N \hat{\mathbf{u}}_i' Z_i \left( \frac{1}{N} \sum_{i=1}^N Z_i' \hat{\mathbf{u}}_i \hat{\mathbf{u}}_i' Z_i \right)^{-1} \frac{1}{N} \sum_{i=1}^N Z_i' \hat{\mathbf{u}}_i \right),$$

where  $\hat{\mathbf{u}}$  are those predicted residuals from the first step and  $\hat{\hat{\mathbf{u}}}$  are predicted from the second stage, and:

$$S \stackrel{a}{\sim} \chi^2(L - K).$$

## *Testing stronger assumptions*

Some of the assumptions that we make are **stronger** than others.

If the problem is overidentified, we can **test** whether results change if we **include or exclude** the orthogonality conditions generated by them.

If they are true, increase **efficiency**, but if not, **inconsistent**!

**Two** ways of testing it:

- **Overidentifying restrictions** (difference in Sargan should be close to zero:  $\Delta S = S - S^A \stackrel{a}{\sim} \chi^2(L - L^A)$ ).
- **Differences in coefficients** (Hausman test:  $\hat{q} = \hat{\delta}_{GMM}^A - \hat{\delta}_{GMM}$ ).

## *Direct test for serial correlation*

The test was proposed by Arellano-Bond (1991).

Tests for the presence of **second order autocorrelation** in the first-differenced residuals.

If differences in residuals are second-order correlated, some **instruments would not be valid!**

The **test** is:

$$m_2 = \frac{\widehat{\Delta \mathbf{v}_{-2}}' \widehat{\Delta \mathbf{v}_*}}{se} \stackrel{a}{\sim} \mathcal{N}(0, 1),$$

where  $\Delta \mathbf{v}_{-2}$  is the second lagged residual in differences, and  $\Delta \mathbf{v}_*$  is the part of the vector of contemporaneous first differences for the periods that overlap with the second lagged vector.

Values close to zero do not reject the hypothesis of **no serial correlation**.