

Note

Joanne Yu

May 4, 2026

1 February 26

1.1 Sample Preparation [1]

1.1.1 For $W_L^\pm W_L^\pm$

```
generate p p > w+{0} w+{0} j j QCD=0, w+ > l+ vl
add process p p > w-{0} w-{0} j j QCD=0, w- > l- vl~
output WW_LL
launch
shower = pythia8
detector = Delphes (R=0.4)
```

1.1.2 For $W_L^\pm W_T^\pm$

```
generate p p > w+{0} w+{T} j j QCD=0, w+ > l+ vl
add process p p > w-{0} w-{T} j j QCD=0, w- > l- vl~
output WW_LT
launch
shower = pythia8
detector = Delphes (R=0.4)
```

1.1.3 For $W_T^\pm W_T^\pm$

```
generate p p > w+{T} w+{T} j j QCD=0, w+ > l+ vl
add process p p > w-{T} w-{T} j j QCD=0, w- > l- vl~
output WW_TT
launch
shower = pythia8
detector = Delphes (R=0.4)
```

1.1.4 For Unpolarized $W^\pm W^\pm$

```
generate p p > w+ w+ j j QCD=0, w+ > l+ vl
add process p p > w- w- j j QCD=0, w- > l- vl~
output WW_UN
launch
shower = pythia8
detector = Delphes (R=0.4)
```

1.2 Restrictions for Candidates

1.2.1 Electrons

Electrons are required to satisfy the “Tight” likelihood-based identification criteria.

- $p_T > 27$ GeV
- $|\eta| < 2.47$ GeV
- Excluding the calorimeter transition region $1.37 < |\eta| < 1.52$

```
ele_mask = (eles.pt > 27) & (abs(eles.eta) < 2.47) & ~((abs(eles.eta) > 1.37) &
(abs(eles.eta) < 1.52))
```

1.2.2 Muons

Muons are required to satisfy the “medium” identification selection.

- $p_T > 27$ GeV
- $|\eta| < 2.5$

```
mu_mask = (muons.pt > 27) & (abs(muons.eta) < 2.5)
```

1.2.3 Jets

Jets are reconstructed using the anti- k_t algorithm with a radius parameter of $R = 0.4$, using the particle-flow object.

- At least two jets with $|\eta| < 4.5$ are required to be present in each event
- $p_T > 65$ GeV for leading jets (written in the paragraph for election of the leading leptons later.)
- $p_T > 35$ GeV for subleading jets

```
jet_mask = (jets.pt > 35) & (abs(jets.eta) < 4.5)
```

1.3 Selection Cuts

The event selection in the signal region (SR) requires:

1. A same-sign lepton pair with $m_{\ell\ell} > 20$ GeV

```
padded_leps = ak.pad_none(all_leps, 2)
l1, l2 = padded_leps[:, 0], padded_leps[:, 1]
```

```
mask1 = (ak.num(all_leps) >= 2) & ak.fill_none((l1.q == l2.q) & ((l1 + l2).mass
> 20), False)
n1 = ak.sum(mask1)
```

2. $E_T^{\text{miss}} > 30$ GeV

```
met_pt = ak.firsts(data.MissingET.MET)
mask2 = mask1 & ak.fill_none((met_pt > 30), False)
n2 = ak.sum(mask2)
```

3. VBS topology: The two highest- p_T jets must satisfy $m_{jj} > 500$ GeV with $|\Delta y_{jj}| > 2.0$

```
padded_jets = ak.pad_none(sel_jets, 2)
j1, j2 = padded_jets[:, 0], padded_jets[:, 1]

vbs_kinematics = (j1.pt > 65) & (j2.pt > 35) & ((j1 + j2).mass > 500) & (abs(j1.
rapidity - j2.rapidity) > 2.0)
mask3 = mask2 & (ak.num(sel_jets) >= 2) & ak.fill_none(vbs_kinematics, False)
n3 = ak.sum(mask3)
```

Cut	LL	pass rate	LT	pass rate	TT	pass rate	Unpol.	pass rate
Total	50000	1.00	50000	1.00	50000	1.00	50000	1.00
Lepton cut	12534	0.25	12453	0.25	12814	0.26	12491	0.25
MET cut	10327	0.21	10614	0.21	11156	0.22	10745	0.21
Jet cut	4403	0.09	5254	0.11	5477	0.11	5252	0.11

Table 1: Event Pass Rates across different polarization states.

1.4 Kinematic Distributions

1.4.1 Definition and Physical Meaning

1. Transverse Mass m_T

In the lab frame, we only know the transverse components of the neutrinos (the Missing Transverse Energy, or MET). Because the longitudinal momentum (p_z) of the neutrinos is unknown, we cannot calculate a true invariant mass. Instead, we project everything onto the 2D transverse plane ($x - y$) to calculate the Cluster Transverse Mass (m_T).

The Mathematics: To calculate the m_T of the WW system, we treat the two visible leptons as a single combined “cluster” ($\ell\ell$) and the MET as the invisible component. Dilepton Transverse Energy ($E_T^{\ell\ell}$): This combines the transverse momentum and the invariant mass of the dilepton system.

$$E_T^{\ell\ell} = \sqrt{|\vec{p}_T^{\ell\ell}|^2 + M_{\ell\ell}^2}, \quad (1)$$

Missing Transverse Energy (E_T^{miss}): This is simply the magnitude of the MET vector.

$$E_T^{\text{miss}} = |\vec{p}_T^{\text{miss}}|, \quad (2)$$

WW Transverse Mass (m_T^{WW}): We subtract the squared vector sum of the transverse momenta from the squared scalar sum of the transverse energies.

$$m_T^{WW} = \sqrt{(E_T^{\ell\ell} + E_T^{\text{miss}})^2 - |\vec{p}_T^{\ell\ell} + \vec{p}_T^{\text{miss}}|^2}. \quad (3)$$

2. Difference in Azimuthal Angle $\Delta\phi_{jj}$

$$\phi = \tan^{-1} \left(\frac{p_y}{p_x} \right) \rightarrow \Delta\phi_{jj} = |\phi_{j1} - \phi_{j2}|, \quad (4)$$

Measures the angle between the two tagging jets in the transverse plane.

3. Difference in Pseudorapidity $\Delta\eta_{\ell\ell}$

$$\eta = \frac{1}{2} \ln \left(\frac{|p| + p_z}{|p| - p_z} \right) \rightarrow \Delta\eta_{\ell\ell} = |\eta_{\ell1} - \eta_{\ell2}|, \quad (5)$$

Measures how far apart the two charged leptons are along the longitudinal axis.

1.4.2 Results




Figure 1: Distribution in the signal region for the difference in pseudorapidity between leading and subleading leptons, the transverse mass, and the difference in azimuthal angle between leading and subleading jets.

For the bottom panel, the ratio is calculated by comparing the normalized distribution of a polarized sample (LL, LT, or TT) against the normalized distribution of the unpolarized (total) sample. The ratio is derived for the plotting:

Before calculating the ratio, the code ensures both the polarized and unpolarized samples are on the same scale by normalizing them. Instead of using the raw number of events, it treats each event as a fraction of the total sample size:

$$w_i = \frac{1}{N_{\text{events}}}, \quad (6)$$

where N_{events} is the number of events that passed all cuts for that specific sample.

For each variable, the data is divided into k bins. Let $H_p(b)$ be the normalized count for a polarized sample in bin b , and $H_u(b)$ be the normalized count for the unpolarized sample:

$$H(b) = \sum_{x_i \in \text{bin } b} w_i, \quad (7)$$

The ratio plotted in the bottom panel represents how much a specific polarization deviates from the average unpolarized behavior in a specific kinematic region:

$$R(b) = \frac{\text{Norm(Polarized)}_b}{\text{Norm(Unpolarized)}_b} = \frac{H_p(b)}{H_u(b)}, \quad (8)$$

- If $R(b) > 1$: The polarized state is more likely to produce events in that specific parameter range than the total population.
- If $R(b) < 1$: The polarized state is suppressed in that range.

2 March 1st

2.1 Additional

To address the low selection efficiency observed in the initial MadGraph results¹, I implemented additional phase-space cuts directly in the run card and increased the number of events generated. This refinement ensures a higher yield of relevant events and optimizes computational resources.

I adjusted the following parameters in the run_card. Including restrictions for the transverse momentum of leptons and jets, missing transverse mass, pseudorapidity for leptons and jets, invariant mass for lepton pairs and jet pairs, and the transverse momentum for leading and subleading jets.

```

27.0 = ptl
35.0 = ptj
20.0 = misset
2.5 = etal
4.5 = etaj
20.0 = mml1
500.0 = mmjj
65.0 = ptj1min
35.0 = ptj2min

```

Event Pass Rates Cut	LL		LT		TT		Unpol	
	Count	Rate	Count	Rate	Count	Rate	Count	Rate
Total	100000	1.00	100000	1.00	100000	1.00	100000	1.00
Lepton cut	26793	0.27	27563	0.28	30021	0.30	28388	0.28
MET cut	23816	0.24	25036	0.25	27755	0.28	25979	0.26
Jet cut	19002	0.19	20602	0.21	22352	0.22	21116	0.21
Cross Sections	LL (pb)		LT (pb)		TT (pb)		Unpol (pb)	
Total	0.000505		0.002843		0.004288		0.007653	
Lepton cut	0.000135		0.000784		0.001287		0.002173	
MET cut	0.000120		0.000712		0.001190		0.001988	
Jet cut	0.000096		0.000586		0.000959		0.001616	

Table 2: Event pass rates and cross sections after pre-cut in MadGraph for LL, LT, TT, and Unpolarized samples.

In table 2, we can see that applying pre-cutting in MadGraph improves the event throughput, which increases the efficiency of data utilization. Furthermore, the change in cross-section after each cut is exactly the ratio times the original cross-section.

2.2 Transformation from pp -cmf to WW -cmf

Since the kinematic distribution in the paper[1] is under WW -cmf, we should consider to transform our original generated events in MadGraph from pp -cmf to WW -cmf.

2.2.1 Work Flow

To transform the cmf of the events, we have to first extract the 4-vector from the original events including leptons, neutrinos, and W bosons, where $W = l + \nu_l$. Then, we calculate the WW system kinematic by:

$$P_{WW} = P_{W+} + P_{W-}, \quad (9)$$

where P is the 4-vector.

To move into the WW -cmf, we must apply a Lorentz transformation to boost our target particles in the opposite direction of the WW system's motion. Therefore, we boost using $-\vec{\beta}$.

$$\vec{\beta}_{WW} = \frac{\vec{p}_{WW}}{E_{WW}}, \quad (10)$$

where $\vec{p}_{WW}$ equals (p_x, p_y, p_z) .

Since we only know $(E_\ell, p_{\ell,x}, p_{\ell,y}, p_{\ell,z})$ and $(p_{\nu,x}, p_{\nu,y})$, we have to calculate the $p_{\nu,z}$ and E_ν . By using the W -mass constraint, which assumes the W boson is perfectly on-shell,

$$(p_\ell + p_\nu)^2 = m_W^2, \quad (11)$$

where $m_W \approx 80.4$ GeV.

By expanding equation 11 using the known lepton kinematics and the measured transverse missing energy, we can end up with a standard quadratic equation for the neutrino's longitudinal momentum and then figure out the $p_{\nu,z}$ and E_ν . However, the transverse part of the momentum is non zero, therefore, the boost is not a purely longitudinal boost, but a 3D boost.

2.2.2 Expectations after the Transformation

- $\Delta\eta_{\ell\ell}$:

$$\Delta\eta' \neq \Delta\eta, \quad (12)$$

The difference in pseudorapidity should change on an event-by-event basis. Because a 3D boost mixes the momentum components, there is no single constant shift applied to the leptons. The $\Delta\eta$ distribution in the boosted frame should look smeared or broaden compared to the lab frame.

- m_T :

$$m_{T,pp-cmf} \rightarrow M_{WW,WW-cmf}, \quad (13)$$

Since we calculated the longitudinal momentum for neutrino, we can then figure out the invariant mass, instead of the transverse mass. While the total momentum in WW -cmf is zero, the invariant mass of the system is simply the total energy of all decay products:

$$M_{WW} = E_{\ell 1}^* + E_{\nu 1}^* + E_{\ell 2}^* + E_{\nu 1}^*. \quad (14)$$

- $\Delta\phi_{jj}$:

When analyzing the Vector Boson Scattering (VBS), the leading and subleading tagging jets have a very specific kinematic signature: they are highly forward. Which means they have massive total energies and longitudinal momenta. The $\gamma E\beta_T$ term becomes massive and so, the $\Delta\phi_{jj}$ distribution is will be heavily distorted.

$$\vec{p}'_{Ti} = \vec{p}_{Ti} + \left[\frac{\gamma - 1}{\beta^2} (\vec{\beta} \cdot \vec{p}_i) - \gamma E_i \right] \vec{\beta}_T. \quad (15)$$

2.2.3 Results




Figure 2: Distribution in the signal region after pre-cut and transformation to the WW -cmf in MadGraph for the difference in pseudorapidity between leading and subleading leptons, the transverse mass, and the difference in azimuthal angle between leading and subleading jets.

- **Peak values for $\Delta\eta_{\ell\ell}$ decreases:** The decrease in the TT peak for $\Delta\eta_{\ell\ell}$ in the WW -cmf occurs because boosting into the center-of-mass frame removes the global longitudinal “push” of the lab frame, unfolding the system and allowing the naturally broader angular distributions of transversely polarized bosons—which follow a $(1 \pm \cos^2 \theta^*)$ distribution—to spread out and lower the normalized peak. Furthermore, because TT leptons are emitted more longitudinally, they are highly sensitive to “smearing” and potential under-boosting caused by the algebraic approximation of p_z , which further flattens the distribution compared to the LL state.
- **M_{WW} goes to zero below ≈ 160 GeV:** The M_{WW} values drop to zero below approximately 160 GeV primarily due to the on-shell threshold; since we force a W -mass constraint of 80.385 GeV for each reconstructed boson, the total invariant mass cannot physically fall below the sum of those two masses (~ 161 GeV). Additionally, the selection criteria—specifically the high p_T requirements for leptons and the $m_{jj} > 500$ GeV VBS topology cut—effectively filter out low-energy phase space where a lower M_{WW} might otherwise occur.
- **$\Delta\phi_{jj}$ goes to zero:** In pp -cmf, the $\Delta\phi_{jj}$ distribution is governed by the hard scatter and recoil dynamics. But in WW -cmf, the distribution is heavily destroyed by the massive Energy term through the 3D boost. The jets are forced to point in similar transverse direction, making them highly collinear in ϕ .

3 March 8th

3.1 WW -cmf in MadGraph

```
generate p p > w+{0} w+{0} j j QCD=0, w+ > l+ vl  
add process p p > w-{0} w-{0} j j QCD=0, w- > l- vl~  
output sample/script/LL  
launch sample/script/LL
```

```
shower = Pythia8  
detector = Delphes  
analysis= OFF  
madspin= OFF
```

```
set run_card nevents 100000  
set run_card ebeam1 6500  
set run_card ebeam2 6500
```

```
set run_card me_frame 3,4,5,6
```

```
set run_card cut_decays True  
set run_card ptj 34.0  
set run_card ptl 26.0  
set run_card miset 29.0
```

```
set run_card etaj 4.6  
set run_card etal 2.6
```

```
set run_card mmjj 450
```

```
set run_card ptj1min 64  
set run_card ptj2min 34
```




Figure 3: Distribution in the signal region after pre-cut in the WW -cmf in MadGraph for the difference in pseudorapidity, transverse mass, and difference in azimuthal angle.

	pp -cmf (fb)				WW -cmf (fb)			
	LL	LT	TT	Total	LL	LT	TT	Total
Total	0.505	2.843	4.288	7.653	0.396	1.263	2.334	3.913
Lepton cut	0.135	0.784	1.287	2.173	0.201	0.650	1.284	2.082
MET cut	0.120	0.712	1.190	1.988	0.186	0.611	1.218	1.966
Jet / VBS cut	0.096	0.586	0.959	1.616	0.148	0.484	0.960	1.547
Jet cut (Thesis)	0.177	1.030	1.718	2.925	0.277	0.874	1.744	2.896
Fraction	5.9%	35.7%	58.4%	100%	9.3%	30.4%	60.3%	100%
Fraction (Thesis)	6.1%	35.2%	58.7%	100%	9.6%	30.2%	60.2%	100%

Table 3: Comparison of cutflows, cross-sections, and polarization fractions for pp -cmf and WW -cmf frames. Results from local MadGraph generations are compared against the thesis references. All cross-sections are expressed in fb.

Table 3 compares the cutflow and polarization results of the local MadGraph generation against the reference thesis. While the absolute cross-sections in the signal region (Jet/VBS cut) are roughly 45% lower than the thesis values, the polarization fractions show excellent agreement. Specifically, the LL, LT, and TT fractions deviate by less than 1% from the reference in both the pp -cmf and WW -cmf frames. This confirms that despite the normalization difference, the underlying angular distributions and physics of the polarized states are correctly modeled. Figure 3 also shows the underlying kinematic distributions are similar to the results in the thesis.

3.2 Sherpa3

In Sherpa3, instead of generating an event that is 100 % purely "LL" or 100 % "TT", Sherpa generates a physical event (a specific arrangement of outgoing jets and leptons) and then calculates a weight for each polarization state for that exact same event.

Also, we must separate the W^+W^+jj and W^-W^-jj processes when using the Pol_Cross_Section feature because of how the generator strictly handles memory allocation for event weights.

3.2.1 Sample Preparation

For W^+ :

```
EVENTS: 150000          #number of events to generate

BEAMS: [2212, 2212]     #incoming colliding particles; 2212 for a proton
BEAM_ENERGIES: 6500     #incoming particle energy for each

#specifies the internal structure model of the proton to be used during the collision
PDF_LIBRARY: LHAPDFSherpa    #tell Shepra to interface with the library
PDF_SET: PDF4LHC21_40_pdfas  #highly accurate PDF set

#93: any light quark, anti-quark, or gluon
#24: w+ boson
PROCESSES:
#2 incoming partons scatter to produce 2 same sign bosons and 2 outgoing partons
- 93 93 -> 24 24 93 93:
#forbidding any strong-force interactions in the hard matrix element
    Order: {QCD: 0, EW: 4}

SELECTORS:
#invariant mass of the 2 outgoing partons must be between 100-13,000 GeV
- [Mass, 93, 93, 100.0, 13000.0]

PARTICLE_DATA:
#treat w+ boson as strictly on-shell during the initial matrix element calculation
24:
    Width: 0.0
#turns on the decay module for heavy particles
HARD_DECAYS:
    Enabled: true
#crucial!!
#ensures the spin-correlations and polarization of w boson are calculated and passed
down to their decay products
    Pol_Cross_Section:
        Enabled: true
        Reference_System: COM      #COM:ww-cmf; Lab: pp-cmf
#disables specific decay routes for the w boson
#status : 0 meaning turns a decay channel off.
    Channels:
```

```

    '24,2,-1': {Status: 0}          #up+anti-down quark (hardronic decay)
    '24,4,-3': {Status: 0}          #charm+anti-strange quark (hardronic decay)
    '24,16,-15': {Status: 0}        #tau neutrino+anti-tau lepton (hardronic decay)
#dictates how and where the final simulated events are saved
EVENT_OUTPUT: HepMC3_GenEvent[Sherpa_Wp]

```

For W^- :

```
EVENTS: 150000
```

```
BEAMS: [2212, 2212]
```

```
BEAM_ENERGIES: 6500
```

```
PDF_LIBRARY: LHAPDFSherpa
```

```
PDF_SET: PDF4LHC21_40_pdfas
```

```
PROCESSES:
```

```

- 93 93 -> -24 -24 93 93:
  Order: {QCD: 0, EW: 4}

```

```
SELECTORS:
```

```
- [Mass, 93, 93, 100.0, 13000.0]
```

```
PARTICLE_DATA:
```

```

-24:
  Width: 0.0

```

```
HARD_DECAYS:
```

```
Enabled: true
```

```
Pol_Cross_Section:
```

```
Enabled: true
```

```
Reference_System: COM
```

```
Channels:
```

```

'-24,-2,1': {Status: 0}
'-24,-4,3': {Status: 0}
'-24,-16,15': {Status: 0}

```

```
EVENT_OUTPUT: HepMC3_GenEvent[Sherpa_Wm]
```

	Cross Sections (fb)				Event Pass Rates	
	LL	LT	TT	Unpol	Raw Events	Fraction
Total	1.239	3.781	6.874	11.915	299,998	1.0000
Lepton cut	0.577	1.710	3.230	5.479	142,376	0.4746
MET cut	0.477	1.473	2.767	4.658	120,052	0.4002
Jet cut	0.239	0.753	1.398	2.349	49,259	0.1642

Table 4: Cross sections for different polarization states and corresponding event pass rates at each selection cut stage.

According to table 4, the event pass rate for Sherpa simulation is much smaller than MadGraph due to less cuts in the event generating step. Perhaps we can explore whether there are other cuts can be added to Sherpa to enhance the event usage.

	Sherpa pp -cmf (fb)				Sherpa WW -cmf (fb)			
	LL	LT	TT	Total	LL	LT	TT	Total
Total	1.060	4.787	6.083	11.915	1.254	3.816	6.873	11.915
n_l cut	0.472	2.180	2.906	5.456	0.575	1.714	3.214	5.456
MET cut	0.381	1.845	2.516	4.635	0.469	1.470	2.755	4.635
VBS cut	0.175	0.915	1.306	2.335	0.237	0.749	1.392	2.335
VBS cut (Thesis)	0.182	1.072	1.732	2.933	0.285	0.909	1.778	2.933
Fraction	7.3%	38.2%	54.5%	100%	10.0%	31.5%	58.5%	100%
Fraction (Thesis)	6.1%	35.9%	58.0%	100%	9.6%	30.6%	59.8%	100%

Table 5: Comparison of cutflows, cross-sections, and polarization fractions for pp -cmf and WW -cmf frames using Sherpa. Results from local generations are compared against the thesis Sherpa ($W^\pm W^\pm jj$) references.

Table 5 shows that the calculated cross-sections are in close agreement with the thesis references. Furthermore, the polarization fractions for the LL, LT, and TT states remain consistent with the reference values across both the pp -cmf and WW -cmf frames.

4 March 12

4.1 Sherpa3 Rerun

It was identified that the initial analysis in section 3.2 lacked a proper detector simulation stage; specifically, Delphes was not applied to the Sherpa-generated events. Furthermore, the standalone Python implementation for jet reconstruction was unreliable for this analysis. Consequently, the Sherpa output files were re-processed through Delphes to ensure a more robust reconstruction of jets within the signal region.

Table 6 summarizes the results following this detector simulation. Notably, the discrepancy persists, which aligns with the results from MadGraph: the cross-sections obtained after the three signal region cuts are approximately 50% lower than those reported in the reference thesis. Despite this difference in absolute magnitude, the relative polarization fractions (LL, LT, and TT) remain broadly consistent with the thesis values.

	Sherpa3 + Delphes pp -cmf (fb)				Sherpa3 + Delphes WW -cmf (fb)			
	LL	LT	TT	Total	LL	LT	TT	Total
Total	1.060	4.787	6.083	11.915	1.254	3.816	6.873	11.915
n_{ll} cut	0.265	1.245	1.740	3.202	0.332	0.985	1.905	3.202
MET cut	0.217	1.057	1.515	2.744	0.275	0.847	1.645	2.744
VBS cut	0.112	0.585	0.870	1.540	0.154	0.479	0.922	1.540
VBS cut (Thesis)	0.182	1.072	1.732	2.933	0.285	0.909	1.778	2.933
Fraction	7.1%	37.4%	55.5%	100%	9.9%	30.8%	59.3%	100%
Fraction (Thesis)	6.1%	35.9%	58.0%	100%	9.6%	30.6%	59.8%	100%

Table 6: Comparison of cutflows, cross-sections, and polarization fractions for pp -cmf and WW -cmf frames using Sherpa after Delphes detector simulation. Local generations are compared against the thesis Sherpa ($W^\pm W^\pm jj$) references.

4.2 PDF Selection

I was using the preset settings for event generation in MadGraph of PDF selection:

```
nn23lo1 = pdlabel
230000 = lhaid
```

But to compare the results with Sherpa, we need to use the same PDF. Therefore, I decided to use the following settings:

In Sherpa3:

```
PDF_SET: PDF4LHC21_40_pdfas
```

In MadGraph:

```
set pdlabel lhapdf
set lhaid 93300
```

And I found that the cross section for different PDF changed:

	LL (fb)		LT (fb)		TT (fb)	
	PDF	Nominal	PDF	Nominal	PDF	Nominal
Total	0.434	0.396	1.372	1.263	2.531	2.334
n_{ll} cut	0.221	0.201	0.702	0.649	1.376	1.284
MET cut	0.205	0.186	0.660	0.611	1.302	1.218
VBS cut	0.163	0.148	0.518	0.484	1.017	0.960

Table 7: Comparison of cross-sections for different polarization states showing the effect of the PDF selection. The “PDF” columns represent the new PDF choice, while “Nominal” represents the baseline generation. All values are in fb.

To ensure consistency between MadGraph and Sherpa3, the default NNPDF2.3 LO (lhaid 230000) was replaced with the modern PDF4LHC21 benchmark (lhaid 93300).

As shown in table 7, updating to the PDF4LHC21 set results in a systematic increase in cross sections across all polarization states. This shift is expected, as the modern PDF set incorporated more precise experimental data and updated theoretical constraints compared to the decade-old nominal set. This transition ensures that the local simulations align with current LHC standards and provide more reliable results.

4.3 Particle-level phase space

After I read the thesis in detail again, I found out that the thesis was comparing the cross section at particle-level, not the detector-level. However, the previous cross section for my simulation were calculated at the detector-level. Instead of rerunning the event generations, we can use the free Delphes GenJet branch. Despite Delphes is a detector simulator, its default configuration actually runs the FastJet clustering algorithm on the PYTHIA truth particles before it applies any detector smearing.

- Particle-Level jets: Must run a jet clustering algorithm on all stable final state particles (Status = 1). (anti- kt algorithm with $R = 0.4$).
- Particle-Level MET: This is calculated as the negative vector sum of the transverse momenta of all stable, weakly interacting particles.
- Particle-Level Leptons: These are stable electrons and muons straight from the PYTHIA output.

```
particles = "Particle.PT", "Particle.Eta", "Particle.Phi", "Particle.Mass"...
genjets = "GenJet.PT", "GenJet.Eta", "GenJet.Phi", "GenJet.Mass"
genmet = "GenMissingET.MET"
```

Sample	WW -cmf			pp -cmf		
	LL	LT+TL	TT	LL	LT+TL	TT
Thesis MadGraph	0.277 fb 9.58%	0.874 fb 30.18%	1.744 fb 60.24%	0.177 fb 6.07%	1.030 fb 35.20%	1.718 fb 58.73%
Local MadGraph	0.256 fb 9.59%	0.814 fb 30.56%	1.595 fb 59.85%	0.166 fb 6.19%	0.954 fb 35.60%	1.560 fb 58.22%
Thesis Sherpa $W^\pm W^\pm jj$	0.285 fb 9.60%	0.909 fb 30.59%	1.778 fb 59.81%	0.182 fb 6.09%	1.072 fb 35.91%	1.732 fb 58.00%
Local Sherpa3	0.252 fb 9.87%	0.780 fb 30.58%	1.519 fb 59.54%	0.192 fb 7.47%	0.977 fb 37.93%	1.406 fb 54.60%

Table 8: Comparison of cross-sections and polarization fractions for WW -cmf and pp -cmf frames. The reference thesis values (MadGraph and Sherpa) are compared against the local MadGraph generations and local Sherpa3 generations at the particle-level VBS cut.

Table 8 provides a comprehensive comparison between local simulation and the reference thesis for both pp -cmf and WW -cmf. The polarization fractions show similar results. While the absolute cross section in the local generations are slightly lower than the thesis reference, which may likely due to higher order corrections or PDF sets. This result also shows that the underlying physics modeling and symmetry factors in MadGrpah or Sherpa3 setups are correctly aligned with the thesis results.

5 3/20

5.1 Background events

Following are the ranked background events mentioned in [2]:

1. $W^\pm Z$ background (dominant)
2. Data-driven backgrounds (~ 18 % combined):
 - Non-prompt leptons: arises from leptons originating from hadron decays and jets that are misidentified as leptons. The source of these events is mainly W +jets and semi-leptonic $t\bar{t}$ processes.
 - Charge-flip background (electron charge misidentification): This background primarily comes from $Z \rightarrow e^+e^-$ decays. A high-energy electron emits a hard photon via bremsstrahlung, which then undergoes pair production into a high-energy positron and a soft electron. The soft electrons get lost in the detector, and the misidentified opposite-charged positron is measured instead, incorrectly passing the same-charge requirement.
3. QCD $W^\pm W^\pm jj$ background (~ 5.2 %)
4. Small prompt backgrounds (< 2 % combined)
5. Interference $W^\pm W^\pm jj$ –INT (~ 1.6 %)

5.1.1 $W^\pm Z jj$ background

- $pp \rightarrow W^\pm Z jj \rightarrow \ell^\pm \nu \ell^\pm \ell^\mp$ (including both pure electroweak $W^\pm Z$ –EW and QCD-induced $W^\pm Z$ –QCD scattering)
- Event generator:
 - $W^\pm Z$ –EW: Madgraph
 - $W^\pm Z$ –QCD: Sherpa2.2.12 + data-driven approach (will Shepra3 simulation be sufficient?) + control region restrictions
- CMF: evaluated in the WW -cmf and pp -cmf
- PDF set: NNPDF3.0_nnlo_as.0118

This background originates from the fully leptonic decay. It mimics the signal region criteria when one of the leptons from the Z boson decay fails baseline selection or falls completely outside the detector’s acceptance. This leaves exactly two same-charged leptons, two jets, and missing transverse momentum. It consists of both pure electroweak and QCD-induced scattering events.

Comprises the electroweak scattering $W^\pm Z$ -EW and the strong scattering $W^\pm Z$ -QCD. $W^\pm Z$ -EW can be simulated by MadGraph well, while $W^\pm Z$ -QCD is known to be poorly modeled by Monte Carlo simulations. Sample generated from Sherpa for $W^\pm Z$ -QCD still needs to be corrected using the same data-driven approach as in [2].

5.1.2 QCD $W^\pm W^\pm jj$ background

- $pp \rightarrow \ell^\pm \nu \ell^\pm \nu jj$ via strong interaction vertices at order $O(\alpha_{\text{EW}}^4 \alpha_S^2)$
- Event generator: MadGraph
- CMF: evaluated in the WW -cmf and pp -cmf
- PDF set: NNPDF3.0_nnlo_as_0118

This is an irreducible background that yields the exact same final state particles as the signal, but it is produced at the order $O(\alpha_{\text{EW}}^4 \alpha_S^2)$ involving strong QCD interaction vertices.

5.1.3 Interference background $W^\pm W^\pm jj$ -INT

- Quantum mechanical interference between the purely electroweak and QCD-induced $W^\pm W^\pm jj$ diagrams that share identical external quarks, calculated at order $O(\alpha_{\text{EW}}^5 \alpha_S)$

- Event generator: MadGraph and Sherpa

For MadGraph, the interference is calculated by:

$$N_{\text{int}} = N_{\text{unpol}} - \sum_{\text{pol}} N_{\text{pol}} \quad (16)$$

For Sherpa, the interference contribution is directly provided by an event weight as for the individual polarization states.

- CMF: evaluated in the WW -cmf and pp -cmf
- PDF set: modeled by MadGraph using NNPDF3.0

5.2 Control Region

5.2.1 $W^\pm Z$ control region

- Two signal leptons and an additional third signal lepton with $p_T > 15$ GeV
- In the case of two electrons they have to fulfill $|\eta| < 1.37$ and $|m_{ee} - m_Z| > 15$ GeV
- At least two leptons satisfy $m_{\ell\ell} \geq 20$ GeV
- $m_{\ell\ell\ell} \geq 106$ GeV
- $p_T^{\text{miss}} \geq 30$ GeV
- At least two signal jets with the highest $p_T > 65$ GeV and the second highest $p_T > 35$ GeV
- $m_{jj} \geq 200$ GeV
- $|\Delta y_{jj}| > 2$
- No jet is b -tagged using the DL1r tagger with the 85% efficiency working point

6 3/29

6.1 Signal events

Adjusting the PDF set for MadGraph into

```
set pdlabel lhpdf
set lhaid 260000
```

which sets directly the same PDF set used in the thesis (NNPDF30_nlo_as_0118)

The following table is the after the correct PDF set selection with the thesis (NNPDF30_nlo_as_0118). We can compare how different overlap removal affect the cross section. But it isn't clearly mentioned in the thesis how the overlap removal is applied.

MadGraph	with step 5	without step 5	corrected	thesis
LL	0.250	0.218	0.229	0.277
LT	0.794	0.662	0.699	0.874
TT	1.558	1.193	1.309	1.744

Table 9: Signal events generated from MadGraph with the PDF set: NNPDF30_nlo_as_0118 and with different overlap removal conditions.

where the column “corrected” is set by the rules mentioned in the thesis 6.3.4 Fiducial Signal Region. If the resulting jet overlaps with an electron within $\Delta R < 0.4$, the electron is removed for $p_{T,e}/p_{T,jet} < 0.5$. The jet is removed if the electron carries more than 50% of the transverse jet momentum.

And the step 5 is a action to remove jets with $R > 0.3$ of the signal leptons.

Sherpa3	with step 5	without step 5	corrected	thesis
LL	0.252	0.233	0.236	0.285
LT	0.780	0.700	0.717	0.909
TT	1.519	1.245	1.325	1.778

Table 10: Signal events generated from Sherpa3 with the PDF set: PDF4LHC21_40_pdfas and with different overlap removal conditions.

We can see from both table that different choice of the overlap removal affect the cross section very much. However, in both the thesis or in the references mentioned in the thesis didn't clearly denote what overlap removal is applied to the signal events. Therefore, the discrepancy between our simulation result and the result in the thesis might due to the choice of the overlap removal.

6.2 Background events

I found some tables and figures that we might be able to compare with the background events before training the DNN model. There are different kinds of cut mentioned in the thesis, which is summarized in the following figure: According to this figure 4, we can compare our simulation result with table G.2-5 in the thesis [3] which includes the number of signal and background events in SR, CR, and low m_{jj} region. Also, it includes both the pp -cmf and WW -cmf. Therefore, by the following equation,

$$N_{\text{expected}} = \sigma \times \mathcal{L} \times \epsilon \quad (17)$$

Requirement	SR	Low- m_{jj} CR	WZ CR
Leading and subleading lepton p_T		> 27 GeV	
Electron $ \eta $	< 2.47 (1.37 in ee), excluding $1.37 \leq \eta \leq 1.52$		
Muon $ \eta $		< 2.5	
Leading (subleading) jet p_T		> 65 (35) GeV	
Additional jet p_T		> 25 GeV	
Jet $ \eta $		< 4.5	
$m_{\ell\ell}$		> 20 GeV	
E_T^{miss}		> 30 GeV	
Charge misid. $Z \rightarrow ee$ veto	$ m_{ee} - m_Z > 15$ GeV		–
b -jet veto	$N_{b\text{-jet}} = 0, p_T^{b\text{-jet}} > 20$ GeV, $ \eta^{b\text{-jet}} < 2.5$		
$N_{\text{veto leptons}}$	$= 0$	$= 0$	$= 1, p_T > 15$ GeV
$m_{\ell\ell\ell}$	–	–	> 106 GeV
m_{jj}	> 500 GeV	$200 < m_{jj} < 500$ GeV	> 200 GeV
$ \Delta y_{jj} $		> 2	

Figure 4: Summary of the event selection for the signal region and control region.

- N_{expected} : number of expected events
- σ : the theoretical cross-section of the process
- $\mathcal{L}$: the integrated luminosity of the data (for the full ATLAS Run 2 dataset, this is 139 fb^{-1})
- ϵ : the efficiency and acceptance. (the event pass rate)

we might be able to validate our results before applying the DNN training.

7 4/10

7.1 Background simulation

7.1.1 Simulation code

We generate the following background events by MadGraph 3.7.0 [4] and all of the PDF sets for each background simulation are selected to be `lhaid 260000`, which is `NNPDF30_nlo_as_0118`. For $W^\pm W^\pm jj$ QCD

```
generate p p > l+ vl l+ vl j j QCD=2 QED=4
add process p p > l- vl~ l- vl~ j j QCD=2 QED=4
```

For $W^\pm Z$ QCD

```
generate p p > l+ l- l+ vl j j QCD=0
add process p p > l+ l- l- vl~ j j QCD=0
```

For $W^\pm Z$ EW

```
generate p p > l+ l- l+ vl j j QCD=2 QED=4
add process p p > l+ l- l- vl~ j j QCD=2 QED=4
```

7.1.2 Results

Table 11: Comparison of signal region cross sections (in fb) between simulation results, Stange Thesis [1] (Table G.2), and ATLAS paper [2] (Table 3). Expected yields from the thesis and paper have been converted to cross sections using the integrated luminosity mentioned in both paper of 139 fb^{-1} .

Process	MadGraph3.7.1 [fb]	Sherpa3 [fb]	Thesis [1][fb]	Paper [2][fb]
Signal Processes ($W^\pm W^\pm jj$ EW)				
$W_L W_L$ (LL)	0.132		0.145	–
$W_L W_T$ (LT)	0.426		0.453	–
$W_T W_T$ (TT)	0.847		0.922	–
sum	1.405	1.353	1.520	–
WW unpolarized	1.378		1.554	1.691
Background Processes				
WZ -EW	0.150	0.409	0.108	0.107
WW -QCD	0.093	0.138	0.175	0.173
WZ -QCD	0.375	0.252	0.554	0.597

The signal region cuts obey the summarized table 2 in paper [2], but adding the detail that lepton pairs should be in same sign. We can see from table 11 that for electroweak processes (both signal and background), MadGraph [4] simulated a great result which is consistence with the reference. But for QCD-induced results, MadGraph [4] does not generate results reasonably. Therefore, we will try to generate some background events from Sherpa[5] to examine the consistency of the result. We can also see that for events generated by Sherpa[5], QCD-induced WW background behaves better than MadGraph, while MadGraph generates better for all electroweak events. However, there is still some inconsistency for the WZ -QCD induced background.

Table 12: Expected yields in the signal region (SR). The ATLAS results are shown alongside Thesis values and our simulation estimates using different generator versions, assuming an integrated luminosity of 139 fb^{-1} .

Process	MG 3.7.0	Sherpa 3.0.3	ATLAS	Thesis 9.1	Thesis G.2
$W^\pm W^\pm jj$ EW	195.31	188.03	235	206.52	211.29
$W^\pm W^\pm jj$ QCD	12.95	19.12	24	24.05	24.27
$W^\pm Z jj$ EW	20.81	56.98	14.9	14.95	15.07
$W^\pm Z jj$ QCD	52.13	34.99	83	82.75	77.0

We can also compare the result by the expected yields in the signal region shown in table 12. Which shows the same trend that Sherpa generates better WW -QCD induced events than MadGraph, while MadGraph generates better electroweak processes ($WWjj$ and $WZjj$) than Sherpa.

However, both event generators exhibit low consistency for the $WZjj$ -QCD induced process.

8 4/18

I took the following steps to try to generate results consistent with Feng-Yang's:

- **Cross section calculation for Sherpa:** Instead of extracting the nominal cross section from the `hepmc` file, we summed the cross sections for the positively charged processes (e.g., W^+) and the negatively charged processes (e.g., W^-), and then multiplied the total by the branching ratio.
- **Delphes `run_card`:** I initially used the ATLAS card but found that it lacked the fat jet finding configuration inside the `run_card` script. Therefore, I switched to the default `run_card` originally provided with Delphes and altered the jet radius to 0.4 to align with the paper.
- **Narrow width approximation (NWA) instead of the full process:** Because there were issues generating the full process in Sherpa (e.g., excessive integration time), we switched to using the narrow width approximation (NWA), which was also used for generating the signal process in the paper. Additionally, we adjusted the events generated by MadGraph to use NWA instead of the full process.

Table 13: Expected yields in the signal region (SR). The ATLAS results are shown alongside Thesis values and our simulation estimates using different generator versions, assuming an integrated luminosity of 139 fb^{-1} .

Process	MG 3.7.0	Sherpa 3.0.3	ATLAS	Thesis 9.1	Thesis G.2
$W^\pm W^\pm jj$ EW	201.89	169.26	235	206.52	211.29
$W^\pm W^\pm jj$ QCD	15.20	17.59	24	24.05	24.27
$W^\pm Zjj$ EW	21.94	32.05	14.9	14.95	15.07
$W^\pm Zjj$ QCD	50.84	78.35	83	82.75	77.0

Comparing this table with last week's, we observe that for MadGraph, the WW processes are now closer to the reference values. For Sherpa, the expected yield for the WW processes has decreased, which might be related to the different Delphes `run_card` used. For the WZ processes, Sherpa demonstrates better consistency than last week's results.

To make further comparisons with the results from the thesis or paper, we need to determine the exact Delphes `run_card` that was used. As our latest results indicate, the choice of `run_card` significantly affects the cross section, as shown in the following table:

Process	Expected Events (139 fb ⁻¹)	
	delphes_card_default.tcl	delphes_card_ATLAS.tcl
WWjj-EW-NWA	201.89	212.46
WWjj-QCD-NWA	15.20	15.86
WZjj-EW-NWA	21.94	24.82
WZjj-QCD-NWA	50.84	53.26

Table 14: Comparison of expected event yields across four processes, evaluated using the ATLAS Delphes card versus the Default Delphes card.

9 4/26

9.1 Summary of DNN Training Input Variables

Table 15 summarizes the variables used as inputs for the neural network training. To improve the training process for the DNN algorithms, some variables are pre-scaled to achieve a more Gaussian-distributed input.

The η values of leptons and jets can be used to define the lepton Zeppenfeld variable:

$$Z_l^* = \left| \frac{\eta_l - \frac{1}{2}(\eta_{j1} + \eta_{j2})}{\eta_{j1} - \eta_{j2}} \right|. \quad (18)$$

This is a measure of how central a lepton is relative to the two tagging jets.

The definition of the massless early-projected transverse mass is:

$$m_{o1}^{ll,MET} = \sqrt{(p_T^{l1} + p_T^{l2} + p_T^{miss})^2 - |\vec{p}_T^{l1} + \vec{p}_T^{l2} + \vec{p}_T^{miss}|^2}, \quad (19)$$

This is an advanced kinematic approximation used to estimate system masses involving missing transverse momentum, assuming massless visible products.

And the mass-preserving transverse mass is defined by:

$$m_T^{ll,MET} = \sqrt{(E_T^{ll} + p_T^{miss})^2 - |\vec{p}_T^{l1} + \vec{p}_T^{l2} + \vec{p}_T^{miss}|^2}, \quad (20)$$

Or generally, it can be defined for object x , for x being the leading lepton, sub-leading lepton, or the de-lepton system:

$$m_T(x, E_T^{miss}) = \sqrt{(E_T^x + E_T^{miss})^2 - |\vec{p}_T^x + \vec{p}_T^{miss}|^2}, \quad (21)$$

where $E_T = \sqrt{m^2 + |p_T|^2}$.

9.2 Dataset separation

The training data is split into train validation, and test sets according to the event number in a ration of 3:1:1.

- The test set: $((\text{EventNumber} - i_{\text{fold}}) \bmod 5) = 0$

Kinematics	Description	Scaling
Low-level variables		
$p_T^{\ell_1}, p_T^{\ell_2}, p_T^{j_1}, p_T^{j_2}$	p_T of the leading/subleading lepton/jet	$\log_{10}(x)$
$\eta^{\ell_1}, \eta^{\ell_2}, \eta^{j_1}, \eta^{j_2}$	η of the leading/subleading lepton/jet	x
ℓ_1 type, ℓ_2 type	Flavor of the leading/subleading lepton	x
$\phi^{\ell_2} - \phi^{\ell_1}, \phi^{j_1} - \phi^{\ell_1}, \phi^{j_2} - \phi^{\ell_1}$	Redefined ϕ of the subleading lepton/ leading jet/ subleading jet	x
p_T^{miss}	Missing transverse momentum	$\log_{10}(x)$
$\phi(p_T^{\text{miss}}) - \phi^{\ell_1}$	Redefined ϕ of the missing transverse energy	x
High-level variables		
$Z_{\ell_1}^*, Z_{\ell_2}^*$	Zeppenfeld variable of the leading/subleading lepton	$\sqrt{x}$
$m_T^{\ell_1, \text{MET}}, m_T^{\ell_2, \text{MET}}$	Transverse mass of the respective lepton and p_T^{miss}	$\sqrt{x}$
$\Delta R_{\ell\ell}, \Delta R_{jj}$	ΔR separation between the two leptons / two jets	x
$\Delta\eta_{\ell\ell}$	Pseudorapidity separation between the two leptons	$\sqrt{x}$
$m_{\ell\ell}, m_{jj}$	Invariant mass of the dilepton / dijet system	$\log_{10}(x)$
$p_T^{\ell\ell}$	Transverse momentum of the dilepton system	$\sqrt{x}$
$m_T^{\ell\ell, \text{MET}}$	Transverse mass of the dilepton system	$\sqrt{x}$
$m_{o1}^{\ell\ell, \text{MET}}$	Early-projected massless invariant mass	$\sqrt{x}$
Δy_{jj}	Rapidity separation between the two jets	x
$\Delta\phi_{jj}$	Azimuthal separation between the two jets	x
$(p_T^{\ell_1} \cdot p_T^{\ell_2}) / (p_T^{j_1} \cdot p_T^{j_2})$	p_T ratio of the lepton and jet systems	$\log_{10}(x + 0.02)$
$\min(\Delta R_{\ell_i, j_k})$	Minimal ΔR separation between any lepton and jet	x

Table 15: Summary of DNN input variables and their pre-scaling functions.

- The validation set: $((\text{EventNumber} - i_{\text{fold}} + 1) \bmod 5) = 0$
- The training set: the remaining events

The index i_{fold} defines the different splits used for cross-validation in the hyperparameter optimization.

9.3 Training process

The training processes are divided into two parts: First, an inclusive network designed to separate the electroweak (EW) signal from all backgrounds. Second, the polarization-specific networks trained exclusively on the EW samples to distinguish specific polarization states. In details, it is trained to separate longitudinal single-boson polarization (LL and LT v.s. TT) or double-boson polarization (LL v.s. LT and TT) from the remaining transverse polarization states.

10 5/1

10.1 5-Fold Method

The k -fold method is a data-splitting and cross-validation strategy used to reliably evaluate neural network performance and prevent biases when making predictions.

10.1.1 Data Categorization

The dataset is divided into three distinct and independent categories, using the i_{Fold} index:

- Test dataset: 20% with $((\text{EventNumber} - i_{\text{Fold}}) \bmod 5) = 0$. Kept completely hidden from the training and optimization process to test the final model.
- Validation dataset: 20% with $((\text{EventNumber} - i_{\text{Fold}} + 1) \bmod 5) = 0$. Used to evaluate the network's performance and tune hyperparameter during training.
- Training dataset: 60%, which is the remaining events that are not test or validation datasets. Used to adapt and optimize the network's trainable parameters.

where the index $i_{\text{Fold}} \in [0, 1, 2, 3, 4]$ and independent of the physical properties of the events.

10.1.2 Cross-Validation and Application

- Training Multiple Networks: The process iteratively trains five separate networks. Each network uses a different overlapping combination of the training folds and evaluates against non-overlapping validation and test sets.
- Evaluating Performance: Training across different independent validation datasets allows the author to calculate a mean network performance and estimate its statistical uncertainty.
- Preventing Mismodeling Bias: Neural networks naturally perform better on the data they were trained on than on unseen data. If predictions evaluated on training data were compared against measured data, it would introduce a systematic bias.
- Final Unbiased Predictions: To guarantee unbiased predictions across the entire dataset, each event's final prediction is generated exclusively by the specific DNN that held that event in its unseen test dataset.

10.1.3 Explanation of Proper Nouns

- **Unseen data** in the context of machine learning and the k -fold method refers specifically to the test dataset, which is used to provide an objective test of how well the final model can generalize to new, real-world examples. It is data that is kept completely hidden from the neural network while it is learning and optimizing its parameters.

$$s_j = \sum_i x_i w_{ij} + \theta_j \quad (22)$$

- **Weights** determine the strength and importance of the connection between two neurons. Mathematically, a weight is a multiplier. The output of a previous neuron (x_i) is multiplied by its assigned weight (w_{ij}) before it enters the next neuron.
- **Bias** (θ_j) is an additional parameter added to the neuron, independent of the input data. Mathematically, bias acts as an offset. By adding a bias, the network shifts the activation function left or right, determining how high the sum of the weighted inputs needs to be before the neuron activates.

10.2 Network Architecture

Based on section 9.2 in the reference thesis [3], the neural networks used are exclusively fully-connected feed-forward neural networks. The architecture is structured as follows:

- **Normalization Layer (Input):** Applies a scaling function to the input variables, normalizing their distributions to have a mean of 0 and a variance of 1. This ensures all inputs have similar scales without losing their kinematic phase space information.
- **Repeating Hidden Sequence:**

1. **Dense Layer:** A standard fully-connected layer.
2. **Batch Normalization Layer:** Used to normalize the output from the preceding dense layer.
3. **Activation Layer:** Applies the activation function, specifically the Swish function defined as:

$$f_{\text{Swish}}(x) = \frac{x}{1 + e^{-x}} \quad (23)$$

For a binary classification the Sigmoid function:

$$f_{\text{Sigmoid}}(x) = \frac{1}{1 + e^{-\beta x}} \quad (24)$$

4. **Dropout Layer:** Introduced for regularization to prevent overtraining.
- **Output Layer:** The repeating sequence concludes with a final dense layer, which computes the ultimate output score of the network.

References

- [1] ATLAS Collaboration. “Evidence for longitudinally polarized W bosons in the electroweak production of same-sign W boson pairs in association with two jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”. In: *Phys. Rev. Lett.* 135 (2025), p. 111802. DOI: 10.1103/bpln-ccq1.
- [2] The ATLAS Collaboration. “Measurement and interpretation of same-sign W boson pair production in association with two jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”. In: *JHEP* 04 (2024), p. 026. DOI: 10.1007/JHEP04(2024)026. arXiv: 2312.00420 [hep-ex].
- [3] Max V. Stange. “Longitudinal $W^\pm$ Boson Polarization in Same-Charged $W^\pm W^\pm$ Scattering in the ATLAS Experiment”. Doctoral thesis. Technische Universität Dresden, Jan. 2025.
- [4] J. Alwall et al. “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”. In: *JHEP* 07 (2014), p. 079. DOI: 10.1007/JHEP07(2014)079. arXiv: 1405.0301 [hep-ph].
- [5] Enrico Bothmann et al. “Event generation with Sherpa 3”. In: (Oct. 2024). arXiv: 2410.22148 [hep-ph].