

The Lax-Wendroff Scheme

João Bettencourt

February 2023

The Lax-Wendroff scheme starts from the linear 1-d advection equation:

$$\frac{\partial u}{\partial t} = -c \frac{\partial u}{\partial x}, \quad (1)$$

and uses the Taylor series expansion of $u(x, t + \Delta t) = u(x_m, t^{n+1}) = u_m^{n+1}$:

$$u_m^{n+1} = u_m^n + \Delta t \frac{\partial u}{\partial t} + \frac{\Delta t^2}{2} \frac{\partial^2 u}{\partial t^2} + O(\Delta t^3). \quad (2)$$

If we replace the time derivatives in (2) by the spatial derivative from (1), we get¹:

$$u_m^{n+1} = u_m^n + \Delta t \left(-c \frac{\partial u}{\partial x}\right) + \frac{\Delta t^2}{2} c^2 \frac{\partial^2 u}{\partial x^2} + O(\Delta t^3). \quad (3)$$

Now we can approximate the spatial derivatives with centred differences to obtain the Lax-Wendroff scheme:

$$u_m^{n+1} = u_m^n - c \frac{\Delta t}{2\Delta x} (u_{m+1}^n - u_{m-1}^n) + c^2 \frac{\Delta t^2}{\Delta x^2} (u_{m+1}^n - 2u_m^n + u_{m-1}^n), \quad (4)$$

where we recognize that the last term in (4) is a dissipative term added to (1) to control what would otherwise be an unstable Euler method. To find out the order of the Lax-Wendroff scheme, we can return to (3) and rewrite it as:

$$\frac{u_m^{n+1} - u_m^n}{\Delta t} = -c \frac{\partial u}{\partial x} + \frac{\Delta t}{2} c^2 \frac{\partial^2 u}{\partial x^2} + O(\Delta t^2),$$

where we then replace the spatial derivatives with their second-order approximations:

$$\begin{aligned} \frac{u_m^{n+1} - u_m^n}{\Delta t} = & -c \left[\frac{u_{m+1}^n - u_{m-1}^n}{2\Delta x} + O(\Delta x^2) \right] + \\ & + \frac{\Delta t}{2} c^2 \left[\frac{u_{m+1}^n - 2u_m^n + u_{m-1}^n}{\Delta x^2} + O(\Delta x^2) \right] + O(\Delta t^2), \end{aligned}$$

to find that the scheme is second order in space as in time.

¹Note that from (1) we have: $\frac{\partial^2 u}{\partial t^2} = -c \frac{\partial}{\partial t} \left(\frac{\partial u}{\partial x} \right) = -c \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial t} \right) = -c \frac{\partial}{\partial x} \left(-c \frac{\partial u}{\partial x} \right) = c^2 \frac{\partial^2 u}{\partial x^2}$