

Optimized Schwarz Method for the Linearized KdV equation and for coupling a NSW-E-Serre model

J. Caldas ¹ R. Cienfuegos ³ **J. Galaz** ³ A. Rousseau ²

¹MERIC, Chile

²Inria and Inria Chile

³Department of Hydraulic and Environmental Engineering, Pontifical Catholic University of Chile

WTE2016

RL1.6 on Resource Assessment and Site Characterization

Advanced modelling for marine energy

Model 1



Model n-1



Model n



RL1.6 on Resource Assessment and Site Characterization

Advanced modelling for marine energy

Model 1



Model n-1



Model n




Ocean propagation
Spectral models
Low resolution
Linear phenomena

Shore interaction
Time domain models
High resolution
Strong non-linear
phenomena


RL1.6 on Resource Assessment and Site Characterization

Advanced modelling for marine energy



RL1.6 on Resource Assessment and Site Characterization

Advanced modelling for marine energy



Question

Can we make these different models **communicate** with each other?

Truncated domain error: A tsunami simulation

Nonlinear long wave model (GeoClaw from the U. Washington)

Challenges

- Long simulations are required for statistics estimates
- Transport of the numerical error from deep to shallow water: shoaling

Truncated domain error: A tsunami simulation

Nonlinear long wave model (GeoClaw from the U. Washington)

Challenges

- Long simulations are required for statistics estimates
- Transport of the numerical error from deep to shallow water: shoaling

Truncated domain error: A tsunami simulation

Nonlinear long wave model (GeoClaw from the U. Washington)

Challenges

- Long simulations are required for statistics estimates
- Transport of the numerical error from deep to shallow water: shoaling

First results

- Develop a self coupling methodology: Domain Decomposition Methods (DDM)

First results

- Develop a self coupling methodology: Domain Decomposition Methods (DDM)
- Study the most fundamental equation for **non-linear** and **dispersive** waves: The KdV equation

$$u_t + u_x + uu_x + u_{xxx} = 0$$

First results

- Develop a self coupling methodology: Domain Decomposition Methods (DDM)
- Study the most fundamental equation for **non-linear** and **dispersive** waves: The KdV equation

$$u_t + u_x + uu_x + u_{xxx} = 0$$

- Simplify even more: keep only the **dispersive** part

$$u_t + u_{xxx} = 0$$

Our work

First results

- Develop a self coupling methodology: Domain Decomposition Methods (DDM)
- Study the most fundamental equation for **non-linear** and **dispersive** waves: The KdV equation

$$u_t + u_x + uu_x + u_{xxx} = 0$$

- Simplify even more: keep only the **dispersive** part


$$u_t + u_{xxx} = 0$$

Ongoing work


Develop a coupled model between the **Non Linear Shallow Water Equations** (NSWE) and the **Serre equations**

A key ingredient: *Transparent* Boundary Conditions (TBC)

- 1 Start with an infinite domain:




A key ingredient: *Transparent* Boundary Conditions (TBC)




- 1 Start with an infinite domain:
- 2 Seek for an exact explicit solution ... and fail (most of the time)

A key ingredient: *Transparent* Boundary Conditions (TBC)




- 1 Start with an infinite domain:
- 2 Seek for an exact explicit solution ... and fail (most of the time)
- 3 **Truncate** the domain:
approximate numerical solution

A key ingredient: *Transparent* Boundary Conditions (TBC)



- 1 Start with an infinite domain:
- 2 Seek for an exact explicit solution ... and fail (most of the time)
- 3 **Truncate** the domain:
approximate numerical solution
- 4 *Apply* Laplace transform to the equation at the boundaries of the **infinite domain** (unknown),
formulate candidate solutions

A key ingredient: *Transparent* Boundary Conditions (TBC)



- 1 Start with an infinite domain:
- 2 Seek for an exact explicit solution ... and fail (most of the time)
- 3 **Truncate** the domain:
approximate numerical solution
- 4 *Apply* Laplace transform to the equation at the boundaries of the **infinite domain** (unknown), formulate candidate solutions
- 5 *Un-apply* it and get new equations for the boundaries

Transparent Boundary Conditions (TBC's)

Exact TBC's (Besse et al, 2015)

Transparent Boundary Conditions (TBC's)

Exact TBC's (Besse et al, 2015)

- Solving these equations is also *very* hard (non-local operator)

$$\begin{cases} u(t, \text{left}) - \mathcal{L}^{-1} \left(\frac{\lambda_1(s)^2}{s} \right) * u_x(t, a) - \mathcal{L}^{-1} \left(\frac{\lambda_1(s)}{s} \right) * u_{xx}(t, a) = 0 \\ u(t, \text{right}) - \mathcal{L}^{-1} \left(\frac{1}{\lambda_1(s)^2} \right) * u_{xx}(t, b) = 0 \\ u_x(t, \text{right}) - \mathcal{L}^{-1} \left(\frac{1}{\lambda_1(s)} \right) * u_{xx}(t, b) = 0 \end{cases}$$

Transparent Boundary Conditions (TBC's)

Exact TBC's (Besse et al, 2015)

- Solving these equations is also *very* hard (non-local operator)

$$\begin{cases} u(t, \text{left}) - \mathcal{L}^{-1} \left(\frac{\lambda_1(s)^2}{s} \right) * u_x(t, a) - \mathcal{L}^{-1} \left(\frac{\lambda_1(s)}{s} \right) * u_{xx}(t, a) = 0 \\ u(t, \text{right}) - \mathcal{L}^{-1} \left(\frac{1}{\lambda_1(s)^2} \right) * u_{xx}(t, b) = 0 \\ u_x(t, \text{right}) - \mathcal{L}^{-1} \left(\frac{1}{\lambda_1(s)} \right) * u_{xx}(t, b) = 0 \end{cases}$$


- Not suited for numerical computations!

Result

Using the approximation $\lambda(s) \approx c$ we obtain the following approximate TBC's


$$\begin{cases} u(t, \text{left}) - cu_x(t, \text{left}) + c^2 u_{xx}(t, \text{left}) = 0 \\ u(t, \text{right}) - c^2 u_{xx}(t, \text{right}) = 0 \\ u_x(t, \text{right}) + cu_{xx}(t, \text{right}) = 0 \end{cases}$$

Using the TBC's for a DDM: The Additive Schwarz Method



DDM Methodology


Using the TBC's for a DDM: The Additive Schwarz Method



DDM Methodology

- Solve each problem separately, using the adjacent domain last information.


Using the TBC's for a DDM: The Additive Schwarz Method



DDM Methodology

- Solve each problem separately, using the adjacent domain last information.
- Iterate until convergence criteria.

Using the TBC's for a DDM: The Additive Schwarz Method




DDM Methodology

- Solve each problem separately, using the adjacent domain last information.
- Iterate until convergence criteria.
- How do we determine the c_L and c_R constants ?


DDM: Determination of optimal $c_L = c_R = c$

- For a fixed mesh discretization (in time and space), fixed interface location and different time-steps (initial condition)



DDM: Determination of optimal $c_L = c_R = c$

- For a fixed mesh discretization (in time and space), fixed interface location and different time-steps (initial condition)




Remark

- There exist **optimal values** for c (e.g. ≈ 4.5) where convergence is observed with (only) 5-7 iterations.

DDM: Determination of optimal $c_L = c_R = c$


- For a fixed mesh discretization (in time and space), fixed interface location and different time-steps (initial condition)



Remark

- There exist **optimal values** for c (e.g. ≈ 4.5) where convergence is observed with (only) 5-7 iterations.
- There is no observed dependency on the **initial condition**.


DDM: Dependency on the location of the interface




Remark

There is no observed dependency on the location of the interface

DDM: Dependency on the discretization size: Δx , Δt




(a) Fixed $\Delta x = \frac{12}{250}$




(b) Fixed $\Delta t = 0.02$

DDM: Dependency on the discretization size: Δx , Δt



(a) Fixed $\Delta x = \frac{12}{250}$




(b) Fixed $\Delta t = 0.02$


Result

There is a strong dependence on Δx and Δt .

DDM: Dependency on the discretization size: Δx , Δt



(a) Fixed $\Delta x = \frac{12}{250}$



(b) Fixed $\Delta t = 0.02$


Result

There is a strong dependence on Δx and Δt . Furthermore the regression


$$c_{opt}(\Delta t, \Delta x) = \kappa + \alpha(\Delta t)^{2/3} + \beta \frac{1}{\Delta x} + \gamma \frac{\Delta t^{2/3}}{\Delta x}$$

fits the numerical experiments with $R^2 = 0.9999894$

DDM: Dependency on the discretization size: Δx , Δt



(a) Fixed $\Delta x = \frac{12}{250}$



(b) Fixed $\Delta t = 0.02$

Result

There is a strong dependence on Δx and Δt . Furthermore the regression

$$c_{opt}(\Delta t, \Delta x) = \kappa + \alpha(\Delta t)^{2/3} + \beta \frac{1}{\Delta x} + \gamma \frac{\Delta t^{2/3}}{\Delta x}$$

fits the numerical experiments with $R^2 = 0.9999894$ for $\kappa = 0.0775$, $\alpha = -0.3353$, $\beta = -0.0012$, $\gamma = 2.7407$

A coupled shallow water (NSWE) - Serre model


The Serre equations

$$h_t + (hu)_x = 0$$


$$u_t + uu_x + gh_x - \frac{1}{3h} \left(h^3 (u_{xt} + uu_{xx} - (u_x)^2) \right)_x = 0$$

- h = the water column height
- u = depth averaged horizontal velocity (1D)

Coupling methodology



Coupling methodology



Numerical method

- Fractional splitting scheme for non-linear/dispersive terms (Bonneton et al, 2011).
- Well-balanced fourth order accurate Finite Volume NSWE model (Berthon and Marche, 2008)
- Fourth order Finite Difference model for the dispersive equation

Three domain coupling: NSWE-Serre-NSWE